



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, 16/18, A61K 38/17, G01N 33/68	A2	(11) International Publication Number: WO 00/31263 (43) International Publication Date: 2 June 2000 (02.06.00)
(21) International Application Number: PCT/US99/28013 (22) International Filing Date: 23 November 1999 (23.11.99) (30) Priority Data: 60/109,592 23 November 1998 (23.11.98) US 60/118,610 4 February 1999 (04.02.99) US 60/127,990 6 April 1999 (06.04.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 60/109,592 (CIP) Filed on 23 November 1998 (23.11.98) US 60/118,610 (CIP) Filed on 4 February 1999 (04.02.99) US 60/127,990 (CIP) Filed on 6 April 1999 (06.04.99) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive, #12, Mountain View,		CA 94040 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). YANG, Junming [CN/US]; 7136 Clarendon Street, San Jose, CA 95129 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: GTPASE ASSOCIATED PROTEINS (57) Abstract The invention provides human GTPase associated proteins (GTPAP) and polynucleotides which identify and encode GTPAP. The invention also provides expression vectors, host cells, antibodies, agonist, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of GTPAP.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

GTPASE ASSOCIATED PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of GTPase associated proteins and to the use of these sequences in the diagnosis, treatment, and prevention of cell proliferative, autoimmune/inflammatory, and immune system disorders.

5

BACKGROUND OF THE INVENTION

Guanine nucleotide binding proteins (GTP-binding proteins) participate in a wide range of regulatory functions in all eukaryotic cells, including metabolism, cellular growth, differentiation, signal transduction, cytoskeletal organization, and intracellular vesicle transport and secretion. In higher organisms they are involved in signaling that regulates such processes as the immune response (Aussel, C. et al (1988) J. Immunol. 140:215-220), apoptosis, differentiation, and cell proliferation including oncogenesis (Dhanasekaran, N. et al. (1998) Oncogene 17:1383-1394). Exchange of bound GDP for GTP followed by hydrolysis of GTP to GDP provides the energy that enables GTP-binding proteins to alter their conformation and interact with other cellular components. The superfamily of GTP-binding proteins consists of several families and may be grouped as translational factors, heterotrimeric GTP-binding proteins involved in transmembrane signaling processes (also called G-proteins), and low molecular weight GTP-binding proteins including the proto-oncogene Ras proteins and products of rab, rap, rho, rac, smg21, smg25, YPT, SEC4, and ARF genes, and tubulins (Kaziro, Y. et al. (1991) Ann. Rev. Biochem. 60:349-400). In all cases, the GTPase activity is regulated through interactions with other proteins.

GTP-binding proteins involved in protein biosynthesis include initiation factor 2 (IF-2), elongation factor 2 (EF-Tu), and elongation factor G (EF-G), observed in prokaryotes; and initiation factor 2 (eIF-2), elongation factor 1 α (EF-1 α) and elongation factor 2 (EF-2) observed in eukaryotes (Kaziro, supra). IF-2 promotes the GTP-dependent binding of the tRNA to the small subunit of the ribosome, the step that initiates protein translation. Similarly, elongation factors promote the binding of tRNA and GTP and the displacement of GDP after hydrolysis as protein biosynthesis proceeds.

Heterotrimeric GTP-binding proteins are composed of 3 subunits (α , β and γ) which, in their inactive conformation, associate as a trimer at the inner face of the plasma membrane. G_α binds GDP or GTP and contains the GTPase activity. The $\beta\gamma$ complex enhances binding of G_α to a receptor. G_γ is necessary for the folding and activity of G_β . (Neer, E.J. et al. (1994) Nature 371:297-300.) Multiple homologs of each subunit have been identified in mammalian tissues, and different combinations of subunits have specific functions and tissue specificities. (Spiegel, A.M. (1997) J.

Inher. Metab. Dis. 20:113-121.) G protein activity is triggered by seven-transmembrane cell surface receptors (G-protein coupled receptors) which respond to lipid analogs, amino acids and their derivatives, peptides, cytokines, and specialized stimuli such as light, taste, and odor. Activation of the receptor by its stimulus causes the replacement of the G protein-bound GDP with GTP. G α -GTP dissociates from the receptor/ $\beta\gamma$ complex and each of these separated components can interact with and regulate downstream effectors. The signaling stops when G α hydrolyzes its bound GTP to GDP and reassociates with the $\beta\gamma$ complex (Neer, supra).

The alpha subunits of heterotrimeric G proteins can be divided into four distinct classes. The α -s class is sensitive to ADP-ribosylation by pertussis toxin which uncouples the receptor:G-protein interaction. This uncoupling blocks signal transduction to receptors that decrease cAMP levels which normally regulate ion channels and activate phospholipases. The inhibitory α -I class is also susceptible to modification by pertussis toxin which prevents α -I from lowering cAMP levels. Two novel classes of α subunits refractory to pertussis toxin modification are α -q, which activates phospholipase C, and α -12, which has sequence homology with the *Drosophila* gene concertina and may contribute to the regulation of embryonic development (Simon, M.I. (1991) Science 252:802-808).

The mammalian G β and G γ subunits, each about 340 amino acids long, share more than 80% homology. The G β subunit (also called transducin) contains seven repeating units, each about 43 amino acids long. The activity of both subunits may be regulated by other proteins such as calmodulin and phosducin or the neural protein GAP 43 (D. Clapham and E. Neer, 1993, Nature 365:403-406). The β and γ subunits are tightly associated. The β subunit sequences are highly conserved between species, implying that they perform a fundamentally important role in the organization and function of G-protein linked systems (Van der Voorn L. (1992) Febs. Lett. 307 (2):131-134). They contain seven tandem repeats of the WD-repeat sequence motif, a motif found in many proteins with regulatory functions. WD-repeat proteins contain from four to eight copies of a loosely conserved repeat of approximately 40 amino acids which participates in protein-protein interactions. Mutations and variant expression of β transducin proteins are linked with various disorders. Mutations in LIS1, a subunit of the human platelet activating factor acetylhydrolase, cause Miller-Dieker lissencephaly. RACK1 binds activated protein kinase C, and RbAp48 binds retinoblastoma protein. CstF is required for polyadenylation of mammalian pre-mRNA in vitro and associates with subunits of cleavage-stimulating factor. Defects in the regulation of β -catenin contribute to the neoplastic transformation of human cells. The WD40 repeats of the human F-box protein β TrCP mediate binding to β -catenin, thus regulating the targeted degradation of β -catenin by

ubiquitin ligase (Neer, *supra*; Hart, M. et al (1999) *Curr. Biol.* 9:207-210). The γ subunit primary structures are more variable than those of the β subunits. They are often post-translationally modified by isoprenylation and carboxyl-methylation of a cysteine residue four amino acids from the C-terminus; this appears to be necessary for the interaction of the $\beta\gamma$ subunit with the membrane and with other GTP-binding proteins. The $\beta\gamma$ subunit has been shown to modulate the activity of isoforms of adenylyl cyclase, phospholipase C, and some ion channels. It is involved in receptor phosphorylation via specific kinases, and has been implicated in the p21ras-dependent activation of the MAP kinase cascade and the recognition of specific receptors by GTP-binding proteins. (Clapham and Neer, *supra*).

10 G-proteins interact with a variety of effectors including adenylyl cyclase (Clapham and Neer, *supra*). The signaling pathway mediated by cAMP is mitogenic in hormone-dependent endocrine tissues such as adrenal cortex, thyroid, ovary, pituitary, and testes. Cancers in these tissues have been related to a mutationally activated form of a $G\alpha$, known as the gsp (Gs protein) oncogene (Dhanasekaran, *supra*). Another effector is phosducin, a retinal phosphoprotein, which forms a specific complex with retinal $G\beta$ and $G\gamma$ ($G\beta\gamma$) and modulates the ability of $G\beta\gamma$ to interact with retinal $G\alpha$ (Clapham and Neer, *supra*).

Irregularities in the GTP-binding protein signaling cascade may result in abnormal activation of leukocytes and lymphocytes, leading to the tissue damage and destruction seen in many inflammatory and autoimmune diseases such as rheumatoid arthritis, biliary cirrhosis, hemolytic anemia, lupus erythematosus, and thyroiditis. Abnormal cell proliferation, including cyclic AMP stimulation of brain, thyroid, adrenal, and gonadal tissue proliferation is regulated by G proteins. Mutations in $G\alpha$ subunits have been found in growth-hormone-secreting pituitary somatotroph tumors, hyperfunctioning thyroid adenomas, and ovarian and adrenal neoplasms (Meij, J.T.A. (1996) *Mol. Cell. Biochem.* 157:31-38; Aussel, *supra*).

25 LMW GTP-binding proteins are GTPases which regulate cell growth, cell cycle control, protein secretion, and intracellular vesicle interaction. They consist of single polypeptides which, like the alpha subunit of the heterotrimeric GTP-binding proteins, are able to bind to and hydrolyze GTP, thus cycling between an inactive and an active state. LMW GTP-binding proteins respond to extracellular signals from receptors and activating proteins by transducing mitogenic signals involved in various cell functions. The binding and hydrolysis of GTP regulates the response of LMW GTP-binding proteins and acts as an energy source during this process (Bokoch, G. M. and Der, C. J. (1993) *FASEB J.* 7:750-759).

At least sixty members of the LMW GTP-binding protein superfamily have been identified

and are currently grouped into the ras, rho, arf, sar1, ran, and rab subfamilies. Activated ras genes were initially found in human cancers, and subsequent studies confirmed that ras function is critical in determining whether cells continue to grow or become differentiated. Ras1 and Ras2 proteins stimulate adenylate cyclase (Kaziro, *supra*), affecting a broad array of cellular processes. Stimulation of cell surface receptors activates Ras which, in turn, activates cytoplasmic kinases. These kinases translocate to the nucleus and activate key transcription factors that control gene expression and protein synthesis (Barbacid, M. (1987) *Ann. Rev Biochem.* 56:779-827, Treisman, R. (1994) *Curr. Opin. Genet. Dev.* 4:96-98). Other members of the LMW GTP-binding protein superfamily have roles in signal transduction that vary with the function of the activated genes and the locations of the GTP-binding proteins that initiate the activity. Rho GTP-binding proteins control signal transduction pathways that link growth factor receptors to actin polymerization, which is necessary for normal cellular growth and division. The rab, arf, and sar1 families of proteins control the translocation of vesicles to and from membranes for protein processing, localization, and secretion. Vesicle- and target- specific identifiers (v-SNAREs and t-SNAREs) bind to each other and dock the vesicle to the acceptor membrane. The budding process is regulated by the closely related ADP ribosylation factors (ARFs) and SAR proteins, while rab proteins allow assembly of SNARE complexes and may play a role in removal of defective complexes (J. Rothman and F. Wieland (1996) *Science* 272:227-234). Ran GTP-binding proteins are located in the nucleus of cells and have a key role in nuclear protein import, the control of DNA synthesis, and cell-cycle progression (Hall, A. (1990) *Science* 249:635-640; Barbacid, M. (1987) *Ann. Rev Biochem.* 56:779-827; Ktistakis, N. (1998) *BioEssays* 20:495-504; and Sasaki, T. and Takai, Y. (1998) *Biochem. Biophys. Res. Commun.* 245:641-645).

The cycling of LMW GTP-binding proteins between the GTP-bound active form and the GDP-bound inactive form is regulated by additional proteins. Guanosine nucleotide exchange factors (GEFs) increase the rate of nucleotide dissociation by several orders of magnitude, thus facilitating release of GDP and loading with GTP. The best characterized is the mammalian homologue of the *Drosophila* Son-of-Sevenless protein. Certain Ras-family proteins are also regulated by guanine nucleotide dissociation inhibitors (GDIs), which inhibit GDP dissociation. The intrinsic rate of GTP hydrolysis of the LMW GTP-binding proteins is typically very slow, but it can be stimulated by several orders of magnitude by GTPase-activating proteins (GAPs) (Geyer, M. and Wittinghofer, A. (1997) *Curr. Opin. Struct. Biol.* 7:786-792). Both GEF and GAP activity may be controlled in response to extracellular stimuli and modulated by accessory proteins such as RalBP1 and POB1. Mutant Ras-family proteins, which bind but can not hydrolyze GTP, are permanently activated, and cause cell proliferation or cancer, as do GEFs that inappropriately activate LMW GTP-binding proteins, such as the human oncogene NET1, a Rho-GEF (Drivas, G. T. et al. (1990) *Mol. Cell. Biol.* -

10:1793-1798; Alberts, A. S. and Treisman, R. (1998) EMBO J. 14:4075-4085).

A novel group of GTP-binding proteins is the GTP1/OBG family, which are found in species ranging from bacteria to yeast to humans. These proteins contain characteristic GTP-binding motifs and are similar to one another but do not show sequence homology to other GTP-binding proteins.

- 5 The exact functions of these proteins are as yet uncertain, but they have been shown to be important for regulation of cell differentiation and development (Okamoto, S. and Ochi, K. (1998). Mol. Microbiol 30:107-119; Sazaka, T. et al. (1992) Biochem. Biophys. Res. Commun. 189:363-370).

The discovery of new GTPase associated proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis,
10 prevention, and treatment of cell proliferative, autoimmune/inflammatory, and immune system disorders.

SUMMARY OF THE INVENTION

The invention features substantially purified polypeptides, GTPase associated proteins,
15 referred to collectively as "GTPAP" and individually as "GTPAP-1," "GTPAP-2," "GTPAP-3,"
"GTPAP-4," "GTPAP-5," "GTPAP-6," "GTPAP-7," "GTPAP-8," "GTPAP-9," "GTPAP-10,"
"GTPAP-11," "GTPAP-12," "GTPAP-13," "GTPAP-14," "GTPAP-15," "GTPAP-16," "GTPAP-17,"
"GTPAP-18," "GTPAP-19," "GTPAP-20," "GTPAP-21," "GTPAP-22," "GTPAP-23," "GTPAP-24,"
"GTPAP-25," "GTPAP-26," "GTPAP-27," "GTPAP-28," and "GTPAP-29." In one aspect, the
20 invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also includes a polypeptide comprising an amino acid sequence that differs by one or more conservative amino acid substitutions from an amino acid sequence selected from the group consisting of SEQ ID NO:1-29.

The invention further provides a substantially purified variant having at least 90% amino acid
25 identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide
30 encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof. The

invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

5 The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the sample. In one aspect, the method further comprises amplifying the polynucleotide prior to hybridization.

10 The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof. The invention also provides an isolated and
15 purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof.

The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group
20 consisting of SEQ ID NO:1-29. In another aspect, the expression vector is contained within a host cell.

The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing a polynucleotide of the invention under conditions suitable for the expression of the polypeptide; and (b) recovering the
25 polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide selected from
30 the group consisting of SEQ ID NO:1-29 and fragments thereof. The invention also provides a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a disorder associated with decreased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially

purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention also provides a method for treating or preventing a disorder associated with increased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding GTPAP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of GTPAP.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding GTPAP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze GTPAP, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same

meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing
5 the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

“GTPAP” refers to the amino acid sequences of substantially purified GTPAP obtained from
10 any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term “agonist” refers to a molecule which intensifies or mimics the biological activity of GTPAP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GTPAP either by directly interacting with
15 GTPAP or by acting on components of the biological pathway in which GTPAP participates.

An “allelic variant” is an alternative form of the gene encoding GTPAP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to
20 allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding GTPAP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as GTPAP or a
25 polypeptide with at least one functional characteristic of GTPAP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding GTPAP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding GTPAP. The encoded protein may also be “altered,” and may contain deletions, insertions, or
30 substitutions of amino acid residues which produce a silent change and result in a functionally equivalent GTPAP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of GTPAP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged -

amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

5 The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein
10 molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

15 The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of GTPAP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of GTPAP either by directly interacting with GTPAP or by acting on components of the biological pathway in which GTPAP participates.

20 The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind GTPAP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired.
25 Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

30 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is

complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic GTPAP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding GTPAP or fragments of GTPAP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the

protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
5	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
10	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
15	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
20	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

25 Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

30 A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

 The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function
35 of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

 A "fragment" is a unique portion of GTPAP or the polynucleotide encoding GTPAP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up
40 to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment

used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:30-58 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:30-58, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:30-58 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:30-58 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:30-58 and the region of SEQ ID NO:30-58 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-29 is encoded by a fragment of SEQ ID NO:30-58. A fragment of SEQ ID NO:1-29 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-29. For example, a fragment of SEQ ID NO:1-29 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-29. The precise length of a fragment of SEQ ID NO:1-29 and the region of SEQ ID NO:1-29 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the

substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62
Reward for match: 1
Penalty for mismatch: -2
Open Gap: 5 and Extension Gap: 2 penalties
Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example,
5 as defined by a particular SEQ ID number, or may be measured over a shorter length, for example,
over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at
least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous
nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported
by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a
10 length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode
similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes
in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid
sequences that all encode substantially the same protein.

15 The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to
the percentage of residue matches between at least two polypeptide sequences aligned using a
standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some
alignment methods take into account conservative amino acid substitutions. Such conservative
substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the
20 site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default
parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e
sequence alignment program (described and referenced above). For pairwise alignments of
polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap
25 penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default
residue weight table. As with polynucleotide alignments, the percent identity is reported by
CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise
comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9
30 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

10 “Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term “humanized antibody” refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

15 “Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

25 Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY;

specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration
5 may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to
10 those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A
15 hybridization complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide
20 sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

25 The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of GTPAP. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other
30 biological, functional, or immunological properties of GTPAP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Probe" refers to nucleic acid sequences encoding GTPAP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5. 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to

5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

20 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be use to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding GTPAP, or fragments thereof, or GTPAP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA,-

RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may

have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human GTPase associated proteins (GTPAP), the polynucleotides encoding GTPAP, and the use of these compositions for the diagnosis, treatment, or prevention of cell proliferative, autoimmune/inflammatory, and immune system disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding GTPAP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each GTPAP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each GTPAP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical

methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding GTPAP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:30-58 and to distinguish between SEQ ID NO:30-58 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express GTPAP as a fraction of total tissues expressing GTPAP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing GTPAP as a fraction of total tissues expressing GTPAP. Column 5 lists the vectors used to subclone each cDNA library. Of particular note is the specific expression of SEQ ID NO:43 in only one library, a human testis tissue library; the specific expression of SEQ ID NO:49 in only 4 libraries, one of which is associated with cell proliferation and 3 of which are associated with inflammation; and the specific expression of SEQ ID NO:40 in only 5 libraries, 3 of which are associated with cell proliferation and one of which is associated with inflammation.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding GTPAP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses GTPAP variants. A preferred GTPAP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the GTPAP amino acid sequence, and which contains at least one functional or structural characteristic of GTPAP.

The invention also encompasses polynucleotides which encode GTPAP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:30-58, which encodes GTPAP.

The invention also encompasses a variant of a polynucleotide sequence encoding GTPAP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding GTPAP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:30-58 which has at least about 70%, or alternatively at least about 90%, or even at least about

95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:30-58. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of GTPAP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding GTPAP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring GTPAP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode GTPAP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring GTPAP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding GTPAP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding GTPAP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode GTPAP and GTPAP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding GTPAP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:30-58 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment

of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding GTPAP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been

size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

5 Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate
10 software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

 In another embodiment of the invention, polynucleotide sequences or fragments thereof
15 which encode GTPAP may be cloned in recombinant DNA molecules that direct expression of GTPAP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express GTPAP.

 The nucleotide sequences of the present invention can be engineered using methods generally
20 known in the art in order to alter GTPAP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction
25 sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

 In another embodiment, sequences encoding GTPAP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, GTPAP itself or a fragment thereof may be synthesized using chemical methods. For
30 example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of GTPAP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active GTPAP, the nucleotide sequences encoding GTPAP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding GTPAP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding GTPAP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding GTPAP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding GTPAP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding GTPAP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or

tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding GTPAP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding GTPAP can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding GTPAP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of GTPAP are needed, e.g. for the production of antibodies, vectors which direct high level expression of GTPAP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of GTPAP. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of GTPAP. Transcription of sequences encoding GTPAP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding GTPAP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader

sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses GTPAP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of GTPAP in cell lines is preferred. For example, sequences encoding GTPAP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *ap^r* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system.

(See, e.g., Rhodes, C.A. (1995) *Methods Mol. Biol.* 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding GTPAP is inserted within a marker gene sequence, transformed cells containing
5 sequences encoding GTPAP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GTPAP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding GTPAP and that express
10 GTPAP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of GTPAP using either
15 specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on GTPAP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See,
20 e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and
25 may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding GTPAP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding GTPAP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available,
30 and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for

ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding GTPAP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein
5 produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode GTPAP may be designed to contain signal sequences which direct secretion of GTPAP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the
10 inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for
15 post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding GTPAP may be ligated to a heterologous sequence resulting in translation of a
20 fusion protein in any of the aforementioned host systems. For example, a chimeric GTPAP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of GTPAP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST),
25 maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity
30 purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the GTPAP encoding sequence and the heterologous protein sequence, so that GTPAP may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10).

A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled GTPAP may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

Fragments of GTPAP may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of GTPAP may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of GTPAP and GTPase associated proteins. In addition, the expression of GTPAP is closely associated with proliferating tissues associated with cancer and fetal development, inflamed tissues, and tissues involved in the immune response. Therefore, GTPAP appears to play a role in cell proliferative, autoimmune/inflammatory, and immune system disorders. In the treatment of disorders associated with increased GTPAP expression or activity, it is desirable to decrease the expression or activity of GTPAP. In the treatment of disorders associated with decreased GTPAP expression or activity, it is desirable to increase the expression or activity of GTPAP.

Therefore, in one embodiment, GTPAP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis,

autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, 5 glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner 10 syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and an immune system disorder, such as acquired immunodeficiency syndrome (AIDS), X-linked agammaglobinemia of Bruton, common variable immunodeficiency (CVI), DiGeorge's syndrome (thymic hypoplasia), thymic dysplasia, isolated IgA deficiency, severe combined immunodeficiency disease (SCID), 15 immunodeficiency with thrombocytopenia and eczema (Wiskott-Aldrich syndrome), Chediak-Higashi syndrome, chronic granulomatous diseases, hereditary angioneurotic edema, and immunodeficiency associated with Cushing's disease, leukemias such as multiple myeloma, and lymphomas such as Hodgkin's disease.

In another embodiment, a vector capable of expressing GTPAP or a fragment or derivative 20 thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified GTPAP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not 25 limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GTPAP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of GTPAP including, but not limited to, those listed above.

In a further embodiment, an antagonist of GTPAP may be administered to a subject to treat or 30 prevent a disorder associated with increased expression or activity of GTPAP. Examples of such disorders include, but are not limited to, those cell proliferative, autoimmune/inflammatory, and immune system disorders described above. In one aspect, an antibody which specifically binds GTPAP may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for

bringing a pharmaceutical agent to cells or tissues which express GTPAP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding GTPAP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of GTPAP including, but not limited to, those described above.

5 In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the
10 various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of GTPAP may be produced using methods which are generally known in the art. In particular, purified GTPAP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GTPAP. Antibodies to GTPAP may
15 also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans,
20 and others may be immunized by injection with GTPAP or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in
25 humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to GTPAP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the
30 entire amino acid sequence of a small, naturally occurring molecule. Short stretches of GTPAP amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to GTPAP may be prepared using any technique which provides for

the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and
5 Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda,
10 S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce GTPAP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

15 Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for GTPAP may also be generated.
20 For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

25 Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between GTPAP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies
30 reactive to two non-interfering GTPAP epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for GTPAP. Affinity is expressed as an

association constant, K_a , which is defined as the molar concentration of GTPAP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple GTPAP epitopes, represents the average affinity, or avidity, of the antibodies for GTPAP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular GTPAP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the GTPAP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of GTPAP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of GTPAP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding GTPAP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding GTPAP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding GTPAP. Thus, complementary molecules or fragments may be used to modulate GTPAP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding GTPAP.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides

encoding GTPAP. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

Genes encoding GTPAP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding GTPAP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in
5 the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing
10 complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding GTPAP. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for
15 the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

20 Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding GTPAP.

25 Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of
30 candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques

for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding GTPAP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA
5 constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase
10 linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

15 Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.
20 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical
25 or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of GTPAP, antibodies to GTPAP, and mimetics, agonists, antagonists, or inhibitors of GTPAP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical
30 carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial,

intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's

solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of GTPAP, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example GTPAP or fragments thereof, antibodies of GTPAP, and agonists, antagonists or inhibitors of GTPAP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be

determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions
5 which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

10 The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and
15 response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about $0.1 \mu\text{g}$ to $100,000 \mu\text{g}$, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and
20 methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

25 In another embodiment, antibodies which specifically bind GTPAP may be used for the diagnosis of disorders characterized by expression of GTPAP, or in assays to monitor patients being treated with GTPAP or agonists, antagonists, or inhibitors of GTPAP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for GTPAP include methods which utilize the antibody and a label to detect
30 GTPAP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring GTPAP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of GTPAP expression. Normal or standard values for GTPAP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to
5 GTPAP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of GTPAP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

10 In another embodiment of the invention, the polynucleotides encoding GTPAP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of GTPAP may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess
15 expression of GTPAP, and to monitor regulation of GTPAP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding GTPAP or closely related molecules may be used to identify nucleic acid sequences which encode GTPAP. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a
20 conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding GTPAP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the GTPAP encoding sequences. The hybridization probes of the subject
25 invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:30-58 or from genomic sequences including promoters, enhancers, and introns of the GTPAP gene.

Means for producing specific hybridization probes for DNAs encoding GTPAP include the cloning of polynucleotide sequences encoding GTPAP or GTPAP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may
30 be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding GTPAP may be used for the diagnosis of disorders associated with expression of GTPAP. Examples of such disorders include, but are not limited to, a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and an immune system disorder, such as acquired immunodeficiency syndrome (AIDS), X-linked agammaglobinemia of Bruton, common variable immunodeficiency (CVI), DiGeorge's syndrome (thymic hypoplasia), thymic dysplasia, isolated IgA deficiency, severe combined immunodeficiency disease (SCID), immunodeficiency with thrombocytopenia and eczema (Wiskott-Aldrich syndrome), Chediak-Higashi syndrome, chronic granulomatous diseases, hereditary angioneurotic edema, and immunodeficiency associated with Cushing's disease, leukemias such as multiple myeloma, and lymphomas such as Hodgkin's disease. The polynucleotide sequences encoding GTPAP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered GTPAP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GTPAP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding GTPAP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a
5 suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding GTPAP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to
10 monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of GTPAP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GTPAP, under conditions suitable for hybridization or
15 amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

20 Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

25 With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development
30 or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding GTPAP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a

polynucleotide encoding GTPAP, or a fragment of a polynucleotide complementary to the polynucleotide encoding GTPAP, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

5 Methods which may also be used to quantify the expression of GTPAP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is
10 presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

 In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify
15 genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

 Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci.
20 USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

 In another embodiment of the invention, nucleic acid sequences encoding GTPAP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence.
25 The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

30 Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the

location of the gene encoding GTPAP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

5 In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides
10 valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention
15 may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

 In another embodiment of the invention, GTPAP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a
20 solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between GTPAP and the agent being tested may be measured.

 Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are
25 synthesized on a solid substrate. The test compounds are reacted with GTPAP, or fragments thereof, and washed. Bound GTPAP is then detected by methods well known in the art. Purified GTPAP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

30 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GTPAP specifically compete with a test compound for binding GTPAP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with GTPAP.

In additional embodiments, the nucleotide sequences which encode GTPAP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

5 Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

10 Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. Nos. 60/109,592, 60/118,610, and 60/127,990 are hereby expressly incorporated
15 by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some
20 tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

25 Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A⁺) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA
30 purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the

recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-
5 1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent E. coli
10 cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a
15 Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

20 Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence
25 scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific)
30 or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled

polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene

families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:30-58. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

100

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding GTPAP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of GTPAP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:30-58 were produced by extension of

an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham

Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

5 The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was
10 quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

 In like manner, the nucleotide sequences of SEQ ID NO:30-58 are used to obtain 5'
15 regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

VI. Labeling and Use of Individual Hybridization Probes

 Hybridization probes derived from SEQ ID NO:30-58 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base
20 pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 µCi of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a
25 SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon
30 membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and

compared.

VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the GTPAP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring GTPAP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of GTPAP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the GTPAP-encoding transcript.

IX. Expression of GTPAP

Expression and purification of GTPAP is achieved using bacterial or virus-based expression

systems. For expression of GTPAP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express GTPAP upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of GTPAP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding GTPAP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, GTPAP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from GTPAP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified GTPAP obtained by these methods can be used directly in the following activity assay.

X. Demonstration of GTPAP Activity

The role of GTPAP can be assayed in vitro by monitoring the mobilization of Ca^{++} as part of the signal transduction pathway. (See, e.g., Grynkievicz, G. et al. (1985) J. Biol. Chem. 260:3440; McColl, S. et al. (1993) J. Immunol. 150:4550-4555; and Aussel, C. et al. (1988) J. Immunol. 140:215.) The assay requires preloading neutrophils or T cells with a fluorescent dye such as FURA-2.

Upon binding Ca^{++} , FURA-2 exhibits an absorption shift that can be observed by scanning the excitation spectrum between 300 and 400 nm, while monitoring the emission at 510 nm. When the cells are exposed to one or more activating stimuli artificially (i.e., anti-CD3 antibody ligation of the T cell receptor) or physiologically (i.e., by allogeneic stimulation), Ca^{++} flux takes place. Ca^{++} flux results from the release of Ca^{++} from intracellular organelles or from Ca^{++} entry into the cell through activated Ca^{++} channels. This flux can be observed and quantified by assaying the cells in a fluorometer or fluorescence activated cell sorter. Measurements of Ca^{++} flux are compared between cells in their normal state and those preloaded with GTPAP. Increased mobilization attributable to increased GTPAP availability results in increased emission.

Alternatively, GTPAP activity is measured by quantifying the amount of a non-hydrolyzable GTP analogue, GTP γ S, bound over a 10 minute incubation period. Varying amounts of GTPAP are incubated at 30°C in 50mM Tris buffer, pH 7.5, containing 1mM dithiothreitol, 1mM EDTA and 1 μ M [^{35}S]GTP γ S. Samples are passed through nitrocellulose filters and washed twice with a buffer consisting of 50mM Tris-HCl, pH 7.8, 1mM NaN_3 , 10mM MgCl_2 , 1mM EDTA, 0.5mM dithiothreitol, 0.01mM PMSF, and 200mM NaCl. The filter-bound counts are measured by liquid scintillation to quantify the amount of bound [^{35}S]GTP γ S. GTPAP activity may also be measured as the amount of GTP hydrolysed over a 10 minute incubation period at 37°C. GTPAP is incubated in 50mM Tris-HCl buffer, pH 7.8, containing 1mM dithiothreitol, 2mM EDTA, 10 μ M [α - ^{32}P]GTP, and 1 μ M H-rab protein. GTPase activity is initiated by adding MgCl_2 to a final concentration of 10 mM. Samples are removed at various time points, mixed with an equal volume of ice-cold 0.5mM EDTA, and frozen. Aliquots are spotted onto polyethyleneimine-cellulose thin layer chromatography plates, which are developed in 1M LiCl, dried, and autoradiographed. The signal detected is proportional to GTPAP activity.

Alternatively, GTPAP activity may be demonstrated as the ability to interact with its associated $\text{G}\alpha$ or LMW GTPase in an in vitro binding assay. The candidate GTPases are expressed as fusion proteins with glutathione S-transferase (GST), and purified by affinity chromatography on glutathione-Sepharose. The GTPases are loaded with GDP by incubating 20 mM Tris buffer, pH 8.0, containing 100 mM NaCl, 2 mM EDTA, 5 mM MgCl_2 , 0.2 mM DTT, 100 μ M AMP-PNP and 10 μ M GDP at 30°C for 20 minutes. GTPAP is expressed as a FLAG fusion proteins in a baculovirus system. Extracts of these baculovirus cells containing GTPAP-FLAG fusion proteins are precleared with GST beads, then incubated with GST-GTPase fusion proteins. The complexes formed are precipitated by glutathione-Sepharose and separated by SDS-polyacrylamide gel electrophoresis. The separated proteins are blotted onto nitrocellulose membranes and probed with commercially available anti-

FLAG antibodies. GTPAP activity is proportional to the amount of GTPAP-FLAG fusion protein detected in the complex.

XI. Functional Assays

GTPAP function is assessed by expressing the sequences encoding GTPAP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of GTPAP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding GTPAP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding GTPAP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of GTPAP Specific Antibodies

GTPAP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

5 Alternatively, the GTPAP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

10 Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-
15 GTPAP activity by, for example, binding the peptide or GTPAP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring GTPAP Using Specific Antibodies

Naturally occurring or recombinant GTPAP is substantially purified by immunoaffinity chromatography using antibodies specific for GTPAP. An immunoaffinity column is constructed by
20 covalently coupling anti-GTPAP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing GTPAP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of GTPAP (e.g., high ionic strength
25 buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/GTPAP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GTPAP is collected.

XIV. Identification of Molecules Which Interact with GTPAP

GTPAP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter
30 reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled GTPAP, washed, and any wells with labeled GTPAP complex are assayed. Data obtained using different concentrations of GTPAP are used to calculate values for the number, affinity, and association of

GTPAP with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be
5 understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	30	708398	SYNORAT04	568987X31 (MMLR3DT01), 708398H1, 708398X11, 708398X15, 708398X16, 708398X17, and 708398X21 (SYNORAT04), 2170523F6 (ENDCNOT03), 3374750H1 (CONNTUT05)
2	31	1259937	MENITUT03	913652R6 (STOMNOT02), 1259937F6 and 1259937H1 (MENITUT03), 1476721F1 (CORPNOT02), 1729248F6 (BRSTTUT08), 2191963H1 (THYRTUT03), 3129757F6 (LUNGTUT12), 3268746X15F1 (BRAINOT20), 3428891F6 (SKINNOT04)
3	32	1452285	PENITUT01	1452285F6 and 1452285H1 (PENITUT01), 2605011H1 (LUNGTUT07), 3505135H1 (ADRENOT11)
4	33	1812894	PROSTUT12	1812894H1, 1812894X12 and 1809113T6 (PROSTUT12), 1904479F6 (OVARNOT07), 2232535X15F1 and 2232535X18F1 (PROSNOT16), 2267486X16C1 (UTRSNOT02), 2508562F6 (CONUTUT01)
5	34	3074884	BONEUNT01	225362F1 (PANCNOT01), 900707R1 (BRSTTUT03), 1339234F6 (COLNTUT03), 1759046R6 (PITUNOT03), 3074884H1 (BONEUNT01), SBDA02767F1
6	35	3452277	UTRSNON03	1684553F6 (PROSNOT15), 1951534H1 (PITUNOT01), 3452277H1 (UTRSNON03), 4092781T6 (BSCNSZT01), SBFA01413F1, SBFA03044F1, SBFA01805F1
7	36	4203832	BRAITUT29	723394F1 (SYNOOAT01), 862290R1, and 862290T1 (BRAITUT03), 1560918F1 (SPLNNOT04), 3509241H1 (CONCNOT01), 4203832H1 (BRAITUT29)
8	37	104368	BMARNOT02	104368H1 (BMARNOT02), SAEA03574F1, SAEA01063F1, SAEA00392F1, SAEA02287F1
9	38	1441680	THYRNOT03	1441680F6, 1441680H1, and 1441680T6 (THYRNOT03), 1904222F6 (OVARNOT07), 2477983F6 (SMCANOT01)
10	39	1494955	PROSNON01	965986R1 (BRSTNOT05), 1429037F1 and 1429037T1 (SINTBST01), 1453487F6 (PENITUT01), 1486114H1 (CORPNOT02), 1494955H1 (PROSNON01), 1995426R6 (BRSTTUT03), 2112074X18F1 and 2112348R6 (BRAITUT03)
11	40	1508161	LUNGNOT14	1508161F6 and 1508161H1 (LUNGNOT14), 3334303H1 (BRAIFET01), 4755656H1 (BRAHNOT01)
12	41	1811877	PROSTUT12	493795H1 (HNT2NOT01), 1573136H1 (LNODNOT03), 1811877F6 and 1811877H1 (PROSTUT12), 1825223F6 (LSUBNOT03), 2454143H1 (ENDANOT01), 2651022H1 (BLADTUT08), 3487062H1 (EPIGNOT01), 4536531H1 (OVARNOT12), 4795253H1 (LIVRTUT09), 4854087H1 (TESTNOT10), 4906149H2 (TLYMNOT08), 5196386H1 (LUNLTUT04)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
13	42	1848674	LUNGFET03	1574127F6, 3857867X306F1, and 3857867X313F1 (LNODNOT03), 1848674H1 (LUNGFET03), 1877170F6 (LEUKNOT03), 2695307H1 (UTRSNOT12), 4148654H1 (SINITUT04), 4984182H1 (HELATXT05), 5288671H1 (LIVRTUS02)
14	43	2012970	TESTNOT03	2012970H1, 2012970R6, 2012970X11F (TESTNOT03)
15	44	2254315	OVARTUT01	022341F1 (ADENINB01), 198476R6 (KIDNNOT02), 2254315H1 (OVARTUT01), 2370170F6 (ADREN07), 2451278F6 (ENDANOT01)
16	45	2415545	HNT3AZT01	775722H1 (COLNNOT05), 870320R6 (LUNGAST01), 889023R1 (STOMTUT01), 895724R1 (BRSTNOT05), 1398541F1 (BRAITUT08), 1662585F6 (BRSTNOT09), 2415545H1 (HNT3AZT01), 2985066H1 (CARGDIT01), 3462702H1 (293TF2T01)
17	46	2707969	PONSAZT01	282552R1, 282552X23, and 282552X7 (CARDNOT01), 889783R1 (STOMTUT01), 1995451R6 (BRSTTUT03), 2707969H1 (PONSAZT01), SAAC00359R1.comp, SAAB00136R1, SAAC00330R1
18	47	2817769	BRSTNOT14	041660R1 (TBLYN0T01), 077378R1 (SYNORAB01), 740028R1 (PANCNOT04), 1593201F6 (BRAINOT14), 1924025R6 (BRSTTUT01), 2817769H1 (BRSTNOT14)
19	48	2917557	THYMFET03	473002F1 and 473002R1 (MMLR1DT01), 690999R6 (LUNGUT02), 997483R1 (KIDNTUT01), 1430662F6 (SINTBST01), 1514017F1 (PANCUT01), 1740475R6 (HIPONON01), 2109547H1 (BRAITUT03), 2917557H1 (THYMFET03), 4309528H1 (BRAUNOT01), 4990135H1 (LIVRTUT11)
20	49	3421335	UCMCNOT04	777588R6 and 777588T6 (COLNNOT05), 3421335H1 (UCMCNOT04)
21	50	605761	BRSTTUT01	605761F1, 605761H1, and 605761R6 (BRSTTUT01), 1271131X15 (TESTTUT02), 1516985F1 (PANCUT01), 1524935H1 (UCMCL5T01), 2234846F6 (PANCUT02)
22	51	483862	HNT2RAT01	483862H1 and 483862R1 (HNT2RAT01), 1750781X305F1, 1750781X307D2 (LIVRTUT01)
23	52	1256777	MENITUT03	264041R6 (HNT2AGT01), 826449R1 (PROSNOT06), 1256777H1 (MENITUT03), 2276061R6 (PROSNON01), 4614049H1 (BRAHNOT01)
24	53	2198779	SPLNFET02	1557708F6 (BLADTUT04), 1922490R6 (BRSTTUT01), 2198779H1 (SPLNFET02), 2541193F7 (BONRTUT01), 3039254F6 (BRSTNOT16), 3057079H1 (LNODNOT08), 3105017H1 (COLNUCT03), 4239592H1 (SYNWDIT01), 5064513H1 (ARTFTDT01)

Table 1 (cont.)

Polypeptide SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
25	54	2226116	SEMVNOT01	1662607F6 (BRSTNOT09), 1662607T6 (BRSTNOT09), 2226116F6 (SEMVNOT01), 2226116H1 (SEMVNOT01), 2930011F6 (TLYMNOT04), 3015747T6 (MUSCNOT07), 4087670H1 (LIVRNOT06)
26	55	2504472	CONUTUT01	420365F1 (BRSTNOT01), 762246R1 (BRAITUT02), 907754R2 (COLNNOT09), 1007508H1 (HEALDIT02), 1302342F6 (PLACNOT02), 1913887H1 (PROSTUT04), 2023822F6 (CONNNOT01), 2023822X11R1 (CONNNOT01), 2504472H1 (CONUTUT01), 2951618F6 (KIDNFET01)
27	56	3029920	HEARFET02	354846T6 (RATRNOT01), 418533R6 (BRSTNOT01), 935073R1 (CERVNOT01), 1340722F1 (COLNTUT03), 1416203T6 (BRAINOT12), 1524567F1 (UCMCL5T01), 1773043H1 (MENTUNON3), 2590310H2 (LUNGNOT22), 3029920H1 (HEARFET02), 4873053H1 (COLDNOT01), 5687696H1 (BRAIUNT01)
28	57	3332415	BRAIFET01	118166R1 (MUSCNOT01), 1257348H1 (MENITUT03), 1288237T6 (BRAINOT11), 1335936F6 (COLNNOT13), 1452268H1 (PENITUT01), 1996016R6 (BRSTTUT03), 2116665R6 (BRSTTUT02), 2206894F6 (SINTFET03), 2540063H1 (BONRTUT01), 2808268H1 (BLADTUT08), 3086221H1 (HEAONOT03), 3127508H1 (LUNGTUT12), 3295812H1 (TLYJJINT01), 3332415H1 (BRAIFET01), 3604705H1 (LUNGNOT30), 4821203H1 (PROSTUT17), 4970353H1 (KIDEUNC10), 5055775H1 (COLATMT01)
29	58	4031536	BRAINOT23	029167X3 (SPLNFET01), 350137R1 (LVENNOT01), 408825X1 (EOSIHET02), 689446X23 (LUNGTUT02), 1963062R6 (BRSTNOT04), 2288043R6 (BRAINON01), 4031536H1 (BRAINOT23)

Table 2

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
1	1002	T30 S224 T405 S499 T533 S558 S701 T737 T845 S864 S6 T152 T268 T412 T442 T464 T514 T528 T693 S814 S815 S823 T880 Y117 Y842	N446	G524-T531: ATP/GTP- binding site motif	GTP-binding protein [Mus musculus] g53169	BLAST MOTIFS
2	338	S21 S77 T86 S200 T246 T299 S77 S306 Y131	N244		CAMP- regulated Guanine nucleotide exchange factor [Rattus norvegicus] g4079657	BLAST
3	211	S159 S199	N33, N74	G16-T23: ATP/GTP- binding site motif	GTP-binding protein [Rattus norvegicus] g206543	BLAST MOTIFS PFAM BLOCKS PRINTS
4	516	T14 S42 T237 S270 S347 S360 T371 T395 T433 S500 T3 S13 S96 T316 S430			Fos-related antigen [Rattus norvegicus] g1016712 Rabaptin-4 [H. sapiens] g3832516	BLAST MOTIFS
5	445	T44 T114 T219 T297 S314 S341 S356 T412 T24 S72 T91 T328 T388 T394		G230-T237: ATP/GTP- binding site motif	GTP-binding protein [H. sapiens] g2765411	BLAST MOTIFS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
6	445	S174 S202 S289 S29 S305 S323 T434 T11 T147 T197 T198 S270 S273 S371 S397 Y125	N73		Regulator of G-protein signaling-9 [H. sapiens] g3284012	BLAST
7	281	S182 S210 S254 S13 T56 S110 S182 S32 T46 S66 S177	N130 N181	G31-T38:ATP/GTP- binding site motif	Putative ras- like protein [H. sapiens] g4092830	MOTIFS PRINTS BLAST PFAM
8	301	S92 T2 T3 Y15 S18 S19 S20 S25 S97 T120 S165 S296 T94 S116 T120 S284		E47-G66, S116-E178, Y188-G272: Phosducin signature	Phosducin- like protein [Rattus rattus] g1323727	MOTIFS BLAST PRINTS
9	485	T6 Y57 S82 T91 S112 S187 T231 T257 S309 T6 T81 S132 S157 S210 S241 T462	N460	L49-S82: Beta G protein	Similar to WD domain Beta transducin- like protein [C. elegans] g5596646	MOTIFS BLAST PRINTS
10	447	S420 S94 T107 S118 T167 T179 T308 S390 S39 S58 T78 T113 S129 T160 T167 Y174 T199 S216 S291 T302 T323 T359 T384 S423 T438	N76 N92 N231 N289 N378 N421	M294-T308: Beta transducin	WS beta- transducin repeat protein [Homo sapiens] g4704417	MOTIFS BLAST
11	199	S90 T55 T140 S190		K6-E130: Ras Guanine exchange factor	Putative guanine nucleotide releasing factor [Drosophila affinis] g2981229	MOTIFS BLAST PFAM

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
12	694	S57 S67 S99 T150 T346 S416 S467 S500 T522 T684 S99 T156 S209 S285 T331 T360 T388 T430 T477 T650 T688		L10-I24, M96-L110: Beta transducin	Transducin- like protein [H. sapiens] g414536	MOTIFS BLAST
13	654	T10 S15 T49 S97 S102 S104 S112 S113 S377 S432 S638 T46 S54 S84 S97 T177 S217 T307 S401 S450 S504 T515 S546 T547 S561 Y618	N353 N362 N502	L197-F211: Beta transducin	Similar to the beta transducin family [C. elegans] g2315521	MOTIFS BLAST
14	180	S14		G23-S30: ATP-GTP binding site	Rab7C (small GTP binding protein) [Lotus japonicus] g1370186	MOTIFS BLAST
15	374	T100 T249 S260 T308 T328 S338 S351 S30 T73 T157 S237 T308	N114 N189 N222	G26-T33: ATP-GTP binding site	ATP (GTP) - binding protein [H. sapiens] g3646130	MOTIFS BLAST
16	649	S67 T344 S366 S63 S68 S75 S122 S177 S265 T282 T332 S373 S380 S563 T569 S634 S20 T94 S128 S314 T382 T385 T458 T559		F307-S544: Probable rabGAP domain	Similar to probable rabGAP [C. elegans] g3925265	MOTIFS BLAST PFAM
17	698	T244 S262 S17 T41 T42 T196 S206 S317 S479 S522 S556 T586 T680 T31 S95 T99 T140 T173 S257 T322 S374 T450 S568 T619	N171 N194 N685		Small GTP- binding protein associated protein [Mus musculus] g725274	MOTIFS BLAST

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
18	396	T325 S115 T133 S232 S275 T336 S22 T221 S232 T320	N60 N230 N286	G29-S36: ATP-GTP binding site	Putative GTP- binding protein [C. elegans] g3880615	MOTIFS BLAST
19	634	T197 S3 S5 S9 T14 S132 T197 T285 T553 T40 T56 S160 T189 S261 S582 Y20 Y396 Y419		G52-T59: ATP-GTP binding site	Putative GTP- binding protein [H. sapiens] g3169010	MOTIFS BLAST
20	196	T60 S73 S90 S99 S73 S193		G19-T26: ATP-GTP binding site	Kidney injury associated protein HW052 Acc No W86322 ADP- ribosylation factor-like protein 3 [Rattus norvegicus] g560006	MOTIFS BLAST
21	446	T10 T24 T93 S122 T243 S263 S270 T305 S317 S325 T357 S372 T379 S100 S170 S223 T227 S285 T348	N79	L323-L337: Beta transducin	Putative WD40 repeat protein [A. thaliana] g4191784	MOTIFS BLAST
22	265	T184 T76 T137 S139 T161 T174 T183 S213	N159	L141, L148, L155 L: zipper gene regulatory motif	TipD; similar to beta transducin family [D. discoideum] g2407788	MOTIFS BLAST

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
23	185	T55 S111 S127 S148 S171 S14 S94 Y103		G10-T17: ATP/GTP binding site (P- loop) A4-S72: Ras domain		MOTIFS PFAM PRINTS
24	554	S388 T488 S30 S75 T111 S149 S220 S237 T255 S305 S325 T339 T359 S363 S509 S172 T195 S211 T378 T438 T470 Y203	N5	N297-D336, P345- D383, G481-Q519: Beta-transducin WD40 repeats	WD-repeat protein [Arabidopsis thaliana] g3924603	BLAST MOTIFS PFAM PRINTS
25	434	S164 S341 T347 S36 S68 S92 T286 S364	N22 N383	G259-S266:ATP/GTP binding site (P- loop): G113-R433: GTP1/OBG domain	Predicted GTP binding protein [C. elegans] g3878629	BLAST MOTIFS PFAM BLOCKS PRINTS
26	826	S122 T243 T247 T427 S454 S519 T528 S623 S701 S715 S809 T58 S143 S266 T411 S505 S577 S603 T661 S735 T753 S791 T815	N23 N264 N576 N600 N789	R48-E91, L97-S143, F197 K237, V273- W319, W378-A416, W604 K642, A659- G697: Beta- transducin WD40 repeats	Predicted WD repeat protein [S. cerevesiae] P42935	BLAST MOTIFS PFAM PRINTS
27	618	T414 S59 T105 S126 T139 T143 S196 T203 S311 S325 T370 T390 S477 T483 S541 T583 T94 S148 T247 Y160 Y383 Y456	N118 N154 N346	G11-T18, G425-S432: ATP/GTP binding site (P-loop) R6-K187: Ras domain	GTP-binding protein APD08 [H.sapiens] Accession W75771	BLAST MOTIFS PFAM PRINTS

Table 2 (cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods and Databases
28	596	S17 S21 S50 S152 S153 T533 S539 T594 S36 S38 S80 T163 T169 S183 S211 T240 S306 T329 T417 S457 S508 T545 S45 T64 S88 T124 S139 S299 S451 S459 S528 S568 Y180 Y364		A178-L355: Rho- family guanine nucleotide exchange factor (RhoGEF) domain	Guanine nucleotide regulatory protein (NET1 homologue) [Mus musculus] g3834631	BLAST MOTIFS PFAM BLOCKS
29	589	T108 S20 T90 S127 S176 S278 S467 T521 S522 T189 S254 T284 T292 T321 T324 T345 T364 T423 S444 T484 T527	N572	L252-S289, G293- N329, G333-D369, G373-D409, E413- D449, G453-D489, G493-D532: Beta- transducin WD40 repeats R160-K206: F-box domain	SEL-10 [C.elegans] g2677836	BLAST MOTIFS PFAM PRINTS

Table 3

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
30	628-711	Reproductive (0.256) Nervous (0.154) Gastrointestinal (0.154)	Cell Proliferation (0.692) Inflammation (0.372)	PSPORT1
31	1094-1129	Reproductive (0.268) Cardiovascular (0.146) Nervous (0.146)	Cell Proliferation (0.731) Inflammation (0.219) Neurological (0.049)	pINCY
32	652-703	Cardiovascular (0.375) Reproductive (0.375) Dermatologic (0.125) Endocrine (0.125)	Cell Proliferation (0.875) Trauma (0.125)	pINCY
33	1224-1292	Reproductive (0.412) Gastrointestinal (0.147) Hematopoietic/Immune (0.147)	Cell Proliferation (0.647) Inflammation (0.264)	pINCY
34	16-65	Nervous (0.211) Reproductive (0.197) Gastrointestinal (0.169)	Cell Proliferation (0.507) Inflammation (0.352)	pINCY
35	947-1043	Reproductive (0.444) Nervous (0.333) Gastrointestinal (0.111) Urologic (0.111)	Cell Proliferation (0.667) Inflammation (0.111) Neurological (0.111)	pINCY
36	840-1001	Nervous (0.340) Reproductive (0.208) Gastrointestinal (0.151)	Cell Proliferation (0.641) Inflammation (0.302) Neurological (0.038)	pINCY

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
37	507-551	Hematopoietic/Immune (0.269) Nervous (0.269) Reproductive (0.154)	Inflammation (0.423) Cell Proliferation (0.269)	PBLUESCRIPT
38	218-262	Cardiovascular (0.357) Nervous (0.214) Gastrointestinal (0.143)	Cell Proliferation (0.572) Inflammation (0.214)	pINCY
39	164-208	Nervous (0.280) Reproductive (0.260) Developmental (0.120)	Cell Proliferation (0.740) Inflammation (0.180)	PSPORT1
40	369-411	Cardiovascular (0.250) Developmental (0.250) Gastrointestinal (0.250)	Cell Proliferation (0.500) Inflammation (0.250)	pINCY
41	272-316	Reproductive (0.392) Gastrointestinal (0.118) Hematopoietic/Immune (0.118)	Cell Proliferation (0.626) Inflammation (0.137)	pINCY
42	664-708	Nervous (0.211) Reproductive (0.211) Cardiovascular (0.158)	Cell Proliferation (0.614) Inflammation (0.281)	pINCY
43	226-270	Reproductive (1.000)	Inflammation (1.000)	PBLUESCRIPT
44	11-55	Reproductive (0.254) Gastrointestinal (0.206) Cardiovascular (0.159)	Cell Proliferation (0.698) Inflammation (0.206)	PSPORT1

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
45	637-681	Reproductive (0.281) Nervous (0.188) Gastrointestinal (0.156)	Cell Proliferation (0.781) Inflammation (0.234)	pINCY
46	1016-1060	Nervous (0.330) Reproductive (0.183) Hematopoietic/Immune (0.122)	Cell Proliferation (0.582) Inflammation (0.235)	pINCY
47	737-781	Nervous (0.218) Reproductive (0.188) Gastrointestinal (0.158)	Cell Proliferation (0.655) Inflammation (0.211)	pINCY
48	469-513	Reproductive (0.222) Hematopoietic/Immune (0.160) Nervous (0.160)	Cell Proliferation (0.543) Inflammation (0.272)	pINCY
49	226-270	Gastrointestinal (0.333) Hematopoietic/Immune (0.333) Reproductive (0.333)	Inflammation (1.000)	pINCY
50	456-500	Reproductive (0.289) Gastrointestinal (0.133) Hematopoietic/Immune (0.133)	Cell Proliferation (0.778) Inflammation (0.156)	PSPORT1
51	252-296	Nervous (0.500) Gastrointestinal (0.200) Cardiovascular (0.100)	Cell Proliferation (1.000) Inflammation (0.200)	PBLUESCRIPT
52	60-104	Nervous (0.326) Reproductive (0.326) Cardiovascular (0.152)	Cell proliferation (0.565) Inflammation (0.369)	pINCY

Table 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
53	488-532	Reproductive (0.232) Nervous (0.195) Hematopoietic/Immune (0.146)	Cell proliferation (0.622) Inflammation (0.427)	pINCY
54	686-730	Reproductive (0.250) Gastrointestinal (0.150) Hematopoietic/Immune (0.150)	Cell proliferation (0.700) Inflammation (0.400)	pINCY
55	543-587 1299-1343	Reproductive (0.282) Nervous (0.155) Gastrointestinal (0.146)	Cell proliferation (0.592) Inflammation (0.359)	pINCY
56	345-389 792-836	Nervous (0.268) Reproductive (0.169) Cardiovascular (0.113) Hematopoietic/Immune (0.113)	Cell proliferation (0.606) Inflammation (0.296)	pINCY
57	163-207	Reproductive (0.270) Gastrointestinal (0.189) Nervous (0.156)	Cell proliferation (0.705) Inflammation (0.254)	pINCY
58	381-425 726-770	Nervous (0.317) Reproductive (0.250) Gastrointestinal (0.117)	Cell proliferation (0.450) Inflammation (0.283)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
30	SYNORAT04	This library was constructed using RNA isolated from the wrist synovial membrane tissue of a 62-year-old female with rheumatoid arthritis.
31	MENITUT03	This library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
32	PENITUT01	This library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.
33	PROSTUT12	This library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
34	BONEUNT01	This library was constructed using RNA isolated from Saos-2, a primary osteogenic sarcoma cell line (ATCC HTB-85) derived from an 11-year-old Caucasian female.
35	UTRSNON03	This library was constructed from 6.4 million independent clones from a uterine library. RNA for these libraries was isolated from uterine myometrial tissue removed from a 41-year-old female during a vaginal hysterectomy with dilation and curettage. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Pathology for the associated tumor tissue indicated uterine leiomyoma. The normalization and hybridization conditions were adapted from Soares et al. (Proc.Natl.Acad.Sci. USA (1994) 91:9928).
36	BRAITUT29	This library was constructed using RNA isolated from brain tumor tissue removed from the parietal lobe of a 43-year-old female during excision of a cerebral meningeal lesion. Pathology indicated high grade glioma. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, benign hypertension, and hyperlipidemia.
37	BMARNOT02	This library was constructed using RNA isolated from the bone marrow of 24 male and female Caucasian donors, 16 to 70 years old. (RNA came from Clontech.)

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
38	THYRNOT03	This library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
39	PROSNON01	This normalized library was constructed from 4.4 million independent clones from a prostate library. Starting RNA was made from prostate tissue removed from a 28-year-old Caucasian male who died from a self-inflicted gunshot wound. The normalization and hybridization conditions were adapted from Soares, M.B. et al. (1994) Proc. Natl. Acad. Sci. USA 91:9228-9232, using a longer (19 hour) reannealing hybridization period.
40	LUNGNOT14	This library was constructed using RNA isolated from lung tissue removed from the left lower lobe of a 47-year-old Caucasian male during a segmental lung resection. Pathology for the associated tumor tissue indicated a grade 4 adenocarcinoma, and the parenchyma showed calcified granuloma. Patient history included benign hypertension and chronic obstructive pulmonary disease. Family history included type II diabetes and acute myocardial infarction.
41	PROSTUT12	This library was constructed using RNA isolated from prostate tumor tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated an adenocarcinoma (Gleason grade 2+2). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA).
42	LUNGFET03	This library was constructed using RNA isolated from lung tissue removed from a Caucasian female fetus who died at 20 weeks' gestation.
43	TESTNOT03	This library was constructed using RNA isolated from testicular tissue removed from a 37-year-old Caucasian male, who died from liver disease. Patient history included cirrhosis, jaundice, and liver failure.
44	OVARTUT01	This library was constructed using RNA isolated from ovarian tumor tissue removed from a 43-year-old Caucasian female during removal of the fallopian tubes and ovaries. Pathology indicated grade 2 mucinous cystadenocarcinoma involving the entire left ovary. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, pancreatic cancer, stress reaction, cerebrovascular disease, breast cancer, and uterine cancer.
45	HNT3AZT01	This library was constructed using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated for three days with 0.35 micromolar 5-aza-2'deoxyctidine (AZ).

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
46	PONSAZT01	This library was constructed using RNA isolated from diseased pons tissue from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
47	BRSTNOT14	This library was constructed using RNA isolated from breast tissue obtained from a 62-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated an invasive grade 3 (of 4), nuclear grade 3 (of 3) adenocarcinoma, ductal type. Patient history included a benign colon neoplasm, hyperlipidemia, cardiac dysrhythmia, and obesity. Family history included atherosclerotic coronary artery disease, myocardial infarction, colon cancer, ovarian cancer, lung cancer, and cerebrovascular disease.
48	THYMFET03	This library was constructed using RNA isolated from thymus tissue removed from a Caucasian male fetus.
49	UCMCNOT04	This library was constructed using RNA isolated from mononuclear cells obtained from the umbilical cord blood of multiple individuals of mixed age and sex. The cells were treated with G-CSF.
50	BRSTTUT01	This library was constructed using RNA isolated from breast tumor tissue removed from a 55-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated invasive grade 4 mammary adenocarcinoma of mixed lobular and ductal type, extensively involving the left breast. Family history included benign hypertension, atherosclerotic coronary artery disease, cerebrovascular disease, and depressive disorder.
51	HNT2RAT01	This library was constructed at Stratagene (STR937231), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 24 hours.
52	MENITUT03	This library was constructed using RNA isolated from brain meningioma tissue removed from a 35-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a benign neoplasm in the right cerebellopontine angle of the brain. Patient history included hypothyroidism. Family history included myocardial infarction and breast cancer.
53	SPLNFET02	This library was constructed using RNA isolated from spleen tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
54	SEMVNOT01	This library was constructed using RNA isolated from seminal vesicle tissue removed from a 58-year-old Caucasian male during radical prostatectomy. Pathology for the associated tumor tissue indicated adenocarcinoma (Gleason grade 3+2) of the prostate. Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA). Family history included a malignant breast neoplasm.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Description
55	CONUTUT01	This library was constructed using RNA isolated from sigmoid mesenteric tumor tissue obtained from a 61-year-old female during a total abdominal hysterectomy and bilateral salpingo-oophorectomy with regional lymph node excision. Pathology indicated a metastatic grade 4 malignant mixed müllerian tumor present in the sigmoid mesentery at two sites.
56	HEARFET02	This library was constructed using RNA isolated from heart tissue removed from a Caucasian male fetus, who was stillborn at 23 weeks' gestation with a hypoplastic left heart.
57	BRAIFET01	This library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who was stillborn at 23 weeks' gestation with a hypoplastic left heart.
58	BRAINOT23	This library was constructed using RNA isolated from right temporal lobe tissue removed from a 45-year-old Black male during a brain lobectomy. Pathology for the associated tumor tissue indicated dysembryoplastic neuroepithelial tumor of the right temporal lobe. The right temporal region dura was consistent with calcifying pseudotumor of the neuraxis. The patient presented with convulsive intractable epilepsy, partial epilepsy, and memory disturbance. Patient history included obesity, meningitis, backache, unspecified sleep apnea, acute stress reaction, acquired knee deformity, and chronic sinusitis. Family history included obesity, benign hypertension, cirrhosis of the liver, alcohol abuse, hyperlipidemia, cerebrovascular disease, and type II diabetes.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less <i>Full Length sequences</i> : Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 <i>Assembled ESTs</i> : fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less <i>Full Length sequences</i> : fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score = 1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score = 120 or greater; Match length = 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score = 3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <u>supra</u> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-29 and fragments thereof.

2. A substantially purified variant having at least 90% amino acid sequence identity to the amino acid sequence of claim 1.

3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.

4. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 3.

5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.

6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 3.

7. A method for detecting a polynucleotide, the method comprising the steps of:
(a) hybridizing the polynucleotide of claim 6 to at least one nucleic acid in a sample, thereby forming a hybridization complex; and
(b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide in the sample.

8. The method of claim 7 further comprising amplifying the polynucleotide prior to hybridization.

9. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:30-58 and fragments thereof.

10. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 9.

11. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 9.

12. An expression vector comprising at least a fragment of the polynucleotide of claim 3.

5

13. A host cell comprising the expression vector of claim 12.

14. A method for producing a polypeptide, the method comprising the steps of:

- 10
- a) culturing the host cell of claim 13 under conditions suitable for the expression of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.

15. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.

15

16. A purified antibody which specifically binds to the polypeptide of claim 1.

17. A purified agonist of the polypeptide of claim 1.

20

18. A purified antagonist of the polypeptide of claim 1.

19. A method for treating or preventing a disorder associated with decreased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 15.

25

20. A method for treating or preventing a disorder associated with increased expression or activity of GTPAP, the method comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 18.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

HILLMAN, Jennifer L.

TANG, Y. Tom

BANDMAN, Olga

LAL, Preeti

YUE, Henry

LU, Dyung Aina M.

BAUGHN, Mariah R.

YANG, Junming

AZIMZAI, Yalda

<120> GTPASE ASSOCIATED PROTEINS

<130> PF-0629 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/109,592; 60/118,610; 60/127,990

<151> 1998-11-23; 1999-02-04; 1999-04-06

<160> 58

<170> PERL Program

<210> 1

<211> 1002

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 708398CD1

<400> 1

Met	Pro	Ser	Lys	Phe	Ser	Cys	Arg	Gln	Leu	Arg	Glu	Ala	Gly	Gln
1				5					10					15
Cys	Phe	Glu	Ser	Phe	Leu	Val	Val	Arg	Gly	Leu	Asp	Met	Glu	Thr
				20					25					30
Asp	Arg	Glu	Arg	Leu	Arg	Thr	Ile	Tyr	Asn	Arg	Asp	Phe	Lys	Ile
				35					40					45
Ser	Phe	Gly	Thr	Pro	Ala	Pro	Gly	Phe	Ser	Ser	Met	Leu	Tyr	Gly
				50					55					60
Met	Lys	Ile	Ala	Asn	Leu	Ala	Tyr	Val	Thr	Lys	Thr	Arg	Val	Arg
				65					70					75
Phe	Phe	Arg	Leu	Asp	Arg	Trp	Ala	Asp	Val	Arg	Phe	Pro	Glu	Lys
				80					85					90
Arg	Arg	Met	Lys	Leu	Gly	Ser	Asp	Ile	Ser	Lys	His	His	Lys	Ser
				95					100					105
Leu	Leu	Ala	Lys	Ile	Phe	Tyr	Asp	Arg	Ala	Glu	Tyr	Leu	His	Gly
				110					115					120
Lys	His	Gly	Val	Asp	Val	Glu	Val	Gln	Gly	Pro	His	Glu	Ala	Arg

	125		130		135
Asp Gly Gln Leu	Leu Ile Arg Leu Asp	Leu Asn Arg Lys Glu	Val		
	140		145		150
Leu Thr Leu Arg	Leu Arg Asn Gly Gly	Thr Gln Ser Val Thr	Leu		
	155		160		165
Thr His Leu Phe	Pro Leu Cys Arg Thr	Pro Gln Phe Ala Phe	Tyr		
	170		175		180
Asn Glu Asp Gln	Glu Leu Pro Cys Pro	Leu Gly Pro Gly Glu	Cys		
	185		190		195
Tyr Glu Leu His	Val His Cys Lys Thr	Ser Phe Val Gly Tyr	Phe		
	200		205		210
Pro Ala Thr Val	Leu Trp Glu Leu Leu	Gly Pro Gly Glu Ser	Gly		
	215		220		225
Ser Glu Gly Ala	Gly Thr Phe Tyr Ile	Ala Arg Phe Leu Ala	Ala		
	230		235		240
Val Ala His Ser	Pro Leu Ala Ala Gln	Leu Lys Pro Met Thr	Pro		
	245		250		255
Phe Lys Arg Thr	Arg Ile Thr Gly Asn	Pro Val Val Thr Asn	Arg		
	260		265		270
Ile Glu Glu Gly	Glu Arg Pro Asp Arg	Ala Lys Gly Tyr Asp	Leu		
	275		280		285
Glu Leu Ser Met	Ala Leu Gly Thr Tyr	Tyr Pro Pro Pro Arg	Leu		
	290		295		300
Arg Gln Leu Leu	Pro Met Leu Leu Gln	Gly Thr Ser Ile Phe	Thr		
	305		310		315
Ala Pro Lys Glu	Ile Ala Glu Ile Lys	Ala Gln Leu Glu Thr	Ala		
	320		325		330
Leu Lys Trp Arg	Asn Tyr Glu Val Lys	Leu Arg Leu Leu Leu	His		
	335		340		345
Leu Glu Glu Leu	Gln Met Glu His Asp	Ile Arg His Tyr Asp	Leu		
	350		355		360
Glu Ser Val Pro	Met Thr Trp Asp Pro	Val Asp Gln Asn Pro	Arg		
	365		370		375
Leu Leu Thr Leu	Glu Val Pro Gly Val	Thr Glu Ser Arg Pro	Ser		
	380		385		390
Val Leu Arg Gly	Asp His Leu Phe Ala	Leu Leu Ser Ser Glu	Thr		
	395		400		405
His Gln Glu Asp	Pro Ile Thr Tyr Lys	Gly Phe Val His Lys	Val		
	410		415		420
Glu Leu Asp Arg	Val Lys Leu Ser Phe	Ser Met Ser Leu Leu	Ser		
	425		430		435
Arg Phe Val Asp	Gly Leu Thr Phe Lys	Val Asn Phe Thr Phe	Asn		
	440		445		450
Arg Gln Pro Leu	Arg Val Gln His Arg	Ala Leu Glu Leu Thr	Gly		
	455		460		465
Arg Trp Leu Leu	Trp Pro Met Leu Phe	Pro Val Ala Pro Arg	Asp		
	470		475		480
Val Pro Leu Leu	Pro Ser Asp Val Lys	Leu Lys Leu Tyr Asp	Arg		
	485		490		495
Ser Leu Glu Ser	Asn Pro Glu Gln Leu	Gln Ala Met Arg His	Ile		
	500		505		510
Val Thr Gly Thr	Thr Arg Pro Ala Pro	Tyr Ile Ile Phe Gly	Pro		
	515		520		525
Pro Gly Thr Gly	Lys Thr Val Thr Leu	Val Glu Ala Ile Lys	Gln		
	530		535		540

3/65

	950		955		960
Leu Leu Gln Gly	Leu Ser Lys Leu Ser	Pro Ser Thr Ser Gly	Pro		
	965		970		975
His Ser His Asp	Tyr Leu Pro Gln Glu	Arg Glu Gly Glu Gly	Gly		
	980		985		990
Leu Ser Leu Gln Val	Glu Pro Glu Trp Arg	Asn Glu			
	995		1000		

<210> 2

<211> 338

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1259937CD1

<400> 2

Met Ala Ala Leu Ala	Gln Glu Asp Gly	Trp Thr Lys Gly	Gln Val
1	5	10	15
Leu Val Lys Val Asn	Ser Ala Gly Asp	Ala Ile Gly Leu	Gln Pro
	20	25	30
Asp Ala Arg Gly Val	Ala Thr Ser Leu	Gly Leu Asn Glu	Arg Leu
	35	40	45
Phe Val Val Asn Pro	Gln Glu Val His	Glu Leu Ile Pro	His Pro
	50	55	60
Asp Gln Leu Gly Pro	Thr Val Gly Ser	Ala Glu Gly Leu	Asp Leu
	65	70	75
Val Ser Ala Lys Asp	Leu Ala Gly Gln	Leu Thr Asp His	Asp Trp
	80	85	90
Ser Leu Phe Asn Ser	Ile His Gln Val	Glu Leu Ile His	Tyr Val
	95	100	105
Leu Gly Pro Gln His	Leu Arg Asp Val	Thr Thr Ala Asn	Leu Glu
	110	115	120
Arg Phe Met Arg Arg	Phe Asn Glu Leu	Gln Tyr Trp Val	Ala Thr
	125	130	135
Glu Leu Cys Leu Cys	Pro Val Pro Gly	Pro Arg Ala Gln	Leu Leu
	140	145	150
Arg Lys Phe Ile Lys	Leu Ala Ala His	Leu Lys Glu Gln	Lys Asn
	155	160	165
Leu Asn Ser Phe Phe	Ala Val Met Phe	Gly Leu Ser Asn	Ser Ala
	170	175	180
Ile Ser Arg Leu Ala	His Thr Trp Glu	Arg Leu Pro His	Lys Val
	185	190	195
Arg Lys Leu Tyr Ser	Ala Leu Glu Arg	Leu Leu Asp Pro	Ser Trp
	200	205	210
Asn His Arg Val Tyr	Arg Leu Ala Leu	Ala Lys Leu Ser	Pro Pro
	215	220	225
Val Ile Pro Phe Met	Pro Leu Leu Leu	Lys Asp Met Thr	Phe Ile
	230	235	240
His Glu Gly Asn His	Thr Leu Val Glu	Asn Leu Ile Asn	Phe Glu
	245	250	255
Lys Met Arg Met Met	Ala Arg Ala Ala	Arg Met Leu His	His Cys
	260	265	270

Arg	Ser	His	Asn	Pro	Val	Pro	Leu	Ser	Pro	Leu	Arg	Ser	Arg	Val
				275					280					285
Ser	His	Leu	His	Glu	Asp	Ser	Gln	Val	Ala	Arg	Ile	Ser	Thr	Cys
				290					295					300
Ser	Glu	Gln	Ser	Leu	Ser	Thr	Arg	Ser	Pro	Ala	Ser	Thr	Trp	Ala
				305					310					315
Tyr	Val	Gln	Gln	Leu	Lys	Val	Ile	Asp	Asn	Gln	Arg	Glu	Leu	Ser
				320					325					330
Arg	Leu	Ser	Arg	Glu	Leu	Glu	Pro							
				335										

<210> 3

<211> 211

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1452285CD1

<400> 3

Met	Gln	Ala	Pro	His	Lys	Glu	His	Leu	Tyr	Lys	Leu	Leu	Val	Ile
1				5					10					15
Gly	Asp	Leu	Gly	Val	Gly	Lys	Thr	Ser	Ile	Ile	Lys	Arg	Tyr	Val
				20					25					30
His	Gln	Asn	Phe	Ser	Ser	His	Tyr	Arg	Ala	Thr	Ile	Gly	Val	Asp
				35					40					45
Phe	Ala	Leu	Lys	Val	Leu	His	Trp	Asp	Pro	Glu	Thr	Val	Val	Arg
				50					55					60
Leu	Gln	Leu	Trp	Asp	Ile	Ala	Gly	Gln	Glu	Arg	Phe	Gly	Asn	Met
				65					70					75
Thr	Arg	Val	Tyr	Tyr	Arg	Glu	Ala	Met	Gly	Ala	Phe	Ile	Val	Phe
				80					85					90
Asp	Val	Thr	Arg	Pro	Ala	Thr	Phe	Glu	Ala	Val	Ala	Lys	Trp	Lys
				95					100					105
Asn	Asp	Leu	Asp	Ser	Lys	Leu	Ser	Leu	Pro	Asn	Gly	Lys	Pro	Val
				110					115					120
Ser	Val	Val	Leu	Leu	Ala	Asn	Lys	Cys	Asp	Gln	Gly	Lys	Asp	Val
				125					130					135
Leu	Met	Asn	Asn	Gly	Leu	Lys	Met	Asp	Gln	Phe	Cys	Lys	Glu	His
				140					145					150
Gly	Phe	Val	Gly	Trp	Phe	Glu	Thr	Ser	Ala	Lys	Glu	Asn	Ile	Asn
				155					160					165
Ile	Asp	Glu	Ala	Ser	Arg	Cys	Leu	Val	Lys	His	Ile	Leu	Ala	Asn
				170					175					180
Glu	Cys	Asp	Leu	Met	Glu	Ser	Ile	Glu	Pro	Asp	Val	Val	Lys	Pro
				185					190					195
His	Leu	Thr	Ser	Thr	Lys	Val	Ala	Ser	Cys	Ser	Gly	Cys	Ala	Lys
				200					205					210
Ser														

<210> 4

<211> 516

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1812894CD1

<400> 4

Met	Glu	Thr	Met	Lys	Ala	Val	Ala	Glu	Val	Ser	Glu	Ser	Thr	Lys
1				5					10					15
Ala	Glu	Ala	Val	Ala	Ala	Val	Gln	Arg	Gln	Cys	Gln	Glu	Glu	Val
				20					25					30
Ala	Ser	Leu	Gln	Ala	Ile	Leu	Lys	Asp	Ser	Ile	Ser	Ser	Tyr	Glu
				35					40					45
Ala	Gln	Ile	Thr	Ala	Leu	Lys	Gln	Glu	Arg	Gln	Gln	Gln	Gln	Gln
				50					55					60
Asp	Cys	Glu	Glu	Lys	Glu	Arg	Glu	Leu	Gly	Arg	Leu	Lys	Gln	Leu
				65					70					75
Leu	Ser	Arg	Ala	Tyr	Pro	Leu	Asp	Ser	Leu	Glu	Lys	Gln	Met	Glu
				80					85					90
Lys	Ala	His	Glu	Asp	Ser	Glu	Lys	Leu	Arg	Glu	Ile	Val	Leu	Pro
				95					100					105
Met	Glu	Lys	Glu	Ile	Glu	Glu	Leu	Lys	Ala	Lys	Leu	Leu	Arg	Ala
				110					115					120
Glu	Glu	Leu	Ile	Gln	Glu	Ile	Gln	Arg	Arg	Pro	Arg	His	Ala	Pro
				125					130					135
Ser	Leu	His	Gly	Ser	Thr	Glu	Leu	Leu	Pro	Leu	Ser	Arg	Asp	Pro
				140					145					150
Ser	Pro	Pro	Leu	Glu	Pro	Leu	Glu	Glu	Leu	Ser	Gly	Asp	Gly	Gly
				155					160					165
Pro	Ala	Ala	Glu	Ala	Phe	Ala	His	Asn	Cys	Asp	Asp	Ser	Ala	Ser
				170					175					180
Ile	Ser	Ser	Phe	Ser	Leu	Gly	Gly	Gly	Val	Gly	Ser	Ser	Ser	Ser
				185					190					195
Leu	Pro	Gln	Ser	Arg	Gln	Gly	Leu	Ser	Pro	Glu	Gln	Glu	Glu	Thr
				200					205					210
Ala	Ser	Leu	Val	Ser	Thr	Gly	Thr	Leu	Val	Pro	Glu	Gly	Ile	Tyr
				215					220					225
Leu	Pro	Pro	Pro	Gly	Tyr	Gln	Leu	Val	Pro	Asp	Thr	Gln	Trp	Glu
				230					235					240
Gln	Leu	Gln	Thr	Glu	Gly	Arg	Gln	Leu	Gln	Lys	Asp	Leu	Glu	Ser
				245					250					255
Val	Ser	Arg	Glu	Arg	Asp	Glu	Leu	Gln	Glu	Gly	Leu	Arg	Arg	Ser
				260					265					270
Asn	Glu	Asp	Cys	Ala	Lys	Gln	Met	Gln	Val	Leu	Leu	Ala	Gln	Val
				275					280					285
Gln	Asn	Ser	Glu	Gln	Leu	Leu	Arg	Thr	Leu	Gln	Gly	Thr	Val	Ser
				290					295					300
Gln	Ala	Gln	Glu	Arg	Val	Gln	Leu	Gln	Met	Ala	Glu	Leu	Val	Thr
				305					310					315
Thr	His	Lys	Cys	Leu	His	His	Glu	Val	Lys	Arg	Leu	Asn	Glu	Glu
				320					325					330
Asn	Gln	Gly	Leu	Arg	Ala	Glu	Gln	Leu	Pro	Ser	Ser	Ala	Pro	Gln
				335					340					345
Gly	Ser	Gln	Gln	Glu	Gln	Gly	Glu	Glu	Glu	Ser	Leu	Pro	Ser	Ser

Val	Pro	Glu	Leu	Gln	Gln	Leu	Leu	Cys	Cys	Thr	Arg	Gln	Glu	Ala	350	355	360
Arg	Ala	Arg	Leu	Gln	Ala	Gln	Glu	His	Gly	Ala	Glu	Arg	Leu	Arg	365	370	375
Ile	Glu	Ile	Val	Thr	Leu	Arg	Glu	Ala	Leu	Glu	Glu	Glu	Thr	Val	380	385	390
Ala	Arg	Ala	Ser	Leu	Glu	Gly	Gln	Leu	Arg	Val	Gln	Arg	Glu	Glu	395	400	405
Thr	Glu	Val	Leu	Glu	Ala	Ser	Leu	Cys	Ser	Leu	Arg	Thr	Glu	Met	410	415	420
Glu	Arg	Val	Gln	Gln	Glu	Gln	Ser	Lys	Ala	Gln	Leu	Pro	Asp	Leu	425	430	435
Leu	Ser	Glu	Gln	Arg	Ala	Lys	Val	Leu	Arg	Leu	Gln	Ala	Glu	Leu	440	445	450
Glu	Thr	Ser	Glu	Gln	Val	Gln	Arg	Asp	Phe	Val	Arg	Leu	Ser	Gln	455	460	465
Ala	Leu	Gln	Val	Arg	Leu	Glu	Arg	Ile	Arg	Gln	Ala	Glu	Thr	Leu	470	475	480
Glu	Gln	Val	Arg	Ser	Ile	Met	Asp	Glu	Ala	Pro	Leu	Thr	Asp	Val	485	490	495
Arg	Asp	Ile	Lys	Asp	Thr										500	505	510
															515		

<210> 5

<211> 445

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3074884CD1

<400> 5

Met	Pro	Glu	Asp	Ala	Asp	Glu	Asn	Ala	Glu	Glu	Glu	Leu	Leu	Arg	1	5	10	15
Gly	Glu	Pro	Leu	Leu	Pro	Ala	Gly	Thr	Gln	Arg	Val	Cys	Leu	Val	20	25	30	35
His	Pro	Asp	Val	Lys	Trp	Gly	Pro	Gly	Lys	Ser	Gln	Met	Thr	Arg	40	45	50	55
Ala	Glu	Trp	Gln	Val	Ala	Glu	Ala	Thr	Ala	Leu	Val	His	Thr	Leu	60	65	70	75
Asp	Gly	Trp	Ser	Val	Val	Gln	Thr	Met	Val	Val	Ser	Thr	Lys	Thr	80	85	90	95
Pro	Asp	Arg	Lys	Leu	Ile	Phe	Gly	Lys	Gly	Asn	Phe	Glu	His	Leu	100	105	110	115
Thr	Glu	Lys	Ile	Arg	Gly	Ser	Pro	Asp	Val	Thr	Cys	Val	Phe	Leu	120	125	130	135
Asn	Val	Glu	Arg	Met	Ala	Ala	Pro	Thr	Lys	Lys	Glu	Leu	Glu	Ala	140	145	150	

Ala	Leu	Ala	Glu	Met	Pro	Leu	His	Arg	Ser	Asn	Leu	Lys	Arg	Asp	
				155					160					165	
Val	Ala	His	Leu	Tyr	Arg	Gly	Val	Gly	Ser	Arg	Tyr	Ile	Met	Gly	
				170					175					180	
Ser	Gly	Glu	Ser	Phe	Met	Gln	Leu	Gln	Gln	Arg	Leu	Leu	Arg	Glu	
				185					190					195	
Lys	Glu	Ala	Lys	Ile	Arg	Lys	Ala	Leu	Asp	Arg	Leu	Arg	Lys	Lys	
				200					205					210	
Arg	His	Leu	Leu	Arg	Arg	Gln	Arg	Thr	Arg	Arg	Glu	Phe	Pro	Val	
				215					220					225	
Ile	Ser	Val	Val	Gly	Tyr	Thr	Asn	Cys	Gly	Lys	Thr	Thr	Leu	Ile	
				230					235					240	
Lys	Ala	Leu	Thr	Gly	Asp	Ala	Ala	Ile	Gln	Pro	Arg	Asp	Gln	Leu	
				245					250					255	
Phe	Ala	Thr	Leu	Asp	Val	Thr	Ala	His	Ala	Gly	Thr	Leu	Pro	Ser	
				260					265					270	
Arg	Met	Thr	Val	Leu	Tyr	Val	Asp	Thr	Ile	Gly	Phe	Leu	Ser	Gln	
				275					280					285	
Leu	Pro	His	Gly	Leu	Ile	Glu	Ser	Phe	Ser	Ala	Thr	Leu	Glu	Asp	
				290					295					300	
Val	Ala	His	Ser	Asp	Leu	Ile	Leu	His	Val	Arg	Asp	Val	Ser	His	
				305					310					315	
Pro	Glu	Ala	Glu	Leu	Gln	Lys	Cys	Ser	Val	Leu	Ser	Thr	Leu	Arg	
				320					325					330	
Gly	Leu	Gln	Leu	Pro	Ala	Pro	Leu	Leu	Asp	Ser	Met	Val	Glu	Val	
				335					340					345	
His	Asn	Lys	Val	Asp	Leu	Val	Pro	Gly	Tyr	Ser	Pro	Thr	Glu	Pro	
				350					355					360	
Asn	Val	Val	Pro	Val	Ser	Ala	Leu	Arg	Gly	His	Gly	Leu	Gln	Glu	
				365					370					375	
Leu	Lys	Ala	Glu	Leu	Asp	Ala	Ala	Val	Leu	Lys	Ala	Thr	Gly	Arg	
				380					385					390	
Gln	Ile	Leu	Thr	Leu	Arg	Val	Arg	Leu	Ala	Gly	Ala	Gln	Leu	Ser	
				395					400					405	
Trp	Leu	Tyr	Lys	Glu	Ala	Thr	Val	Gln	Glu	Val	Asp	Val	Ile	Pro	
				410					415					420	
Glu	Asp	Gly	Ala	Ala	Asp	Val	Arg	Val	Ile	Ile	Ser	Asn	Ser	Ala	
				425					430					435	
Tyr	Gly	Lys	Phe	Arg	Lys	Leu	Phe	Pro	Gly						
				440					445						

<210> 6

<211> 445

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3452277CD1

<400> 6

Met	Tyr	Tyr	Gln	Gln	Ala	Leu	Met	Arg	Ser	Thr	Val	Lys	Ser	Ser	
1				5					10					15	
Val	Ser	Leu	Gly	Gly	Ile	Val	Lys	Tyr	Ser	Glu	Gln	Phe	Ser	Ser	

Asn Asp Ala Ile Met Ser Gly Cys Leu Pro Ser Asn Pro Trp Ile	20	25	30
Thr Asp Asp Thr Gln Phe Trp Asp Leu Asn Ala Lys Leu Val Glu	35	40	45
Ile Pro Thr Lys Met Arg Val Glu Arg Trp Ala Phe Asn Phe Ser	50	55	60
Glu Leu Ile Arg Asp Pro Lys Gly Arg Gln Ser Phe Gln Tyr Phe	65	70	75
Leu Lys Lys Glu Phe Ser Gly Glu Asn Leu Gly Phe Trp Glu Ala	80	85	90
Cys Glu Asp Leu Lys Tyr Gly Asp Gln Ser Lys Val Lys Glu Lys	95	100	105
Ala Glu Glu Ile Tyr Lys Leu Phe Leu Ala Pro Gly Ala Arg Arg	110	115	120
Trp Ile Asn Ile Asp Gly Lys Thr Met Asp Ile Thr Val Lys Gly	125	130	135
Leu Lys His Pro His Arg Tyr Val Leu Asp Ala Ala Gln Thr His	140	145	150
Ile Tyr Met Leu Met Lys Lys Asp Ser Tyr Ala Arg Tyr Leu Lys	155	160	165
Ser Pro Ile Tyr Lys Asp Met Leu Ala Lys Ala Ile Glu Pro Gln	170	175	180
Glu Thr Thr Lys Lys Ser Ser Thr Leu Pro Phe Met Arg Arg His	185	190	195
Leu Arg Ser Ser Pro Ser Pro Val Ile Leu Arg Gln Leu Glu Glu	200	205	210
Glu Ala Lys Ala Arg Glu Ala Ala Asn Thr Val Asp Ile Thr Gln	215	220	225
Pro Gly Gln His Met Ala Pro Ser Pro His Leu Thr Val Tyr Thr	230	235	240
Gly Thr Cys Met Pro Pro Ser Pro Ser Ser Pro Phe Ser Ser Ser	245	250	255
Cys Arg Ser Pro Arg Lys Pro Phe Ala Ser Pro Ser Arg Phe Ile	260	265	270
Arg Arg Pro Ser Thr Thr Ile Cys Pro Ser Pro Ile Arg Val Ala	275	280	285
Leu Glu Ser Ser Ser Gly Leu Glu Gln Lys Gly Glu Cys Ser Gly	290	295	300
Ser Met Ala Pro Arg Gly Pro Ser Val Thr Glu Ser Ser Glu Ala	305	310	315
Ser Leu Asp Thr Ser Trp Pro Arg Ser Arg Pro Arg Ala Pro Pro	320	325	330
Lys Ala Arg Met Ala Leu Ser Phe Ser Arg Phe Leu Arg Arg Gly	335	340	345
Cys Leu Ala Ser Pro Val Phe Ala Arg Leu Ser Pro Lys Cys Pro	350	355	360
Ala Val Ser His Gly Arg Val Gln Pro Leu Gly Asp Val Gly Gln	365	370	375
Gln Leu Pro Arg Leu Lys Ser Lys Arg Val Ala Asn Phe Phe Gln	380	385	390
Ile Lys Met Asp Val Pro Thr Gly Ser Gly Thr Cys Leu Met Asp	395	400	405
Ser Glu Asp Ala Gly Thr Gly Glu Ser Gly Asp Arg Ala Thr Glu	410	415	420
	425	430	435

Lys Glu Val Ile Cys Pro Trp Glu Ser Leu
 440 445

<210> 7

<211> 281

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4203832CD1

<400> 7

Met	Lys	Leu	Ala	Ala	Met	Ile	Lys	Lys	Met	Cys	Pro	Ser	Asp	Ser	1	5	10	15
Glu	Leu	Ser	Ile	Pro	Ala	Lys	Asn	Cys	Tyr	Arg	Met	Val	Ile	Leu	20	25	30	35
Gly	Ser	Ser	Lys	Val	Gly	Lys	Thr	Ala	Ile	Val	Ser	Arg	Phe	Leu	40	45	50	55
Thr	Gly	Arg	Phe	Glu	Asp	Ala	Tyr	Thr	Pro	Thr	Ile	Glu	Asp	Phe	60	65	70	75
His	Arg	Lys	Phe	Tyr	Ser	Ile	Arg	Gly	Glu	Val	Tyr	Gln	Leu	Asp	80	85	90	95
Ile	Leu	Asp	Thr	Ser	Gly	Asn	His	Pro	Phe	Pro	Ala	Met	Arg	Cys	100	105	110	115
Leu	Ser	Ile	Leu	Thr	Gly	Asp	Val	Phe	Ile	Leu	Val	Phe	Ser	Leu	120	125	130	135
Asp	Asn	Arg	Asp	Ser	Phe	Glu	Glu	Val	Gln	Arg	Leu	Arg	Gln	Gln	140	145	150	155
Ile	Leu	Asp	Thr	Lys	Ser	Cys	Leu	Lys	Asn	Lys	Thr	Lys	Glu	Asn	160	165	170	175
Val	Asp	Val	Pro	Leu	Val	Ile	Cys	Gly	Asn	Lys	Gly	Asp	Arg	Asp	180	185	190	195
Phe	Tyr	Arg	Glu	Val	Asp	Gln	Arg	Glu	Ile	Glu	Gln	Leu	Val	Gly	200	205	210	215
Asp	Asp	Pro	Gln	Arg	Cys	Ala	Tyr	Phe	Glu	Ile	Ser	Ala	Lys	Lys	220	225	230	235
Asn	Ser	Ser	Leu	Asp	Gln	Met	Phe	Arg	Ala	Leu	Phe	Ala	Met	Ala	240	245	250	255
Lys	Leu	Pro	Ser	Glu	Met	Ser	Pro	Asp	Leu	His	Arg	Lys	Val	Ser	260	265	270	275
Val	Gln	Tyr	Cys	Asp	Val	Leu	His	Lys	Lys	Ala	Leu	Arg	Asn	Lys	280	285	290	295
Lys	Leu	Leu	Arg	Ala	Gly	Ser	Gly	Gly	Gly	Gly	Gly	Asp	Pro	Gly	300	305	310	315
Asp	Ala	Phe	Gly	Ile	Val	Ala	Pro	Phe	Ala	Arg	Arg	Pro	Ser	Val	320	325	330	335
His	Ser	Asp	Leu	Met	Tyr	Ile	Arg	Glu	Lys	Ala	Ser	Ala	Gly	Ser	340	345	350	355
Gln	Ala	Lys	Asp	Lys	Glu	Arg	Cys	Val	Ile	Ser					360	365	370	375

<210> 8

<211> 301

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 104368CD1

<400> 8

Met	Thr	Thr	Leu	Asp	Asp	Lys	Leu	Leu	Gly	Glu	Lys	Leu	Gln	Tyr	
1				5					10					15	
Tyr	Tyr	Ser	Ser	Ser	Glu	Asp	Glu	Asp	Ser	Asp	His	Glu	Asp	Lys	
				20					25					30	
Asp	Arg	Gly	Arg	Cys	Ala	Pro	Ala	Ser	Ser	Ser	Val	Pro	Ala	Glu	
				35					40					45	
Ala	Glu	Leu	Ala	Gly	Glu	Gly	Ile	Ser	Val	Asn	Thr	Gly	Pro	Lys	
				50					55					60	
Gly	Val	Ile	Asn	Asp	Trp	Arg	Arg	Phe	Lys	Gln	Leu	Glu	Thr	Glu	
				65					70					75	
Gln	Arg	Glu	Glu	Gln	Cys	Arg	Glu	Met	Glu	Arg	Leu	Ile	Lys	Lys	
				80					85					90	
Leu	Ser	Met	Thr	Cys	Arg	Ser	His	Leu	Asp	Glu	Glu	Glu	Glu	Gln	
				95					100					105	
Gln	Lys	Gln	Lys	Asp	Leu	Gln	Glu	Lys	Ile	Ser	Gly	Lys	Met	Thr	
				110					115					120	
Leu	Lys	Glu	Phe	Ala	Ile	Met	Asn	Glu	Asp	Gln	Asp	Asp	Glu	Glu	
				125					130					135	
Phe	Leu	Gln	Gln	Tyr	Arg	Lys	Gln	Arg	Met	Glu	Glu	Met	Arg	Gln	
				140					145					150	
Gln	Leu	His	Lys	Gly	Pro	Gln	Phe	Lys	Gln	Val	Phe	Glu	Ile	Ser	
				155					160					165	
Ser	Gly	Glu	Gly	Phe	Leu	Asp	Met	Ile	Asp	Lys	Glu	Gln	Lys	Ser	
				170					175					180	
Ile	Val	Ile	Met	Val	His	Ile	Tyr	Glu	Asp	Gly	Ile	Pro	Gly	Thr	
				185					190					195	
Glu	Ala	Met	Asn	Gly	Cys	Met	Ile	Cys	Leu	Ala	Ala	Glu	Tyr	Pro	
				200					205					210	
Ala	Val	Lys	Phe	Cys	Lys	Val	Lys	Ser	Ser	Val	Ile	Gly	Ala	Ser	
				215					220					225	
Ser	Gln	Phe	Thr	Arg	Asn	Ala	Leu	Pro	Ala	Leu	Leu	Ile	Tyr	Lys	
				230					235					240	
Gly	Gly	Glu	Leu	Ile	Gly	Asn	Phe	Val	Arg	Val	Thr	Asp	Gln	Leu	
				245					250					255	
Gly	Asp	Asp	Phe	Phe	Ala	Val	Asp	Leu	Glu	Ala	Phe	Leu	Gln	Glu	
				260					265					270	
Phe	Gly	Leu	Leu	Pro	Glu	Lys	Glu	Val	Leu	Val	Leu	Thr	Ser	Val	
				275					280					285	
Arg	Asn	Ser	Ala	Thr	Cys	His	Ser	Glu	Asp	Ser	Asp	Leu	Glu	Ile	
				290					295					300	

Asp

<210> 9

<211> 485

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1441680CD1

<400> 9

Met	Arg	Ala	Val	Leu	Thr	Trp	Arg	Asp	Lys	Ala	Glu	His	Cys	Ile	1	5	10	15
Asn	Asp	Ile	Ala	Phe	Lys	Pro	Asp	Gly	Thr	Gln	Leu	Ile	Leu	Ala	20	25	30	
Ala	Gly	Ser	Arg	Leu	Leu	Val	Tyr	Asp	Thr	Ser	Asp	Gly	Thr	Leu	35	40	45	
Leu	Gln	Pro	Leu	Lys	Gly	His	Lys	Asp	Thr	Val	Tyr	Cys	Val	Ala	50	55	60	
Tyr	Ala	Lys	Asp	Gly	Lys	Arg	Phe	Ala	Ser	Gly	Ser	Ala	Asp	Lys	65	70	75	
Ser	Val	Ile	Ile	Trp	Thr	Ser	Lys	Leu	Glu	Gly	Ile	Leu	Lys	Tyr	80	85	90	
Thr	His	Asn	Asp	Ala	Ile	Gln	Cys	Val	Ser	Tyr	Asn	Pro	Ile	Thr	95	100	105	
His	Gln	Leu	Ala	Ser	Cys	Ser	Ser	Ser	Asp	Phe	Gly	Leu	Trp	Ser	110	115	120	
Pro	Glu	Gln	Lys	Ser	Val	Ser	Lys	His	Lys	Ser	Ser	Ser	Lys	Ile	125	130	135	
Ile	Cys	Cys	Ser	Trp	Thr	Asn	Asp	Gly	Gln	Tyr	Leu	Ala	Leu	Gly	140	145	150	
Met	Phe	Asn	Gly	Ile	Ile	Ser	Ile	Arg	Asn	Lys	Asn	Gly	Glu	Glu	155	160	165	
Lys	Val	Lys	Ile	Glu	Arg	Pro	Gly	Gly	Ser	Leu	Ser	Pro	Ile	Trp	170	175	180	
Ser	Ile	Cys	Trp	Asn	Pro	Ser	Arg	Glu	Glu	Arg	Asn	Asp	Ile	Leu	185	190	195	
Ala	Val	Ala	Asp	Trp	Gly	Gln	Lys	Val	Ser	Phe	Tyr	Gln	Leu	Ser	200	205	210	
Gly	Lys	Gln	Ile	Gly	Lys	Asp	Arg	Ala	Leu	Asn	Phe	Asp	Pro	Cys	215	220	225	
Cys	Ile	Ser	Tyr	Phe	Thr	Lys	Gly	Glu	Tyr	Ile	Leu	Leu	Gly	Gly	230	235	240	
Ser	Asp	Lys	Gln	Val	Ser	Leu	Phe	Thr	Lys	Asp	Gly	Val	Arg	Leu	245	250	255	
Gly	Thr	Val	Gly	Glu	Gln	Asn	Ser	Trp	Val	Trp	Thr	Cys	Gln	Ala	260	265	270	
Lys	Pro	Asp	Ser	Asn	Tyr	Val	Val	Val	Gly	Cys	Gln	Asp	Gly	Thr	275	280	285	
Ile	Ser	Phe	Tyr	Gln	Leu	Ile	Phe	Ser	Thr	Val	His	Gly	Val	Tyr	290	295	300	
Lys	Asp	Arg	Tyr	Ala	Tyr	Arg	Asp	Ser	Met	Thr	Asp	Val	Ile	Val	305	310	315	
Gln	His	Leu	Ile	Thr	Glu	Gln	Lys	Val	Arg	Ile	Lys	Cys	Lys	Glu	320	325	330	
Leu	Val	Lys	Lys	Ile	Ala	Ile	Tyr	Arg	Asn	Arg	Leu	Ala	Ile	Gln	335	340	345	
Leu	Pro	Glu	Lys	Ile	Leu	Ile	Tyr	Glu	Leu	Tyr	Ser	Glu	Asp	Leu	350	355	360	

Ser	Asp	Met	His	Tyr	Arg	Val	Lys	Glu	Lys	Ile	Ile	Lys	Lys	Phe
				365					370					375
Glu	Cys	Asn	Leu	Leu	Val	Val	Cys	Ala	Asn	His	Ile	Ile	Leu	Cys
				380					385					390
Gln	Glu	Lys	Arg	Leu	Gln	Cys	Leu	Ser	Phe	Ser	Gly	Val	Lys	Glu
				395					400					405
Arg	Glu	Trp	Gln	Met	Glu	Ser	Leu	Ile	Arg	Tyr	Ile	Lys	Val	Ile
				410					415					420
Gly	Gly	Pro	Pro	Gly	Arg	Glu	Gly	Leu	Leu	Val	Gly	Leu	Lys	Lys
				425					430					435
Met	Tyr	Leu	Leu	Val	Tyr	Ser	Phe	Ile	Leu	Ile	Val	Lys	Asp	Tyr
				440					445					450
Phe	Ser	Leu	Ser	Thr	Asp	Val	Leu	Gly	Asn	Leu	Thr	Trp	Lys	His
				455					460					465
Val	Cys	Lys	Lys	His	Tyr	Trp	Val	Phe	His	Leu	Phe	Ser	Trp	Tyr
				470					475					480
Tyr	Ile	Phe	Val	Gln										
				485										

<210> 10

<211> 447

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1494955CD1

<400> 10

Met	Glu	Leu	Ser	Gln	Met	Ser	Glu	Leu	Met	Gly	Leu	Ser	Val	Leu
1				5					10					15
Leu	Gly	Leu	Leu	Ala	Leu	Met	Ala	Thr	Ala	Ala	Val	Ala	Arg	Gly
				20					25					30
Trp	Leu	Arg	Ala	Gly	Glu	Glu	Arg	Ser	Gly	Arg	Pro	Ala	Cys	Gln
				35					40					45
Lys	Ala	Asn	Gly	Phe	Pro	Pro	Asp	Lys	Ser	Ser	Gly	Ser	Lys	Lys
				50					55					60
Gln	Lys	Gln	Tyr	Gln	Arg	Ile	Arg	Lys	Glu	Lys	Pro	Gln	Gln	His
				65					70					75
Asn	Phe	Thr	His	Arg	Leu	Leu	Ala	Ala	Ala	Leu	Lys	Ser	His	Ser
				80					85					90
Gly	Asn	Ile	Ser	Cys	Met	Asp	Phe	Ser	Ser	Asn	Gly	Lys	Tyr	Leu
				95					100					105
Ala	Thr	Cys	Ala	Asp	Asp	Arg	Thr	Ile	Arg	Ile	Trp	Ser	Thr	Lys
				110					115					120
Asp	Phe	Leu	Gln	Arg	Glu	His	Arg	Ser	Met	Arg	Ala	Asn	Val	Glu
				125					130					135
Leu	Asp	His	Ala	Thr	Leu	Val	Arg	Phe	Ser	Pro	Asp	Cys	Arg	Ala
				140					145					150
Phe	Ile	Val	Trp	Leu	Ala	Asn	Gly	Asp	Thr	Leu	Arg	Val	Phe	Lys
				155					160					165
Met	Thr	Lys	Arg	Glu	Asp	Gly	Gly	Tyr	Thr	Phe	Thr	Ala	Thr	Pro
				170					175					180
Glu	Asp	Phe	Pro	Lys	Lys	His	Lys	Ala	Pro	Val	Ile	Asp	Ile	Gly

	185		190		195
Ile Ala Asn Thr	Gly Lys Phe Ile Met	Thr Ala Ser Ser Asp Thr			
	200		205		210
Thr Val Leu Ile	Trp Ser Leu Lys Gly	Gln Val Leu Ser Thr Ile			
	215		220		225
Asn Thr Asn Gln	Met Asn Asn Thr His	Ala Ala Val Ser Pro Cys			
	230		235		240
Gly Arg Phe Val	Ala Ser Cys Gly Phe	Thr Pro Asp Val Lys Val			
	245		250		255
Trp Glu Val Cys	Phe Gly Lys Lys Gly	Glu Phe Gln Glu Val Val			
	260		265		270
Arg Ala Phe Glu	Leu Lys Gly His Ser	Ala Ala Val His Ser Phe			
	275		280		285
Ala Phe Ser Asn	Asp Ser Arg Arg Met	Ala Ser Val Ser Lys Asp			
	290		295		300
Gly Thr Trp Lys	Leu Trp Asp Thr Asp	Val Glu Tyr Lys Lys Lys			
	305		310		315
Gln Asp Pro Tyr	Leu Leu Lys Thr Gly	Arg Phe Glu Glu Ala Ala			
	320		325		330
Gly Ala Ala Pro	Cys Arg Leu Ala Leu	Ser Pro Asn Ala Gln Val			
	335		340		345
Leu Ala Leu Ala	Ser Gly Ser Ser Ile	His Leu Tyr Asn Thr Arg			
	350		355		360
Arg Gly Glu Lys	Glu Glu Cys Phe Glu	Arg Val His Gly Glu Cys			
	365		370		375
Ile Ala Asn Leu	Ser Phe Asp Ile Thr	Gly Arg Phe Leu Ala Ser			
	380		385		390
Cys Gly Asp Arg	Ala Val Arg Leu Phe	His Asn Thr Pro Gly His			
	395		400		405
Arg Ala Met Val	Glu Glu Met Gln Gly	His Leu Lys Arg Ala Ser			
	410		415		420
Asn Glu Ser Thr	Arg Gln Arg Leu Gln	Gln Gln Leu Thr Gln Ala			
	425		430		435
Gln Glu Thr Leu	Lys Ser Leu Gly Ala	Leu Lys Lys			
	440		445		

<210> 11

<211> 199

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1508161CD1

<400> 11

Met Pro Val Lys Lys	Lys His Arg Ala Arg	Met Ile Glu Tyr Phe
1	5	10
Ile Asp Val Ala Arg	Glu Cys Phe Asn Ile	Gly Asn Phe Asn Ser
	20	25
Leu Met Ala Ile Ile	Ser Gly Met Asn Met	Ser Pro Val Ser Arg
	35	40
Leu Lys Lys Thr Trp	Ala Lys Val Lys Thr	Ala Lys Phe Asp Ile
	50	55

Leu	Glu	His	Gln	Met	Asp	Pro	Ser	Ser	Asn	Phe	Tyr	Asn	Tyr	Arg
				65					70					75
Thr	Ala	Leu	Arg	Gly	Ala	Ala	Gln	Arg	Ser	Leu	Thr	Ala	His	Ser
				80					85					90
Ser	Arg	Glu	Lys	Ile	Val	Ile	Pro	Phe	Phe	Ser	Leu	Leu	Ile	Lys
				95					100					105
Asp	Ile	Tyr	Phe	Leu	Asn	Glu	Gly	Cys	Ala	Asn	Arg	Leu	Pro	Asn
				110					115					120
Gly	His	Val	Asn	Phe	Glu	Lys	Phe	Trp	Glu	Leu	Ala	Lys	Gln	Val
				125					130					135
Ser	Glu	Phe	Met	Thr	Trp	Lys	Gln	Val	Glu	Cys	Pro	Phe	Glu	Arg
				140					145					150
Asp	Arg	Lys	Ile	Leu	Gln	Tyr	Leu	Leu	Thr	Val	Pro	Val	Phe	Ser
				155					160					165
Glu	Asp	Ala	Leu	Tyr	Leu	Ala	Ser	Tyr	Glu	Ser	Glu	Gly	Pro	Glu
				170					175					180
Asn	His	Ile	Glu	Lys	Asp	Arg	Trp	Lys	Ser	Leu	Arg	Ser	Ser	Leu
				185					190					195
Leu	Gly	Arg	Val											

<210> 12

<211> 694

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1811877CD1

<400> 12

Met	Ala	Phe	Asp	Pro	Thr	Ser	Thr	Leu	Leu	Ala	Thr	Gly	Gly	Cys
1				5					10					15
Asp	Gly	Ala	Val	Arg	Val	Trp	Asp	Ile	Val	Arg	His	Tyr	Gly	Thr
				20					25					30
His	His	Phe	Arg	Gly	Ser	Pro	Gly	Val	Val	His	Leu	Val	Ala	Phe
				35					40					45
His	Pro	Asp	Pro	Thr	Arg	Leu	Leu	Leu	Phe	Ser	Ser	Ala	Thr	Asp
				50					55					60
Ala	Ala	Ile	Arg	Val	Trp	Ser	Leu	Gln	Asp	Arg	Ser	Cys	Leu	Ala
				65					70					75
Val	Leu	Thr	Ala	His	Tyr	Ser	Ala	Val	Thr	Ser	Leu	Ala	Phe	Ser
				80					85					90
Ala	Asp	Gly	His	Thr	Met	Leu	Ser	Ser	Gly	Arg	Asp	Lys	Ile	Cys
				95					100					105
Ile	Ile	Trp	Asp	Leu	Gln	Ser	Cys	Gln	Ala	Thr	Arg	Thr	Val	Pro
				110					115					120
Val	Phe	Glu	Ser	Val	Glu	Ala	Ala	Val	Leu	Leu	Pro	Glu	Glu	Pro
				125					130					135
Val	Ser	Gln	Leu	Gly	Val	Lys	Ser	Pro	Gly	Leu	Tyr	Phe	Leu	Thr
				140					145					150
Ala	Gly	Asp	Gln	Gly	Thr	Leu	Arg	Val	Trp	Glu	Ala	Ala	Ser	Gly
				155					160					165
Gln	Cys	Val	Tyr	Thr	Gln	Ala	Gln	Pro	Pro	Gly	Pro	Gly	Gln	Glu
				170					175					180

Leu Thr His Cys Thr	Leu Ala His Thr	Ala Gly Val Val	Leu Thr
185		190	195
Ala Thr Ala Asp His	Asn Leu Leu Leu	Tyr Glu Ala Arg Ser	Leu
200		205	210
Arg Leu Gln Lys Gln	Phe Ala Gly Tyr	Ser Glu Glu Val	Leu Asp
215		220	225
Val Arg Phe Leu Gly	Pro Glu Asp Ser	His Val Val Val	Ala Ser
230		235	240
Asn Ser Pro Cys Leu	Lys Val Phe Glu	Leu Gln Thr Ser	Ala Cys
245		250	255
Gln Ile Leu His Gly	His Thr Asp Ile	Val Leu Ala Leu	Asp Val
260		265	270
Phe Arg Lys Gly Trp	Leu Phe Ala Ser	Cys Ala Lys Asp	Gln Ser
275		280	285
Val Arg Ile Trp Arg	Met Asn Lys Ala	Gly Gln Val Met	Cys Val
290		295	300
Ala Gln Gly Ser Gly	His Thr His Ser	Val Gly Thr Val	Cys Cys
305		310	315
Ser Arg Leu Lys Glu	Ser Phe Leu Val	Thr Gly Ser Gln	Asp Cys
320		325	330
Thr Val Lys Leu Trp	Pro Leu Pro Lys	Ala Leu Leu Ser	Lys Asn
335		340	345
Thr Ala Pro Asp Asn	Gly Pro Ile Leu	Leu Gln Ala Gln	Thr Thr
350		355	360
Gln Arg Cys His Asp	Lys Asp Ile Asn	Ser Val Ala Ile	Ala Pro
365		370	375
Asn Asp Lys Leu Leu	Ala Thr Gly Ser	Gln Asp Arg Thr	Ala Lys
380		385	390
Leu Trp Ala Leu Pro	Gln Cys Gln Leu	Leu Gly Val Phe	Ser Gly
395		400	405
His Arg Arg Gly Leu	Trp Cys Val Gln	Phe Ser Pro Met	Asp Gln
410		415	420
Val Leu Ala Thr Ala	Ser Ala Asp Gly	Thr Ile Lys Leu	Trp Ala
425		430	435
Leu Gln Asp Phe Ser	Cys Leu Lys Thr	Phe Glu Gly His	Asp Ala
440		445	450
Ser Val Leu Lys Val	Ala Phe Val Ser	Arg Gly Thr Gln	Leu Leu
455		460	465
Ser Ser Gly Ser Asp	Gly Leu Val Lys	Leu Trp Thr Ile	Lys Asn
470		475	480
Asn Glu Cys Val Arg	Thr Leu Asp Ala	His Glu Asp Lys	Val Trp
485		490	495
Gly Leu His Cys Ser	Arg Leu Asp Asp	His Ala Leu Thr	Gly Ala
500		505	510
Ser Asp Ser Arg Val	Ile Leu Trp Lys	Asp Val Thr Glu	Ala Glu
515		520	525
Gln Ala Glu Glu Gln	Ala Arg Gln Glu	Glu Gln Val Val	Arg Gln
530		535	540
Gln Glu Leu Asp Asn	Leu Leu His Glu	Lys Arg Tyr Leu	Arg Ala
545		550	555
Leu Gly Leu Ala Ile	Ser Leu Asp Arg	Pro His Thr Val	Leu Thr
560		565	570
Val Ile Gln Ala Ile	Arg Arg Asp Pro	Glu Ala Cys Glu	Lys Leu
575		580	585
Glu Ala Thr Met Leu	Arg Leu Arg Arg	Asp Gln Lys Glu	Ala Leu

	590		595		600
Leu Arg Phe Cys	Val Thr Trp Asn Thr	Asn Ser Arg His Cys	His		
	605		610		615
Glu Ala Gln Ala	Val Leu Gly Val Leu	Leu Arg Arg Glu Ala	Pro		
	620		625		630
Glu Glu Leu Leu	Ala Tyr Glu Gly Val	Arg Ala Ala Leu Glu	Ala		
	635		640		645
Leu Leu Pro Tyr	Thr Glu Arg His Phe	Gln Arg Leu Ser Arg	Thr		
	650		655		660
Leu Gln Ala Ala	Ala Phe Leu Asp Phe	Leu Trp His Asn Met	Lys		
	665		670		675
Leu Pro Val Pro	Ala Ala Ala Pro Thr	Pro Trp Glu Thr His	Lys		
	680		685		690
Gly Ala Leu Pro					

<210> 13

<211> 654

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1848674CD1

<400> 13

Met Glu Arg Ser Gly	Pro Ser Glu Val Thr	Gly Ser Asp Ala Ser
1	5	10 15
Gly Pro Asp Pro Gln	Leu Ala Val Thr Met	Gly Phe Thr Gly Phe
	20	25 30
Gly Lys Lys Ala Arg	Thr Phe Asp Leu Glu	Ala Met Phe Glu Gln
	35	40 45
Thr Arg Arg Thr Ala	Val Glu Arg Ser Arg	Lys Thr Leu Glu Ala
	50	55 60
Arg Glu Lys Glu Glu	Glu Met Asn Arg Glu	Lys Glu Leu Arg Arg
	65	70 75
Gln Asn Glu Asp Ile	Glu Pro Thr Ser Ser	Arg Ser Asn Val Val
	80	85 90
Arg Asp Cys Ser Lys	Ser Ser Ser Arg Asp	Thr Ser Ser Ser Glu
	95	100 105
Ser Glu Gln Ser Ser	Asp Ser Ser Asp Asp	Glu Leu Ile Gly Pro
	110	115 120
Pro Leu Pro Pro Lys	Met Val Gly Lys Pro	Val Asn Phe Met Glu
	125	130 135
Glu Asp Ile Leu Gly	Pro Leu Pro Pro Pro	Leu Asn Glu Glu Glu
	140	145 150
Glu Glu Ala Glu Glu	Glu Glu Glu Glu Glu	Glu Glu Glu Glu Asn
	155	160 165
Pro Val His Lys Ile	Pro Asp Ser His Glu	Ile Thr Leu Lys His
	170	175 180
Gly Thr Lys Thr Val	Ser Ala Leu Gly Leu	Asp Pro Ser Gly Ala
	185	190 195
Arg Leu Val Thr Gly	Gly Tyr Asp Tyr Asp	Val Lys Phe Trp Asp
	200	205 210
Phe Ala Gly Met Asp	Ala Ser Phe Lys Ala	Phe Arg Ser Leu Gln

Pro Cys Glu Cys	215	Leu Gln Tyr Ser Asn Thr	220	225
	230		235	240
Gly Asp Met Ile	245	Glu Cys Ile Lys Gly Asp	250	255
	260		265	270
Ile Asp Arg Asp	275	Lys Gly His Thr Ala Met	280	285
	290		295	300
Gln Tyr Ile Val	305	Thr Trp Glu Val Glu Asn	310	315
	320		325	330
Leu His Thr Gly	335	Tyr Ser Arg Asp Gly Asn	340	345
	350		355	360
Thr Cys Ser Asn	365	His Tyr Lys Gln Ala His	370	375
	380		385	390
Pro Lys Lys Gln	395	Asp Ser Leu Lys Leu Trp	400	405
	410		415	420
Lys Lys Val Ile	425	Cys Phe Ser Pro Asp Asp	430	435
	440		445	450
Leu Ile Ala Ala	455	Thr Phe Gln Arg Val Tyr	460	465
	470		475	480
Arg Asn Leu Thr	485	Val Arg Cys Leu Trp His	490	495
	500		505	510
Asp Ser Gly Thr	515	Lys Gln Ala Glu Thr Leu	520	525
	530		535	540
Asn Val Leu Ala	545	Gln Leu Glu Lys Asp Arg	550	555
	560		565	570
Asp Ile Arg Gln	575	Glu Pro Pro Val Ala Gly	580	585
	590		595	600
Pro Thr Met Phe	605	Ala Lys Ala Ala Glu Asp	610	615
	620		625	630

Thr Met Phe Ala Gln Val Glu Ser Asp Asp Glu Glu Ala Lys Asn
 635 640 645
 Glu Pro Glu Trp Lys Lys Arg Lys Ile
 650

<210> 14
 <211> 180
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2012970CD1

<400> 14
 Met Glu Ala Asn Met Pro Lys Arg Lys Glu Pro Gly Arg Ser Leu
 1 5 10 15
 Arg Ile Lys Val Ile Ser Met Gly Asn Ala Glu Val Gly Lys Ser
 20 25 30
 Cys Ile Ile Lys Arg Tyr Cys Glu Lys Arg Phe Val Ser Lys Tyr
 35 40 45
 Leu Ala Thr Ile Gly Ile Asp Tyr Gly Val Thr Lys Val His Val
 50 55 60
 Arg Asp Arg Glu Ile Lys Val Asn Ile Phe Asp Met Ala Gly His
 65 70 75
 Pro Phe Phe Tyr Glu Val Arg Asn Glu Phe Tyr Lys Asp Thr Gln
 80 85 90
 Gly Val Ile Leu Val Tyr Asp Val Gly Gln Lys Asp Ser Phe Asp
 95 100 105
 Ala Leu Asp Ala Trp Leu Ala Glu Met Lys Gln Glu Leu Gly Pro
 110 115 120
 His Gly Asn Met Glu Asn Ile Ile Phe Val Val Cys Ala Asn Lys
 125 130 135
 Ile Asp Cys Thr Lys His Arg Cys Val Asp Glu Ser Glu Gly Arg
 140 145 150
 Leu Trp Ala Glu Ser Lys Gly Phe Leu Tyr Phe Glu Thr Ser Ala
 155 160 165
 Gln Thr Gly Glu Gly Ile Asn Glu Met Phe Gln Ile His Leu Gly
 170 175 180

<210> 15
 <211> 374
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2254315CD1

<400> 15
 Met Ala Ala Ser Ala Ala Ala Ala Glu Leu Gln Ala Ser Gly Gly
 1 5 10 15
 Pro Arg His Pro Val Cys Leu Leu Val Leu Gly Met Ala Gly Ser

	20		25		30									
Gly	Lys	Thr	Thr	Phe	Val	Gln	Arg	Leu	Thr	Gly	His	Leu	His	Ala
	35								40					45
Gln	Gly	Thr	Pro	Pro	Tyr	Val	Ile	Asn	Leu	Asp	Pro	Ala	Val	His
	50								55					60
Glu	Val	Pro	Phe	Pro	Ala	Asn	Ile	Asp	Ile	Arg	Asp	Thr	Val	Lys
	65								70					75
Tyr	Lys	Glu	Val	Met	Lys	Gln	Tyr	Gly	Leu	Gly	Pro	Asn	Gly	Gly
	80								85					90
Ile	Val	Thr	Ser	Leu	Asn	Leu	Phe	Ala	Thr	Arg	Phe	Asp	Gln	Val
	95								100					105
Met	Lys	Phe	Ile	Glu	Lys	Ala	Gln	Asn	Met	Ser	Lys	Tyr	Val	Leu
	110								115					120
Ile	Asp	Thr	Pro	Gly	Gln	Ile	Glu	Val	Phe	Thr	Trp	Ser	Ala	Ser
	125								130					135
Gly	Thr	Ile	Ile	Thr	Glu	Ala	Leu	Ala	Ser	Ser	Phe	Pro	Thr	Val
	140								145					150
Val	Ile	Tyr	Val	Met	Asp	Thr	Ser	Arg	Ser	Thr	Asn	Pro	Val	Thr
	155								160					165
Phe	Met	Ser	Asn	Met	Leu	Tyr	Ala	Cys	Ser	Ile	Leu	Tyr	Lys	Thr
	170								175					180
Lys	Leu	Pro	Phe	Ile	Val	Val	Met	Asn	Lys	Thr	Asp	Ile	Ile	Asp
	185								190					195
His	Ser	Phe	Ala	Val	Glu	Trp	Met	Gln	Asp	Phe	Glu	Ala	Phe	Gln
	200								205					210
Asp	Ala	Leu	Asn	Gln	Glu	Thr	Thr	Tyr	Val	Ser	Asn	Leu	Thr	Arg
	215								220					225
Ser	Met	Ser	Leu	Val	Leu	Asp	Glu	Phe	Tyr	Ser	Ser	Leu	Arg	Val
	230								235					240
Val	Gly	Val	Ser	Ala	Val	Leu	Gly	Thr	Gly	Leu	Asp	Glu	Leu	Phe
	245								250					255
Val	Gln	Val	Thr	Ser	Ala	Ala	Glu	Glu	Tyr	Glu	Arg	Glu	Tyr	Arg
	260								265					270
Pro	Glu	Tyr	Glu	Arg	Leu	Lys	Lys	Ser	Leu	Ala	Asn	Ala	Glu	Ser
	275								280					285
Gln	Gln	Gln	Arg	Glu	Gln	Leu	Glu	Arg	Leu	Arg	Lys	Asp	Met	Gly
	290								295					300
Ser	Val	Ala	Leu	Asp	Ala	Gly	Thr	Ala	Lys	Asp	Ser	Leu	Ser	Pro
	305								310					315
Val	Leu	His	Pro	Ser	Asp	Leu	Ile	Leu	Thr	Arg	Gly	Thr	Leu	Asp
	320								325					330
Glu	Glu	Asp	Glu	Glu	Ala	Asp	Ser	Asp	Thr	Asp	Asp	Ile	Asp	His
	335								340					345
Arg	Val	Thr	Glu	Glu	Ser	His	Glu	Glu	Pro	Ala	Phe	Gln	Asn	Phe
	350								355					360
Met	Gln	Glu	Ser	Met	Ala	Gln	Tyr	Trp	Lys	Arg	Asn	Asn	Lys	
	365								370					

<210> 16

<211> 649

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2415545CD1

<400> 16

Met	Glu	Gly	Ala	Gly	Tyr	Arg	Val	Val	Phe	Glu	Lys	Gly	Gly	Val	1	5	10	15
Tyr	Leu	His	Thr	Ser	Ala	Lys	Lys	Tyr	Gln	Asp	Arg	Asp	Ser	Leu	20	25	30	
Ile	Ala	Gly	Val	Ile	Arg	Val	Val	Glu	Lys	Asp	Asn	Asp	Val	Leu	35	40	45	
Leu	His	Trp	Ala	Pro	Val	Glu	Glu	Ala	Gly	Asp	Ser	Thr	Gln	Ile	50	55	60	
Leu	Phe	Ser	Lys	Lys	Asp	Ser	Ser	Gly	Gly	Asp	Ser	Cys	Ala	Ser	65	70	75	
Glu	Glu	Glu	Pro	Thr	Phe	Asp	Pro	Gly	Tyr	Glu	Pro	Asp	Trp	Ala	80	85	90	
Val	Ile	Ser	Thr	Val	Arg	Pro	Gln	Pro	Cys	His	Ser	Glu	Pro	Thr	95	100	105	
Arg	Gly	Ala	Glu	Pro	Ser	Cys	Pro	Gln	Gly	Ser	Trp	Ala	Phe	Ser	110	115	120	
Val	Ser	Leu	Gly	Glu	Leu	Lys	Ser	Ile	Arg	Arg	Ser	Lys	Pro	Gly	125	130	135	
Leu	Ser	Trp	Ala	Tyr	Leu	Val	Leu	Val	Thr	Gln	Ala	Gly	Gly	Ser	140	145	150	
Leu	Pro	Ala	Leu	His	Phe	His	Arg	Gly	Gly	Thr	Arg	Ala	Leu	Leu	155	160	165	
Arg	Val	Leu	Ser	Arg	Tyr	Leu	Leu	Leu	Ala	Ser	Ser	Pro	Gln	Asp	170	175	180	
Ser	Arg	Leu	Tyr	Leu	Val	Phe	Pro	His	Asp	Ser	Ser	Ala	Leu	Ser	185	190	195	
Asn	Ser	Phe	His	His	Leu	Gln	Leu	Phe	Asp	Gln	Asp	Ser	Ser	Asn	200	205	210	
Val	Val	Ser	Arg	Phe	Leu	Gln	Asp	Pro	Tyr	Ser	Thr	Thr	Phe	Ser	215	220	225	
Ser	Phe	Ser	Arg	Val	Thr	Asn	Phe	Phe	Arg	Gly	Ala	Leu	Gln	Pro	230	235	240	
Gln	Pro	Glu	Gly	Ala	Ala	Ser	Asp	Leu	Pro	Pro	Pro	Pro	Asp	Asp	245	250	255	
Glu	Pro	Glu	Pro	Gly	Phe	Glu	Val	Ile	Ser	Cys	Val	Glu	Leu	Gly	260	265	270	
Pro	Arg	Pro	Thr	Val	Glu	Arg	Gly	Pro	Pro	Val	Thr	Glu	Glu	Glu	275	280	285	
Trp	Ala	Arg	His	Val	Gly	Pro	Glu	Gly	Arg	Leu	Gln	Gln	Val	Pro	290	295	300	
Glu	Leu	Lys	Asn	Arg	Ile	Phe	Ser	Gly	Gly	Leu	Ser	Pro	Ser	Leu	305	310	315	
Arg	Arg	Glu	Ala	Trp	Lys	Phe	Leu	Leu	Gly	Tyr	Leu	Ser	Trp	Glu	320	325	330	
Gly	Thr	Ala	Glu	Glu	His	Lys	Ala	His	Ile	Arg	Lys	Lys	Thr	Asp	335	340	345	
Glu	Tyr	Phe	Arg	Met	Lys	Leu	Gln	Trp	Lys	Ser	Val	Ser	Pro	Glu	350	355	360	
Gln	Glu	Arg	Arg	Asn	Ser	Leu	Leu	His	Gly	Tyr	Arg	Ser	Leu	Ile	365	370	375	
Glu	Arg	Asp	Val	Ser	Arg	Thr	Asp	Arg	Thr	Asn	Lys	Phe	Tyr	Glu				

	380		385		390
Gly Pro Glu Asn	Pro Gly Leu Gly Leu	Leu Asn Asp Ile Leu	Leu		
	395		400		405
Thr Tyr Cys Met	Tyr His Phe Asp Leu	Gly Tyr Val Gln Gly	Met		
	410		415		420
Ser Asp Leu Leu	Ser Pro Ile Leu Tyr	Val Ile Gln Asn Glu	Val		
	425		430		435
Asp Ala Phe Trp	Cys Phe Cys Gly Phe	Met Glu Leu Val Gln	Gly		
	440		445		450
Asn Phe Glu Glu	Ser Gln Glu Thr Met	Lys Arg Gln Leu Gly	Arg		
	455		460		465
Leu Leu Leu Leu	Leu Arg Val Leu Asp	Pro Leu Leu Cys Asp	Phe		
	470		475		480
Leu Asp Ser Gln	Asp Ser Gly Ser Leu	Cys Phe Cys Phe Arg	Trp		
	485		490		495
Leu Leu Ile Trp	Phe Lys Arg Glu Phe	Pro Phe Pro Asp Val	Leu		
	500		505		510
Arg Leu Trp Glu	Val Leu Trp Thr Gly	Leu Pro Gly Pro Asn	Leu		
	515		520		525
His Leu Leu Val	Ala Cys Ala Ile Leu	Asp Met Glu Arg Asp	Thr		
	530		535		540
Leu Met Leu Ser	Gly Phe Gly Ser Asn	Glu Ile Leu Lys His	Ile		
	545		550		555
Asn Glu Leu Thr	Met Lys Leu Ser Val	Glu Asp Val Leu Thr	Arg		
	560		565		570
Ala Glu Ala Leu	His Arg Gln Leu Thr	Ala Cys Thr Arg Ala	Ala		
	575		580		585
Pro Gln Arg Ala	Gly Asp Pro Gly Ala	Gly Pro Ala Thr Gln	Ser		
	590		595		600
Pro Thr Ala Pro	Arg Pro Pro Pro Pro	Arg Cys Leu Cys Thr	Pro		
	605		610		615
Thr Arg Ala Pro	Pro Thr Pro Pro Pro	Ser Thr Asp Thr Ala	Pro		
	620		625		630
Gln Pro Asp Ser	Ser Leu Glu Ile Leu	Pro Glu Glu Glu Asp	Glu		
	635		640		645
Gly Ala Asp Ser					

<210> 17

<211> 698

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2707969CD1

<400> 17

Met Cys His Asp	Asp Asp Asp Lys Asp	Pro Val Leu Arg	Val Phe
1	5	10	15
Asp Ser Arg Val	Asp Lys Ile Arg Leu	Leu Asn Val Arg	Thr Pro
	20	25	30
Thr Leu Arg Thr	Ser Met Tyr Gln Lys	Cys Thr Thr Val	Asp Glu
	35	40	45
Ala Glu Lys Ala	Ile Glu Leu Arg Leu	Ala Lys Ile Asp	His Thr

				50					55					60
Ala	Ile	His	Pro	His	Leu	Leu	Asp	Met	Lys	Ile	Gly	Gln	Gly	Lys
				65					70					75
Tyr	Glu	Pro	Gly	Phe	Phe	Pro	Lys	Leu	Gln	Ser	Asp	Val	Leu	Ser
				80					85					90
Thr	Gly	Pro	Ala	Ser	Asn	Lys	Trp	Thr	Lys	Arg	Asn	Ala	Pro	Ala
				95					100					105
Gln	Trp	Arg	Arg	Lys	Asp	Arg	Gln	Lys	Gln	His	Thr	Glu	His	Leu
				110					115					120
Arg	Leu	Asp	Asn	Asp	Gln	Arg	Glu	Lys	Tyr	Ile	Gln	Glu	Ala	Arg
				125					130					135
Thr	Met	Gly	Ser	Thr	Ile	Arg	Gln	Pro	Lys	Leu	Ser	Asn	Leu	Ser
				140					145					150
Pro	Ser	Val	Ile	Ala	Gln	Thr	Asn	Trp	Lys	Phe	Val	Glu	Gly	Leu
				155					160					165
Leu	Lys	Glu	Cys	Arg	Asn	Lys	Thr	Lys	Arg	Met	Leu	Val	Glu	Lys
				170					175					180
Met	Gly	Arg	Glu	Ala	Val	Glu	Leu	Gly	His	Gly	Glu	Val	Asn	Ile
				185					190					195
Thr	Gly	Val	Glu	Glu	Asn	Thr	Leu	Ile	Ala	Ser	Leu	Cys	Asp	Leu
				200					205					210
Leu	Glu	Arg	Ile	Trp	Ser	His	Gly	Leu	Gln	Val	Lys	Gln	Gly	Lys
				215					220					225
Ser	Ala	Leu	Trp	Ser	His	Leu	Leu	His	Tyr	Gln	Asp	Asn	Arg	Gln
				230					235					240
Arg	Lys	Leu	Thr	Ser	Gly	Ser	Leu	Ser	Thr	Ser	Gly	Ile	Leu	Leu
				245					250					255
Asp	Ser	Glu	Arg	Arg	Lys	Ser	Asp	Ala	Ser	Ser	Leu	Met	Pro	Pro
				260					265					270
Leu	Arg	Ile	Ser	Leu	Ile	Gln	Asp	Met	Arg	His	Ile	Gln	Asn	Ile
				275					280					285
Gly	Glu	Ile	Lys	Thr	Asp	Val	Gly	Lys	Ala	Arg	Ala	Trp	Val	Arg
				290					295					300
Leu	Ser	Met	Glu	Lys	Lys	Leu	Leu	Ser	Arg	His	Leu	Lys	Gln	Leu
				305					310					315
Leu	Ser	Asp	His	Glu	Leu	Thr	Lys	Lys	Leu	Tyr	Lys	Arg	Tyr	Ala
				320					325					330
Phe	Leu	Arg	Cys	Asp	Asp	Glu	Lys	Glu	Gln	Phe	Leu	Tyr	His	Leu
				335					340					345
Leu	Ser	Phe	Asn	Ala	Val	Asp	Tyr	Phe	Cys	Phe	Thr	Asn	Val	Phe
				350					355					360
Thr	Thr	Ile	Leu	Ile	Pro	Tyr	His	Ile	Leu	Ile	Val	Pro	Ser	Lys
				365					370					375
Lys	Leu	Gly	Gly	Ser	Met	Phe	Thr	Ala	Asn	Pro	Trp	Ile	Cys	Ile
				380					385					390
Ser	Gly	Glu	Leu	Gly	Glu	Thr	Gln	Ile	Met	Gln	Ile	Pro	Arg	Asn
				395					400					405
Val	Leu	Glu	Met	Thr	Phe	Glu	Cys	Gln	Asn	Leu	Gly	Lys	Leu	Thr
				410					415					420
Thr	Val	Gln	Ile	Gly	His	Asp	Asn	Ser	Gly	Leu	Tyr	Ala	Lys	Trp
				425					430					435
Leu	Val	Glu	Tyr	Val	Met	Val	Arg	Asn	Glu	Ile	Thr	Gly	His	Thr
				440					445					450
Tyr	Lys	Phe	Pro	Cys	Gly	Arg	Trp	Leu	Gly	Lys	Gly	Met	Asp	Asp
				455					460					465

Gly	Ser	Leu	Glu	Arg	Ile	Leu	Val	Gly	Glu	Leu	Leu	Thr	Ser	Gln	
				470					475					480	
Pro	Glu	Val	Asp	Glu	Arg	Pro	Cys	Arg	Thr	Pro	Pro	Leu	Gln	Gln	
				485					490					495	
Ser	Pro	Ser	Val	Ile	Arg	Arg	Leu	Val	Thr	Ile	Ser	Pro	Asn	Asn	
				500					505					510	
Lys	Pro	Lys	Leu	Asn	Thr	Gly	Gln	Ile	Gln	Glu	Ser	Ile	Gly	Glu	
				515					520					525	
Ala	Val	Asn	Gly	Ile	Val	Lys	His	Phe	His	Lys	Pro	Glu	Lys	Glu	
				530					535					540	
Arg	Gly	Ser	Leu	Thr	Leu	Leu	Leu	Cys	Gly	Glu	Cys	Gly	Leu	Val	
				545					550					555	
Ser	Ala	Leu	Glu	Gln	Ala	Phe	Gln	His	Gly	Phe	Lys	Ser	Pro	Arg	
				560					565					570	
Leu	Phe	Lys	Asn	Val	Phe	Ile	Trp	Asp	Phe	Leu	Glu	Lys	Ala	Gln	
				575					580					585	
Thr	Tyr	Tyr	Glu	Thr	Leu	Glu	Lys	Asn	Glu	Val	Val	Pro	Glu	Glu	
				590					595					600	
Asn	Trp	His	Thr	Arg	Ala	Arg	Asn	Phe	Cys	Arg	Phe	Val	Thr	Ala	
				605					610					615	
Ile	Asn	Asn	Thr	Pro	Arg	Asn	Ile	Gly	Lys	Asp	Gly	Lys	Phe	Gln	
				620					625					630	
Met	Leu	Val	Cys	Leu	Gly	Ala	Arg	Asp	His	Leu	Leu	His	His	Trp	
				635					640					645	
Ile	Ala	Leu	Leu	Ala	Asp	Cys	Pro	Ile	Thr	Ala	His	Met	Tyr	Glu	
				650					655					660	
Asp	Val	Ala	Leu	Ile	Lys	Asp	His	Thr	Leu	Val	Asn	Ser	Leu	Ile	
				665					670					675	
Arg	Val	Leu	Gln	Thr	Leu	Gln	Glu	Phe	Asn	Ile	Thr	Leu	Glu	Thr	
				680					685					690	
Ser	Leu	Val	Lys	Gly	Ile	Asp	Ile								
				695											

<210> 18

<211> 396

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2817769CD1

<400> 18

Met	Pro	Pro	Lys	Lys	Gly	Gly	Asp	Gly	Ile	Lys	Pro	Pro	Pro	Ile	
1				5					10					15	
Ile	Gly	Arg	Phe	Gly	Thr	Ser	Leu	Lys	Ile	Gly	Ile	Val	Gly	Leu	
				20					25					30	
Pro	Asn	Val	Gly	Lys	Ser	Thr	Phe	Phe	Asn	Val	Leu	Thr	Asn	Ser	
				35					40					45	
Gln	Ala	Ser	Ala	Glu	Asn	Phe	Pro	Phe	Cys	Thr	Ile	Asp	Pro	Asn	
				50					55					60	
Glu	Ser	Arg	Val	Pro	Val	Pro	Asp	Glu	Arg	Phe	Asp	Phe	Leu	Cys	
				65					70					75	
Gln	Tyr	His	Lys	Pro	Ala	Ser	Lys	Ile	Pro	Ala	Phe	Leu	Asn	Val	
				80					85					90	

<400> 19
Met Ser Ser Asp Ser Glu Tyr Asp Ser Asp Asp Asp Arg Thr Lys

1	5	10	15
Glu Glu Arg Ala Tyr Asp Lys Ala Lys Arg Arg Ile Glu Lys Arg			
	20	25	30
Arg Leu Glu His Ser Lys Asn Val Asn Thr Glu Lys Leu Arg Ala			
	35	40	45
Pro Ile Ile Cys Val Leu Gly His Val Asp Thr Gly Lys Thr Lys			
	50	55	60
Ile Leu Asp Lys Leu Arg His Thr His Val Gln Asp Gly Glu Ala			
	65	70	75
Gly Gly Ile Thr Gln Gln Ile Gly Ala Thr Asn Val Pro Leu Glu			
	80	85	90
Ala Ile Asn Glu Gln Thr Lys Met Ile Lys Asn Phe Asp Arg Glu			
	95	100	105
Asn Val Arg Ile Pro Gly Met Leu Ile Ile Asp Thr Pro Gly His			
	110	115	120
Glu Ser Phe Ser Asn Leu Arg Asn Arg Gly Ser Ser Leu Cys Asp			
	125	130	135
Ile Ala Ile Leu Val Val Asp Ile Met His Gly Leu Glu Pro Gln			
	140	145	150
Thr Ile Glu Ser Ile Asn Leu Leu Lys Ser Lys Lys Cys Pro Phe			
	155	160	165
Ile Val Ala Leu Asn Lys Ile Asp Arg Leu Tyr Asp Trp Lys Lys			
	170	175	180
Ser Pro Asp Ser Asp Val Ala Ala Thr Leu Lys Lys Gln Lys Lys			
	185	190	195
Asn Thr Lys Asp Glu Phe Glu Glu Arg Ala Lys Ala Ile Ile Val			
	200	205	210
Glu Phe Ala Gln Gln Gly Leu Asn Ala Ala Leu Phe Tyr Glu Asn			
	215	220	225
Lys Asp Pro Arg Thr Phe Val Ser Leu Val Pro Thr Ser Ala His			
	230	235	240
Thr Gly Asp Gly Met Gly Ser Leu Ile Tyr Leu Leu Val Glu Leu			
	245	250	255
Thr Gln Thr Met Leu Ser Lys Arg Leu Ala His Cys Glu Glu Leu			
	260	265	270
Arg Ala Gln Val Met Glu Val Lys Ala Leu Pro Gly Met Gly Thr			
	275	280	285
Thr Ile Asp Val Ile Leu Ile Asn Gly Arg Leu Lys Glu Gly Asp			
	290	295	300
Thr Ile Ile Val Pro Gly Val Glu Gly Pro Ile Val Thr Gln Ile			
	305	310	315
Arg Gly Leu Leu Leu Pro Pro Pro Met Lys Glu Leu Arg Val Lys			
	320	325	330
Asn Gln Tyr Glu Lys His Lys Glu Val Glu Ala Ala Gln Gly Val			
	335	340	345
Lys Ile Leu Gly Lys Asp Leu Glu Lys Thr Leu Ala Gly Leu Pro			
	350	355	360
Leu Leu Val Ala Tyr Lys Glu Asp Glu Ile Pro Val Leu Lys Asp			
	365	370	375
Glu Leu Ile His Glu Leu Lys Gln Thr Leu Asn Ala Ile Lys Leu			
	380	385	390
Glu Glu Lys Gly Val Tyr Val Gln Ala Ser Thr Leu Gly Ser Leu			
	395	400	405
Glu Ala Leu Leu Glu Phe Leu Lys Thr Ser Glu Val Pro Tyr Ala			
	410	415	420

Gly	Ile	Asn	Ile	Gly	Pro	Val	His	Lys	Lys	Asp	Val	Met	Lys	Ala	
				425					430					435	
Ser	Val	Met	Leu	Glu	His	Asp	Pro	Gln	Tyr	Ala	Val	Ile	Leu	Ala	
				440					445					450	
Phe	Asp	Val	Arg	Ile	Glu	Arg	Asp	Ala	Gln	Glu	Met	Ala	Asp	Ser	
				455					460					465	
Leu	Gly	Val	Arg	Ile	Phe	Ser	Ala	Glu	Ile	Ile	Tyr	His	Leu	Phe	
				470					475					480	
Asp	Ala	Phe	Thr	Lys	Tyr	Arg	Gln	Asp	Tyr	Lys	Lys	Gln	Lys	Gln	
				485					490					495	
Glu	Glu	Phe	Lys	His	Ile	Ala	Val	Phe	Pro	Cys	Lys	Ile	Lys	Ile	
				500					505					510	
Leu	Pro	Gln	Tyr	Ile	Phe	Asn	Ser	Arg	Asp	Pro	Ile	Val	Met	Gly	
				515					520					525	
Val	Thr	Val	Glu	Ala	Gly	Gln	Val	Lys	Gln	Gly	Thr	Pro	Met	Cys	
				530					535					540	
Val	Pro	Ser	Lys	Asn	Phe	Val	Asp	Ile	Gly	Ile	Val	Thr	Ser	Ile	
				545					550					555	
Glu	Ile	Asn	His	Lys	Gln	Val	Asp	Val	Ala	Lys	Lys	Gly	Gln	Glu	
				560					565					570	
Val	Cys	Val	Lys	Ile	Glu	Pro	Ile	Pro	Gly	Glu	Ser	Pro	Lys	Met	
				575					580					585	
Phe	Gly	Arg	His	Phe	Glu	Ala	Thr	Asp	Ile	Leu	Val	Ser	Lys	Ile	
				590					595					600	
Ser	Arg	Gln	Ser	Ile	Asp	Ala	Leu	Lys	Asp	Trp	Phe	Arg	Asp	Glu	
				605					610					615	
Met	Gln	Lys	Ser	Asp	Trp	Gln	Leu	Ile	Val	Glu	Leu	Lys	Lys	Val	
				620					625					630	
Phe	Glu	Ile	Ile												

<210> 20

<211> 196

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3421335CD1

<400> 20

Met	Gly	Ser	Val	Asn	Ser	Arg	Gly	His	Lys	Ala	Glu	Ala	Gln	Val	
1				5					10					15	
Val	Met	Met	Gly	Leu	Asp	Ser	Ala	Gly	Lys	Thr	Thr	Leu	Leu	Tyr	
				20					25					30	
Lys	Leu	Lys	Gly	His	Gln	Leu	Val	Glu	Thr	Leu	Pro	Thr	Val	Gly	
				35					40					45	
Phe	Asn	Val	Glu	Pro	Leu	Lys	Ala	Pro	Gly	His	Val	Ser	Leu	Thr	
				50					55					60	
Leu	Trp	Asp	Val	Gly	Gly	Gln	Ala	Pro	Leu	Arg	Ala	Ser	Trp	Lys	
				65					70					75	
Asp	Tyr	Leu	Glu	Gly	Thr	Asp	Ile	Leu	Val	Tyr	Val	Leu	Asp	Ser	
				80					85					90	
Thr	Asp	Glu	Ala	Arg	Leu	Pro	Glu	Ser	Ala	Ala	Glu	Leu	Thr	Glu	
				95					100					105	

Val	Leu	Asn	Asp	Pro	Asn	Met	Ala	Gly	Val	Pro	Phe	Leu	Val	Leu
				110					115					120
Ala	Asn	Lys	Gln	Glu	Ala	Pro	Asp	Ala	Leu	Pro	Leu	Leu	Lys	Ile
				125					130					135
Arg	Asn	Arg	Leu	Ser	Leu	Glu	Arg	Phe	Gln	Asp	His	Cys	Trp	Glu
				140					145					150
Leu	Arg	Gly	Cys	Ser	Ala	Leu	Thr	Gly	Glu	Gly	Leu	Pro	Glu	Ala
				155					160					165
Leu	Gln	Ser	Leu	Trp	Ser	Leu	Leu	Lys	Ser	Arg	Ser	Cys	Met	Cys
				170					175					180
Leu	Gln	Ala	Arg	Ala	His	Gly	Ala	Glu	Arg	Gly	Asp	Ser	Lys	Arg
				185					190					195

Ser

<210> 21

<211> 446

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 605761CD1

<400> 21

Met	Ala	Ala	Arg	Lys	Gly	Arg	Arg	Arg	Thr	Cys	Glu	Thr	Gly	Glu
1				5					10					15
Pro	Met	Glu	Ala	Glu	Ser	Gly	Asp	Thr	Ser	Ser	Glu	Gly	Pro	Ala
				20					25					30
Gln	Val	Tyr	Leu	Pro	Gly	Arg	Gly	Pro	Pro	Leu	Arg	Glu	Gly	Glu
				35					40					45
Glu	Leu	Val	Met	Asp	Glu	Glu	Ala	Tyr	Val	Leu	Tyr	His	Arg	Ala
				50					55					60
Gln	Thr	Gly	Ala	Pro	Cys	Leu	Ser	Phe	Asp	Ile	Val	Arg	Asp	His
				65					70					75
Leu	Gly	Asp	Asn	Arg	Thr	Glu	Leu	Pro	Leu	Thr	Leu	Tyr	Leu	Cys
				80					85					90
Ala	Gly	Thr	Gln	Ala	Glu	Ser	Ala	Gln	Ser	Asn	Arg	Leu	Met	Met
				95					100					105
Leu	Arg	Met	His	Asn	Leu	His	Gly	Thr	Lys	Pro	Pro	Pro	Ser	Glu
				110					115					120
Gly	Ser	Asp	Glu	Glu	Glu	Glu	Glu	Glu	Asp	Glu	Glu	Asp	Glu	Glu
				125					130					135
Glu	Arg	Lys	Pro	Gln	Leu	Glu	Leu	Ala	Met	Val	Pro	His	Tyr	Gly
				140					145					150
Gly	Ile	Asn	Arg	Val	Arg	Val	Ser	Trp	Leu	Gly	Glu	Glu	Pro	Val
				155					160					165
Ala	Gly	Val	Trp	Ser	Glu	Lys	Gly	Gln	Val	Glu	Val	Phe	Ala	Leu
				170					175					180
Arg	Arg	Leu	Leu	Gln	Val	Val	Glu	Glu	Pro	Gln	Ala	Leu	Ala	Ala
				185					190					195
Phe	Leu	Arg	Asp	Glu	Gln	Ala	Gln	Met	Lys	Pro	Ile	Phe	Ser	Phe
				200					205					210
Ala	Gly	His	Met	Gly	Glu	Gly	Phe	Ala	Leu	Asp	Trp	Ser	Pro	Arg
				215					220					225

Val Thr Gly Arg	Leu Leu Thr Gly Asp	Cys Gln Lys Asn Ile His	
	230	235	240
Leu Trp Thr Pro	Thr Asp Gly Gly Ser	Trp His Val Asp Gln Arg	
	245	250	255
Pro Phe Val Gly	His Thr Arg Ser Val	Glu Asp Leu Gln Trp Ser	
	260	265	270
Pro Thr Glu Asn	Thr Val Phe Ala Ser	Cys Ser Ala Asp Ala Ser	
	275	280	285
Ile Arg Ile Trp	Asp Ile Arg Ala Ala	Pro Ser Lys Ala Cys Met	
	290	295	300
Leu Thr Thr Ala	Thr Ala His Asp Gly	Asp Val Asn Val Ile Ser	
	305	310	315
Trp Ser Arg Arg	Glu Pro Phe Leu Leu	Ser Gly Gly Asp Asp Gly	
	320	325	330
Ala Leu Lys Ile	Trp Asp Leu Arg Gln	Phe Lys Ser Gly Ser Pro	
	335	340	345
Val Ala Thr Phe	Lys Gln His Val Ala	Pro Val Thr Ser Val Glu	
	350	355	360
Trp His Pro Gln	Asp Ser Gly Val Phe	Ala Ala Ser Gly Ala Asp	
	365	370	375
His Gln Ile Thr	Gln Trp Asp Leu Ala	Val Glu Arg Asp Pro Glu	
	380	385	390
Ala Gly Asp Val	Glu Ala Asp Pro Gly	Leu Ala Asp Leu Pro Gln	
	395	400	405
Gln Leu Leu Phe	Val His Gln Gly Glu	Thr Glu Leu Lys Glu Leu	
	410	415	420
His Trp His Pro	Gln Cys Pro Gly Leu	Leu Val Ser Thr Ala Leu	
	425	430	435
Ser Gly Phe Thr	Ile Phe Arg Thr Ile	Ser Val	
	440	445	

<210> 22

<211> 265

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 483862CD1

<400> 22

Met Ser Ser Gly	Leu Arg Ala Ala Asp	Phe Pro Arg Trp Lys Arg	
1	5	10	15
His Ile Ser Glu	Gln Leu Arg Arg Arg	Asp Arg Leu Gln Arg Gln	
	20	25	30
Ala Phe Glu Glu	Ile Ile Leu Gln Tyr	Asn Lys Leu Leu Glu Lys	
	35	40	45
Ser Asp Leu His	Ser Val Leu Ala Gln	Lys Leu Gln Ala Glu Lys	
	50	55	60
His Asp Val Pro	Asn Arg His Glu Ile	Ser Pro Gly His Asp Gly	
	65	70	75
Thr Trp Asn Asp	Asn Gln Leu Gln Glu	Met Ala Gln Leu Arg Ile	
	80	85	90
Lys His Gln Glu	Glu Leu Thr Glu Leu	His Lys Lys Arg Gly Glu	

	95		100		105									
Leu	Ala	Gln	Leu	Val	Ile	Asp	Leu	Asn	Asn	Gln	Met	Gln	Arg	Lys
	110								115					120
Asp	Arg	Glu	Met	Gln	Met	Asn	Glu	Ala	Lys	Ile	Ala	Glu	Cys	Leu
	125								130					135
Gln	Thr	Ile	Ser	Asp	Leu	Glu	Thr	Glu	Cys	Leu	Asp	Leu	Arg	Thr
	140								145					150
Lys	Leu	Cys	Asp	Leu	Glu	Arg	Ala	Asn	Gln	Thr	Leu	Lys	Asp	Glu
	155								160					165
Tyr	Asp	Ala	Leu	Gln	Ile	Thr	Phe	Thr	Ala	Leu	Glu	Gly	Lys	Leu
	170								175					180
Arg	Lys	Thr	Thr	Glu	Glu	Asn	Gln	Glu	Leu	Val	Thr	Arg	Trp	Met
	185								190					195
Ala	Glu	Lys	Ala	Gln	Glu	Ala	Asn	Arg	Leu	Asn	Ala	Glu	Asn	Glu
	200								205					210
Lys	Asp	Ser	Arg	Arg	Arg	Gln	Ala	Arg	Leu	Gln	Lys	Glu	Leu	Ala
	215								220					225
Glu	Ala	Ala	Lys	Glu	Pro	Leu	Pro	Val	Glu	Gln	Asp	Asp	Asp	Ile
	230								235					240
Glu	Val	Ile	Val	Asp	Glu	Thr	Ser	Asp	His	Thr	Glu	Glu	Thr	Ser
	245								250					255
Pro	Val	Arg	Ala	Ile	Ser	Arg	Ala	Ala	Thr					
	260								265					

<210> 23

<211> 185

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1256777CD1

<400> 23

Met	Leu	Lys	Ala	Lys	Ile	Leu	Phe	Val	Gly	Pro	Cys	Glu	Ser	Gly
1				5					10					15
Lys	Thr	Val	Leu	Ala	Asn	Phe	Leu	Thr	Glu	Ser	Ser	Asp	Ile	Thr
				20					25					30
Glu	Tyr	Ser	Pro	Thr	Gln	Gly	Val	Arg	Ile	Leu	Glu	Phe	Glu	Asn
				35					40					45
Pro	His	Val	Thr	Ser	Asn	Asn	Lys	Gly	Thr	Gly	Cys	Glu	Phe	Glu
				50					55					60
Leu	Trp	Asp	Cys	Gly	Gly	Asp	Ala	Lys	Phe	Glu	Ser	Cys	Trp	Pro
				65					70					75
Ala	Leu	Met	Lys	Asp	Ala	His	Gly	Val	Val	Ile	Val	Phe	Asn	Ala
				80					85					90
Asp	Ile	Pro	Ser	His	Arg	Lys	Glu	Met	Glu	Met	Trp	Tyr	Ser	Cys
				95					100					105
Phe	Val	Gln	Gln	Pro	Ser	Leu	Gln	Asp	Thr	Gln	Cys	Met	Leu	Ile
				110					115					120
Ala	His	His	Lys	Pro	Gly	Ser	Gly	Asp	Asp	Lys	Gly	Ser	Leu	Ser
				125					130					135
Leu	Ser	Pro	Pro	Leu	Asn	Lys	Leu	Lys	Leu	Val	His	Ser	Asn	Leu
				140					145					150

Glu	Asp	Asp	Pro	Glu	Glu	Ile	Arg	Met	Glu	Phe	Ile	Lys	Tyr	Leu
				155					160					165
Lys	Ser	Ile	Ile	Asn	Ser	Met	Ser	Glu	Ser	Arg	Asp	Arg	Glu	Glu
				170					175					180
Met	Ser	Ile	Met	Thr										
				185										

<210> 24

<211> 554

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2198779CD1

<400> 24

Met	Gly	Ser	Arg	Asn	Ser	Ser	Ser	Ala	Gly	Ser	Gly	Ser	Gly	Asp
1				5					10					15
Pro	Ser	Glu	Gly	Leu	Pro	Arg	Arg	Gly	Ala	Gly	Leu	Arg	Arg	Ser
				20					25					30
Glu	Glu	Glu	Glu	Glu	Glu	Asp	Glu	Asp	Val	Asp	Leu	Ala	Gln	Val
				35					40					45
Leu	Ala	Tyr	Leu	Leu	Arg	Arg	Gly	Gln	Val	Arg	Leu	Val	Gln	Gly
				50					55					60
Gly	Gly	Ala	Ala	Asn	Leu	Gln	Phe	Ile	Gln	Ala	Leu	Leu	Asp	Ser
				65					70					75
Glu	Glu	Glu	Asn	Asp	Arg	Ala	Trp	Asp	Gly	Arg	Leu	Gly	Asp	Arg
				80					85					90
Tyr	Asn	Pro	Pro	Val	Asp	Ala	Thr	Pro	Asp	Thr	Arg	Glu	Leu	Glu
				95					100					105
Phe	Asn	Glu	Ile	Lys	Thr	Gln	Val	Glu	Leu	Ala	Thr	Gly	Gln	Leu
				110					115					120
Gly	Leu	Arg	Arg	Ala	Ala	Gln	Lys	His	Ser	Phe	Pro	Arg	Met	Leu
				125					130					135
His	Gln	Arg	Glu	Arg	Gly	Leu	Cys	His	Arg	Gly	Ser	Phe	Ser	Leu
				140					145					150
Gly	Glu	Gln	Ser	Arg	Val	Ile	Ser	His	Phe	Leu	Pro	Asn	Asp	Leu
				155					160					165
Gly	Phe	Thr	Asp	Ser	Tyr	Ser	Gln	Lys	Ala	Phe	Cys	Gly	Ile	Tyr
				170					175					180
Ser	Lys	Asp	Gly	Gln	Ile	Phe	Met	Ser	Ala	Cys	Gln	Asp	Gln	Thr
				185					190					195
Ile	Arg	Leu	Tyr	Asp	Cys	Arg	Tyr	Gly	Arg	Phe	Arg	Lys	Phe	Lys
				200					205					210
Ser	Ile	Lys	Ala	Arg	Asp	Val	Gly	Trp	Ser	Val	Leu	Asp	Val	Ala
				215					220					225
Phe	Thr	Pro	Asp	Gly	Asn	His	Phe	Leu	Tyr	Ser	Ser	Trp	Ser	Asp
				230					235					240
Tyr	Ile	His	Ile	Cys	Asn	Ile	Tyr	Gly	Glu	Gly	Asp	Thr	His	Thr
				245					250					255
Ala	Leu	Asp	Leu	Arg	Pro	Asp	Glu	Arg	Arg	Phe	Ala	Val	Phe	Ser
				260					265					270
Ile	Ala	Val	Ser	Ser	Asp	Gly	Arg	Glu	Val	Leu	Gly	Gly	Ala	Asn
				275					280					285

Asp Gly Cys Leu Tyr Val Phe Asp Arg Glu Gln Asn Arg Arg Thr	290	295	300
Leu Gln Ile Glu Ser His Glu Asp Asp Val Asn Ala Val Ala Phe	305	310	315
Ala Asp Ile Ser Ser Gln Ile Leu Phe Ser Gly Gly Asp Asp Ala	320	325	330
Ile Cys Lys Val Trp Asp Arg Arg Thr Met Arg Glu Asp Asp Pro	335	340	345
Lys Pro Val Gly Ala Leu Ala Gly His Gln Asp Gly Ile Thr Phe	350	355	360
Ile Asp Ser Lys Gly Asp Ala Arg Tyr Leu Ile Ser Asn Ser Lys	365	370	375
Asp Gln Thr Ile Lys Leu Trp Asp Ile Arg Arg Phe Ser Ser Arg	380	385	390
Glu Gly Met Glu Ala Ser Arg Gln Ala Ala Thr Gln Gln Asn Trp	395	400	405
Asp Tyr Arg Trp Gln Gln Val Pro Lys Lys Gly Phe Thr Leu His	410	415	420
Pro Tyr Pro Ala Trp Arg Lys Leu Lys Leu Pro Gly Asp Ser Ser	425	430	435
Leu Met Thr Tyr Arg Gly His Gly Val Leu His Thr Leu Ile Arg	440	445	450
Cys Arg Phe Ser Pro Ile His Ser Thr Gly Gln Gln Phe Ile Tyr	455	460	465
Ser Gly Cys Ser Thr Gly Lys Val Val Val Tyr Asp Leu Leu Ser	470	475	480
Gly His Ile Val Lys Lys Leu Thr Asn His Lys Ala Cys Val Arg	485	490	495
Asp Val Ser Trp His Pro Phe Glu Glu Lys Ile Val Ser Ser Ser	500	505	510
Trp Asp Gly Asn Leu Arg Leu Trp Gln Tyr Arg Gln Ala Glu Tyr	515	520	525
Phe Gln Asp Asp Met Pro Glu Ser Glu Glu Cys Ala Ser Ala Pro	530	535	540
Ala Pro Val Pro Gln Ser Ser Thr Pro Phe Ser Ser Pro Gln	545	550	

<210> 25

<211> 434

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2226116CD1

<400> 25

Met Arg Pro Ser Ser Ser Val Ser Val Ser Cys Pro Ala Leu Asn	1	5	10	15
Gln Val Ser His Phe Ala Asn Leu Thr Ser Val Gly Ala Met Ala	20	25	30	
Pro Ala Arg Cys Phe Ser Ala Arg Leu Arg Thr Val Phe Gln Gly	35	40	45	
Val Gly His Trp Ala Leu Ser Thr Trp Ala Gly Leu Lys Pro Ser				

Arg	Leu	Leu	Pro	Gln	Arg	Ala	Ser	Pro	Arg	Leu	Leu	Ser	Val	Gly	50	55	60
															65	70	75
Arg	Ala	Asp	Leu	Ala	Lys	His	Gln	Glu	Leu	Pro	Gly	Lys	Lys	Leu			
															80	85	90
Leu	Ser	Glu	Lys	Lys	Leu	Lys	Arg	Tyr	Phe	Val	Asp	Tyr	Arg	Arg			
															95	100	105
Val	Leu	Val	Cys	Gly	Gly	Asn	Gly	Gly	Ala	Gly	Ala	Ser	Cys	Phe			
															110	115	120
His	Ser	Glu	Pro	Arg	Lys	Glu	Phe	Gly	Gly	Pro	Asp	Gly	Gly	Asp			
															125	130	135
Gly	Gly	Asn	Gly	Gly	His	Val	Ile	Leu	Arg	Val	Asp	Gln	Gln	Val			
															140	145	150
Lys	Ser	Leu	Ser	Ser	Val	Leu	Ser	Arg	Tyr	Gln	Gly	Phe	Ser	Gly			
															155	160	165
Glu	Asp	Gly	Gly	Ser	Lys	Asn	Cys	Phe	Gly	Arg	Ser	Gly	Ala	Val			
															170	175	180
Leu	Tyr	Ile	Arg	Val	Pro	Val	Gly	Thr	Leu	Val	Lys	Glu	Gly	Gly			
															185	190	195
Arg	Val	Val	Ala	Asp	Leu	Ser	Cys	Val	Gly	Asp	Glu	Tyr	Ile	Ala			
															200	205	210
Ala	Leu	Gly	Gly	Ala	Gly	Gly	Lys	Gly	Asn	Arg	Phe	Phe	Leu	Ala			
															215	220	225
Asn	Asn	Asn	Arg	Ala	Pro	Val	Thr	Cys	Thr	Pro	Gly	Gln	Pro	Gly			
															230	235	240
Gln	Gln	Arg	Val	Leu	His	Leu	Glu	Leu	Lys	Thr	Val	Ala	His	Ala			
															245	250	255
Gly	Met	Val	Gly	Phe	Pro	Asn	Ala	Gly	Lys	Ser	Ser	Leu	Leu	Arg			
															260	265	270
Ala	Ile	Ser	Asn	Ala	Arg	Pro	Ala	Val	Ala	Ser	Tyr	Pro	Phe	Thr			
															275	280	285
Thr	Leu	Lys	Pro	His	Val	Gly	Ile	Val	His	Tyr	Glu	Gly	His	Leu			
															290	295	300
Gln	Ile	Ala	Val	Ala	Asp	Ile	Pro	Gly	Ile	Ile	Arg	Gly	Ala	His			
															305	310	315
Gln	Asn	Arg	Gly	Leu	Gly	Ser	Ala	Phe	Leu	Arg	His	Ile	Glu	Arg			
															320	325	330
Cys	Arg	Phe	Leu	Leu	Phe	Val	Val	Asp	Leu	Ser	Gln	Pro	Glu	Pro			
															335	340	345
Trp	Thr	Gln	Val	Asp	Asp	Leu	Lys	Tyr	Glu	Leu	Glu	Met	Tyr	Glu			
															350	355	360
Lys	Gly	Leu	Ser	Ala	Arg	Pro	His	Ala	Ile	Val	Ala	Asn	Lys	Ile			
															365	370	375
Asp	Leu	Pro	Glu	Ala	Gln	Ala	Asn	Leu	Ser	Gln	Leu	Arg	Asp	His			
															380	385	390
Leu	Gly	Gln	Glu	Val	Ile	Val	Leu	Ser	Ala	Leu	Thr	Gly	Glu	Asn			
															395	400	405
Leu	Glu	Gln	Leu	Leu	Leu	His	Leu	Lys	Val	Leu	Tyr	Asp	Ala	Tyr			
															410	415	420
Ala	Glu	Ala	Glu	Leu	Gly	Gln	Gly	Arg	Gln	Pro	Leu	Arg	Trp				
															425	430	

<210> 26

<211> 826

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2504472CD1

<400> 26

Met	Val	Ala	Pro	Val	Leu	Glu	Thr	Ser	His	Val	Phe	Cys	Cys	Pro
1				5					10					15
Asn	Arg	Val	Arg	Gly	Val	Leu	Asn	Trp	Ser	Ser	Gly	Pro	Arg	Gly
				20					25					30
Leu	Leu	Ala	Phe	Gly	Thr	Ser	Cys	Ser	Val	Val	Leu	Tyr	Asp	Pro
				35					40					45
Leu	Lys	Arg	Val	Val	Val	Thr	Asn	Leu	Asn	Gly	His	Thr	Ala	Arg
				50					55					60
Val	Asn	Cys	Ile	Gln	Trp	Ile	Cys	Lys	Gln	Asp	Gly	Ser	Pro	Ser
				65					70					75
Thr	Glu	Leu	Val	Ser	Gly	Gly	Ser	Asp	Asn	Gln	Val	Ile	His	Trp
				80					85					90
Glu	Ile	Glu	Asp	Asn	Gln	Leu	Leu	Lys	Ala	Val	His	Leu	Gln	Gly
				95					100					105
His	Glu	Gly	Pro	Val	Tyr	Ala	Val	His	Ala	Val	Tyr	Gln	Arg	Arg
				110					115					120
Thr	Ser	Asp	Pro	Ala	Leu	Cys	Thr	Leu	Ile	Val	Ser	Ala	Ala	Ala
				125					130					135
Asp	Ser	Ala	Val	Arg	Leu	Trp	Ser	Lys	Lys	Gly	Pro	Glu	Val	Met
				140					145					150
Cys	Leu	Gln	Thr	Leu	Asn	Phe	Gly	Asn	Gly	Phe	Ala	Leu	Ala	Leu
				155					160					165
Cys	Leu	Ser	Phe	Leu	Pro	Asn	Thr	Asp	Val	Pro	Ile	Leu	Ala	Cys
				170					175					180
Gly	Asn	Asp	Asp	Cys	Arg	Ile	His	Ile	Phe	Ala	Gln	Gln	Asn	Asp
				185					190					195
Gln	Phe	Gln	Lys	Val	Leu	Ser	Leu	Cys	Gly	His	Glu	Asp	Trp	Ile
				200					205					210
Arg	Gly	Val	Glu	Trp	Ala	Ala	Phe	Gly	Arg	Asp	Leu	Phe	Leu	Ala
				215					220					225
Ser	Cys	Ser	Gln	Asp	Cys	Leu	Ile	Arg	Ile	Trp	Lys	Leu	Tyr	Ile
				230					235					240
Lys	Ser	Thr	Ser	Leu	Glu	Thr	Gln	Asp	Asp	Asp	Asn	Ile	Arg	Leu
				245					250					255
Lys	Glu	Asn	Thr	Phe	Thr	Ile	Glu	Asn	Glu	Ser	Val	Lys	Ile	Ala
				260					265					270
Phe	Ala	Val	Thr	Leu	Glu	Thr	Val	Leu	Ala	Gly	His	Glu	Asn	Trp
				275					280					285
Val	Asn	Ala	Val	His	Trp	Gln	Pro	Val	Phe	Tyr	Lys	Asp	Gly	Val
				290					295					300
Leu	Gln	Gln	Pro	Val	Arg	Leu	Leu	Ser	Ala	Ser	Met	Asp	Lys	Thr
				305					310					315
Met	Ile	Leu	Trp	Ala	Pro	Asp	Glu	Glu	Ser	Gly	Val	Trp	Leu	Glu
				320					325					330
Gln	Val	Arg	Val	Gly	Glu	Val	Gly	Gly	Asn	Thr	Leu	Gly	Phe	Tyr
				335					340					345
Asp	Cys	Gln	Phe	Asn	Glu	Asp	Gly	Ser	Met	Ile	Ile	Ala	His	Ala

	350		355		360
Phe His Gly Ala	Leu His Leu Trp Lys	Gln Asn Thr Val Asn	Pro		
	365		370		375
Arg Glu Trp Thr	Pro Glu Ile Val Ile	Ser Gly His Phe Asp	Gly		
	380		385		390
Val Gln Asp Leu	Val Trp Asp Pro Glu	Gly Glu Phe Ile Ile	Thr		
	395		400		405
Val Gly Thr Asp	Gln Thr Thr Arg Leu	Phe Ala Pro Trp Lys	Arg		
	410		415		420
Lys Asp Gln Ser	Gln Val Thr Trp His	Glu Ile Ala Arg Pro	Gln		
	425		430		435
Ile His Gly Tyr	Asp Leu Lys Cys Leu	Ala Met Ile Asn Arg	Phe		
	440		445		450
Gln Phe Val Ser	Gly Ala Asp Glu Lys	Val Leu Arg Val Phe	Ser		
	455		460		465
Ala Pro Arg Asn	Phe Val Glu Asn Phe	Cys Ala Ile Thr Gly	Gln		
	470		475		480
Ser Leu Asn His	Val Leu Cys Asn Gln	Asp Ser Asp Leu Pro	Glu		
	485		490		495
Gly Ala Thr Val	Pro Ala Leu Gly Leu	Ser Asn Lys Ala Val	Phe		
	500		505		510
Gln Gly Asp Ile	Ala Ser Gln Pro Ser	Asp Glu Glu Glu Leu	Leu		
	515		520		525
Thr Ser Thr Gly	Phe Glu Tyr Gln Gln	Val Ala Phe Gln Pro	Ser		
	530		535		540
Ile Leu Thr Glu	Pro Pro Thr Glu Asp	His Leu Leu Gln Asn	Thr		
	545		550		555
Leu Trp Pro Glu	Val Gln Lys Leu Tyr	Gly His Gly Tyr Glu	Ile		
	560		565		570
Phe Cys Val Thr	Cys Asn Ser Ser Lys	Thr Leu Leu Ala Ser	Ala		
	575		580		585
Cys Lys Ala Ala	Lys Lys Glu His Ala	Ala Ile Ile Leu Trp	Asn		
	590		595		600
Thr Thr Ser Trp	Lys Gln Val Gln Asn	Leu Val Phe His Ser	Leu		
	605		610		615
Thr Val Thr Gln	Met Ala Phe Ser Pro	Asn Glu Lys Phe Leu	Leu		
	620		625		630
Ala Val Ser Arg	Asp Arg Thr Trp Ser	Leu Trp Lys Lys Gln	Asp		
	635		640		645
Thr Ile Ser Pro	Glu Phe Glu Pro Val	Phe Ser Leu Phe Ala	Phe		
	650		655		660
Thr Asn Lys Ile	Thr Ser Val His Ser	Arg Ile Ile Trp Ser	Cys		
	665		670		675
Asp Trp Ser Pro	Asp Ser Lys Tyr Phe	Phe Thr Gly Ser Arg	Asp		
	680		685		690
Lys Lys Val Val	Val Trp Gly Glu Cys	Asp Ser Thr Asp Asp	Cys		
	695		700		705
Ile Glu His Asn	Ile Gly Pro Cys Ser	Ser Val Leu Asp Val	Gly		
	710		715		720
Gly Ala Val Thr	Ala Val Ser Val Cys	Pro Val Leu His Pro	Ser		
	725		730		735
Gln Arg Tyr Val	Val Ala Val Gly Leu	Glu Cys Gly Lys Ile	Cys		
	740		745		750
Leu Tyr Thr Trp	Lys Lys Thr Asp Gln	Val Pro Glu Ile Asn	Asp		
	755		760		765

Trp	Thr	His	Cys	Val	Glu	Thr	Ser	Gln	Ser	Gln	Ser	His	Thr	Leu
				770					775					780
Ala	Ile	Arg	Lys	Leu	Cys	Trp	Lys	Asn	Cys	Ser	Gly	Lys	Thr	Glu
				785					790					795
Gln	Lys	Glu	Ala	Glu	Gly	Ala	Glu	Trp	Leu	His	Phe	Ala	Ser	Cys
				800					805					810
Gly	Glu	Asp	His	Thr	Val	Lys	Ile	His	Arg	Val	Asn	Lys	Cys	Ala
				815					820					825

Leu

<210> 27

<211> 618

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3029920CD1

<400> 27

Met	Lys	Lys	Asp	Val	Arg	Ile	Leu	Leu	Val	Gly	Glu	Pro	Arg	Val
1				5					10					15
Gly	Lys	Thr	Ser	Leu	Ile	Met	Ser	Leu	Val	Ser	Glu	Glu	Phe	Pro
				20					25					30
Glu	Glu	Val	Pro	Pro	Arg	Ala	Glu	Glu	Ile	Thr	Ile	Pro	Ala	Asp
				35					40					45
Val	Thr	Pro	Glu	Arg	Val	Pro	Thr	His	Ile	Val	Asp	Tyr	Ser	Glu
				50					55					60
Ala	Glu	Gln	Ser	Asp	Glu	Gln	Leu	His	Gln	Glu	Ile	Ser	Gln	Ala
				65					70					75
Asn	Val	Ile	Cys	Ile	Val	Tyr	Ala	Val	Asn	Asn	Lys	His	Ser	Ile
				80					85					90
Asp	Lys	Val	Thr	Ser	Arg	Trp	Ile	Pro	Leu	Ile	Asn	Glu	Arg	Thr
				95					100					105
Asp	Lys	Asp	Ser	Arg	Leu	Pro	Leu	Ile	Leu	Val	Gly	Asn	Lys	Ser
				110					115					120
Asp	Leu	Val	Glu	Tyr	Ser	Ser	Met	Glu	Thr	Ile	Leu	Pro	Ile	Met
				125					130					135
Asn	Gln	Tyr	Thr	Glu	Ile	Glu	Thr	Cys	Val	Glu	Cys	Ser	Ala	Lys
				140					145					150
Asn	Leu	Lys	Asn	Ile	Ser	Glu	Leu	Phe	Tyr	Tyr	Ala	Gln	Lys	Ala
				155					160					165
Val	Leu	His	Pro	Thr	Gly	Pro	Leu	Tyr	Cys	Pro	Glu	Glu	Lys	Glu
				170					175					180
Met	Lys	Pro	Ala	Cys	Ile	Lys	Ala	Leu	Thr	Arg	Ile	Phe	Lys	Ile
				185					190					195
Ser	Asp	Gln	Asp	Asn	Asp	Gly	Thr	Leu	Asn	Asp	Ala	Glu	Leu	Asn
				200					205					210
Phe	Phe	Gln	Arg	Ile	Cys	Phe	Asn	Thr	Pro	Leu	Ala	Pro	Gln	Ala
				215					220					225
Leu	Glu	Asp	Val	Lys	Asn	Val	Val	Arg	Lys	His	Ile	Ser	Asp	Gly
				230					235					240
Val	Ala	Asp	Ser	Gly	Leu	Thr	Leu	Lys	Gly	Phe	Leu	Phe	Leu	His
				245					250					255

Thr	Leu	Phe	Ile	Gln	Arg	Gly	Arg	His	Glu	Thr	Thr	Trp	Thr	Val
				260					265					270
Leu	Arg	Arg	Phe	Gly	Tyr	Asp	Asp	Asp	Leu	Asp	Leu	Thr	Pro	Glu
				275					280					285
Tyr	Leu	Phe	Pro	Leu	Leu	Lys	Ile	Pro	Pro	Asp	Cys	Thr	Thr	Glu
				290					295					300
Leu	Asn	His	His	Ala	Tyr	Leu	Phe	Leu	Gln	Ser	Thr	Phe	Asp	Lys
				305					310					315
His	Asp	Leu	Asp	Arg	Asp	Cys	Ala	Leu	Ser	Pro	Asp	Glu	Leu	Lys
				320					325					330
Asp	Leu	Phe	Lys	Val	Phe	Pro	Tyr	Ile	Pro	Trp	Gly	Pro	Asp	Val
				335					340					345
Asn	Asn	Thr	Val	Cys	Thr	Asn	Glu	Arg	Gly	Trp	Ile	Thr	Tyr	Gln
				350					355					360
Gly	Phe	Leu	Ser	Gln	Trp	Thr	Leu	Thr	Thr	Tyr	Leu	Asp	Val	Gln
				365					370					375
Arg	Cys	Leu	Glu	Tyr	Leu	Gly	Tyr	Leu	Gly	Tyr	Ser	Ile	Leu	Thr
				380					385					390
Glu	Gln	Glu	Ser	Gln	Ala	Ser	Ala	Val	Thr	Val	Thr	Arg	Asp	Lys
				395					400					405
Lys	Ile	Asp	Leu	Gln	Lys	Lys	Gln	Thr	Gln	Arg	Asn	Val	Phe	Arg
				410					415					420
Cys	Asn	Val	Ile	Gly	Val	Lys	Asn	Cys	Gly	Lys	Ser	Gly	Val	Leu
				425					430					435
Gln	Ala	Leu	Leu	Gly	Arg	Asn	Leu	Met	Arg	Gln	Lys	Lys	Ile	Arg
				440					445					450
Glu	Asp	His	Lys	Ser	Tyr	Tyr	Ala	Ile	Asn	Thr	Val	Tyr	Val	Tyr
				455					460					465
Gly	Gln	Glu	Lys	Tyr	Leu	Leu	Leu	His	Asp	Ile	Ser	Glu	Ser	Glu
				470					475					480
Phe	Leu	Thr	Glu	Ala	Glu	Ile	Ile	Cys	Asp	Val	Val	Cys	Leu	Val
				485					490					495
Tyr	Asp	Val	Ser	Asn	Pro	Lys	Ser	Phe	Glu	Tyr	Cys	Ala	Arg	Ile
				500					505					510
Phe	Lys	Gln	His	Phe	Met	Asp	Ser	Arg	Ile	Pro	Cys	Leu	Ile	Val
				515					520					525
Ala	Ala	Lys	Ser	Asp	Leu	His	Glu	Val	Lys	Gln	Glu	Tyr	Ser	Ile
				530					535					540
Ser	Pro	Thr	Asp	Phe	Cys	Arg	Lys	His	Lys	Met	Pro	Pro	Pro	Gln
				545					550					555
Ala	Phe	Thr	Cys	Asn	Thr	Ala	Asp	Ala	Pro	Ser	Lys	Asp	Ile	Phe
				560					565					570
Val	Lys	Leu	Thr	Thr	Met	Ala	Met	Tyr	Pro	His	Val	Thr	Gln	Ala
				575					580					585
Asp	Leu	Lys	Ser	Ser	Thr	Phe	Trp	Leu	Arg	Ala	Ser	Phe	Gly	Ala
				590					595					600
Thr	Val	Phe	Ala	Val	Leu	Gly	Phe	Ala	Met	Tyr	Lys	Ala	Leu	Leu
				605					610					615
Lys	Gln	Arg												

<210> 28

<211> 596

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3332415CD1

<400> 28

Met	Glu	Pro	Glu	Leu	Asp	Ala	Gln	Lys	Gln	Pro	Arg	Pro	Arg	Arg	1	5	10	15
Arg	Ser	Arg	Arg	Ala	Ser	Gly	Leu	Ser	Thr	Glu	Gly	Ala	Thr	Gly	20	25	30	
Pro	Ser	Ala	Asp	Thr	Ser	Gly	Ser	Glu	Leu	Asp	Gly	Arg	Cys	Ser	35	40	45	
Leu	Arg	Arg	Gly	Ser	Ser	Phe	Thr	Phe	Leu	Thr	Pro	Gly	Pro	Asn	50	55	60	
Trp	Asp	Phe	Thr	Leu	Lys	Arg	Lys	Arg	Arg	Glu	Lys	Asp	Asp	Asp	65	70	75	
Val	Val	Ser	Leu	Ser	Ser	Leu	Asp	Leu	Lys	Glu	Pro	Ser	Asn	Lys	80	85	90	
Arg	Val	Arg	Pro	Leu	Ala	Arg	Val	Thr	Ser	Leu	Ala	Asn	Leu	Ile	95	100	105	
Ser	Pro	Val	Arg	Asn	Gly	Ala	Val	Arg	Arg	Phe	Gly	Gln	Thr	Ile	110	115	120	
Gln	Ser	Phe	Thr	Leu	Arg	Gly	Asp	His	Arg	Ser	Pro	Ala	Ser	Ala	125	130	135	
Gln	Lys	Phe	Ser	Ser	Arg	Ser	Thr	Val	Pro	Thr	Pro	Ala	Lys	Arg	140	145	150	
Arg	Ser	Ser	Ala	Leu	Trp	Ser	Glu	Met	Leu	Asp	Ile	Thr	Met	Lys	155	160	165	
Glu	Ser	Leu	Thr	Thr	Arg	Glu	Ile	Arg	Arg	Gln	Glu	Ala	Ile	Tyr	170	175	180	
Glu	Met	Ser	Arg	Gly	Glu	Gln	Asp	Leu	Ile	Glu	Asp	Leu	Lys	Leu	185	190	195	
Ala	Arg	Lys	Ala	Tyr	His	Asp	Pro	Met	Leu	Lys	Leu	Ser	Ile	Met	200	205	210	
Ser	Glu	Glu	Glu	Leu	Thr	His	Ile	Phe	Gly	Asp	Leu	Asp	Ser	Tyr	215	220	225	
Ile	Pro	Leu	His	Glu	Asp	Leu	Leu	Thr	Arg	Ile	Gly	Glu	Ala	Thr	230	235	240	
Lys	Pro	Asp	Gly	Thr	Val	Glu	Gln	Ile	Gly	His	Ile	Leu	Val	Ser	245	250	255	
Trp	Leu	Pro	Arg	Leu	Asn	Ala	Tyr	Arg	Gly	Tyr	Cys	Ser	Asn	Gln	260	265	270	
Leu	Ala	Ala	Lys	Ala	Leu	Leu	Asp	Gln	Lys	Lys	Gln	Asp	Pro	Arg	275	280	285	
Val	Gln	Asp	Phe	Leu	Gln	Arg	Cys	Leu	Glu	Ser	Pro	Phe	Ser	Arg	290	295	300	
Lys	Leu	Asp	Leu	Trp	Ser	Phe	Leu	Asp	Ile	Pro	Arg	Ser	Arg	Leu	305	310	315	
Val	Lys	Tyr	Pro	Leu	Leu	Leu	Lys	Glu	Ile	Leu	Lys	His	Thr	Pro	320	325	330	
Lys	Glu	His	Pro	Asp	Val	Gln	Leu	Leu	Glu	Asp	Ala	Ile	Leu	Ile	335	340	345	
Ile	Gln	Gly	Val	Leu	Ser	Asp	Ile	Asn	Leu	Lys	Lys	Gly	Glu	Ser	350	355	360	
Glu	Cys	Gln	Tyr	Tyr	Ile	Asp	Lys	Leu	Glu	Tyr	Leu	Asp	Glu	Lys	365	370	375	

Gln	Arg	Asp	Pro	Arg	Ile	Glu	Ala	Ser	Lys	Val	Leu	Leu	Cys	His	
				380					385					390	
Gly	Glu	Leu	Arg	Ser	Lys	Ser	Gly	His	Lys	Leu	Tyr	Ile	Phe	Leu	
				395					400					405	
Phe	Gln	Asp	Ile	Leu	Val	Leu	Thr	Arg	Pro	Val	Thr	Arg	Asn	Glu	
				410					415					420	
Arg	His	Ser	Tyr	Gln	Val	Tyr	Arg	Gln	Pro	Ile	Pro	Val	Gln	Glu	
				425					430					435	
Leu	Val	Leu	Glu	Asp	Leu	Gln	Asp	Gly	Asp	Val	Arg	Met	Gly	Gly	
				440					445					450	
Ser	Phe	Arg	Gly	Ala	Phe	Ser	Asn	Ser	Glu	Lys	Ala	Lys	Asn	Ile	
				455					460					465	
Phe	Arg	Ile	Arg	Phe	His	Asp	Pro	Ser	Pro	Ala	Gln	Ser	His	Thr	
				470					475					480	
Leu	Gln	Ala	Asn	Asp	Val	Phe	His	Lys	Gln	Gln	Trp	Phe	Asn	Cys	
				485					490					495	
Ile	Arg	Ala	Ala	Ile	Ala	Pro	Phe	Gln	Ser	Ala	Gly	Ser	Pro	Pro	
				500					505					510	
Glu	Leu	Gln	Gly	Leu	Pro	Glu	Leu	His	Glu	Glu	Cys	Glu	Gly	Asn	
				515					520					525	
His	Pro	Ser	Ala	Arg	Lys	Leu	Thr	Ala	Gln	Arg	Arg	Ala	Ser	Thr	
				530					535					540	
Val	Ser	Ser	Val	Thr	Gln	Val	Glu	Val	Asp	Glu	Asn	Ala	Tyr	Arg	
				545					550					555	
Cys	Gly	Ser	Gly	Met	Gln	Met	Ala	Glu	Asp	Ser	Lys	Ser	Leu	Lys	
				560					565					570	
Thr	His	Gln	Thr	Gln	Pro	Gly	Ile	Arg	Arg	Ala	Arg	Asp	Lys	Ala	
				575					580					585	
Leu	Ser	Gly	Gly	Lys	Arg	Lys	Glu	Thr	Leu	Val					
				590					595						

<210> 29

<211> 589

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4031536CD1

<400> 29

Met	Ser	Lys	Pro	Gly	Lys	Pro	Thr	Leu	Asn	His	Gly	Leu	Val	Pro	
1				5					10					15	
Val	Asp	Leu	Lys	Ser	Ala	Lys	Glu	Pro	Leu	Pro	His	Gln	Thr	Val	
				20					25					30	
Met	Arg	Ile	Phe	Ser	Ile	Ser	Ile	Ile	Ala	Gln	Gly	Leu	Pro	Phe	
				35					40					45	
Cys	Arg	Arg	Arg	Met	Lys	Arg	Lys	Leu	Asp	His	Gly	Ser	Glu	Val	
				50					55					60	
Arg	Ser	Phe	Ser	Leu	Gly	Lys	Lys	Pro	Cys	Lys	Val	Ser	Glu	Tyr	
				65					70					75	
Thr	Ser	Thr	Thr	Gly	Leu	Val	Pro	Cys	Ser	Ala	Thr	Pro	Thr	Thr	
				80					85					90	
Phe	Gly	Asp	Leu	Arg	Ala	Ala	Asn	Gly	Gln	Gly	Gln	Gln	Arg	Arg	

	95	100	105
Arg Ile Thr Ser Val Gln Pro Pro Thr Gly Leu Gln Glu Trp Leu			
	110	115	120
Lys Met Phe Gln Ser Trp Ser Gly Pro Glu Lys Leu Leu Ala Leu			
	125	130	135
Asp Glu Leu Ile Asp Ser Cys Glu Pro Thr Gln Val Lys His Met			
	140	145	150
Met Gln Val Ile Glu Pro Gln Phe Gln Arg Asp Phe Ile Ser Leu			
	155	160	165
Leu Pro Lys Glu Leu Ala Leu Tyr Val Leu Ser Phe Leu Glu Pro			
	170	175	180
Lys Asp Leu Leu Gln Ala Ala Gln Thr Cys Arg Tyr Trp Arg Ile			
	185	190	195
Leu Ala Glu Asp Asn Leu Leu Trp Arg Glu Lys Cys Lys Glu Glu			
	200	205	210
Gly Ile Asp Glu Pro Leu His Ile Lys Arg Arg Lys Val Ile Lys			
	215	220	225
Pro Gly Phe Ile His Ser Pro Trp Lys Ser Ala Tyr Ile Arg Gln			
	230	235	240
His Arg Ile Asp Thr Asn Trp Arg Arg Gly Glu Leu Lys Ser Pro			
	245	250	255
Lys Val Leu Lys Gly His Asp Asp His Val Ile Thr Cys Leu Gln			
	260	265	270
Phe Cys Gly Asn Arg Ile Val Ser Gly Ser Asp Asp Asn Thr Leu			
	275	280	285
Lys Val Trp Ser Ala Val Thr Gly Lys Cys Leu Arg Thr Leu Val			
	290	295	300
Gly His Thr Gly Gly Val Trp Ser Ser Gln Met Arg Asp Asn Ile			
	305	310	315
Ile Ile Ser Gly Ser Thr Asp Arg Thr Leu Lys Val Trp Asn Ala			
	320	325	330
Glu Thr Gly Glu Cys Ile His Thr Leu Tyr Gly His Thr Ser Thr			
	335	340	345
Val Arg Cys Met His Leu His Glu Lys Arg Val Val Ser Gly Ser			
	350	355	360
Arg Asp Ala Thr Leu Arg Val Trp Asp Ile Glu Thr Gly Gln Cys			
	365	370	375
Leu His Val Leu Met Gly His Val Ala Ala Val Arg Cys Val Gln			
	380	385	390
Tyr Asp Gly Arg Arg Val Val Ser Gly Ala Tyr Asp Phe Met Val			
	395	400	405
Lys Val Trp Asp Pro Glu Thr Glu Thr Cys Leu His Thr Leu Gln			
	410	415	420
Gly His Thr Asn Arg Val Tyr Ser Leu Gln Phe Asp Gly Ile His			
	425	430	435
Val Val Ser Gly Ser Leu Asp Thr Ser Ile Arg Val Trp Asp Val			
	440	445	450
Glu Thr Gly Asn Cys Ile His Thr Leu Thr Gly His Gln Ser Leu			
	455	460	465
Thr Ser Gly Met Glu Leu Lys Asp Asn Ile Leu Val Ser Gly Asn			
	470	475	480
Ala Asp Ser Thr Val Lys Ile Trp Asp Ile Lys Thr Gly Gln Cys			
	485	490	495
Leu Gln Thr Leu Gln Gly Pro Asn Lys His Gln Ser Ala Val Thr			
	500	505	510

Cys	Leu	Gln	Phe	Asn	Lys	Asn	Phe	Val	Ile	Thr	Ser	Ser	Asp	Asp
				515					520					525
Gly	Thr	Val	Lys	Leu	Trp	Asp	Leu	Lys	Thr	Gly	Glu	Phe	Ile	Arg
				530					535					540
Asn	Leu	Val	Thr	Leu	Glu	Ser	Gly	Gly	Ser	Gly	Gly	Val	Val	Trp
				545					550					555
Arg	Ile	Arg	Ala	Ser	Asn	Thr	Lys	Leu	Val	Cys	Ala	Val	Gly	Ser
				560					565					570
Arg	Asn	Gly	Thr	Glu	Glu	Thr	Lys	Leu	Leu	Val	Leu	Asp	Phe	Asp
				575					580					585

Val Asp Met Lys

<210> 30

<211> 3375

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 708398CB1

<400> 30

```

ggggagaagg gagcccgag gatcaggggt cagagttagg gggettcct cctgctgcac 60
cctcatctca gggccgcaa cttccagctg cagcggcgac tttcagtttc atttccacgg 120
accctcctgc ctgggcccgc agccgccgcc gcgatgccca gtaagttcag ctgccggcag 180
ctccgggagg cgggccagtg tttcgagagt ttcctggctg ttcggggact ggacatggag 240
acagatcgcg agcggctgcg gaccatttat aaccgcgact tcaagatcag ctttgggacc 300
cccgcccttg gcttctcctc catgctgtat ggaatgaaga ttgcaaactt ggcctacgtc 360
accaagactc gggtcaggtt cttcagactc gaccgctggg ccgacgtgcg gttcccagaa 420
aagaggagaa tgaagctggg gtcagatata agcaaacacc acaagtcact gctagccaag 480
atcttttatg acagggtga gtatcttcat gggaaacatg gtgtggatgt ggaagtccag 540
gggccccatg aagcccgaga tgggcagctc cttatccgcc tggatttgaa ccgcaaagag 600
gtgctgacct tgaggcttcg gaatggcgga acccagctctg ttaccctcac tcacctcttc 660
ccactctgcc ggacacccca gtttgctttc tacaatgaag accaggagt ggcctgtcca 720
ctgggccccg gtgaatgcta tgaactccat gtccattgta agaccagctt tgtgggctac 780
ttcccagcca cagtgtcttg ggagctgctg ggacctgggg agtcgggttc agaaggagcc 840
ggcacattct acattgcccg cttcttggtt gccgtcgccc acagccccct ggctgcacag 900
ctgaagccca tgactccctt caagcggacc cggatcaccg gaaaccctgt ggtgaccaat 960
cggatagagg aaggagagag acctgaccgc gctaagggtt atgacctgga gttaagtatg 1020
gcgctgggga catactacce acctccccgc ctcaggcagc tgctccccat gcttcttcag 1080
ggaacaagta tcttcactgc ccctaaggag atcgagaga tcaaggccca gctggagaca 1140
gccctgaagt ggaggaacta tgaggtgaag ctgcggctgc tgctgcacct ggaggaactg 1200
cagatggagc atgatatccg gcactatgac ctggagtcgg tgcccatgac ctgggacctt 1260
gtggaccaga accccaggct gctcacgctg gaggttcctg gagtgactga gagccgcccc 1320
tcagtgtctc ggggcgacca cctgtttgcc cttttgtcct cggagacaca ccaggaggac 1380
cccatcacat ataagggtt tgtgcacaag gtggaattgg accgtgtcaa gctgagcttt 1440
tccatgagcc tcctgagccg ctttgtggat gggctgacct tcaagggtgaa ctttaccttc 1500
aaccgccagc cgctgcgagt ccagcaccgt gccctggagc tgacagggcg ctggctgctg 1560
tggcccatgc tctttcctgt ggcacctcgg gacgtcccgc tgctgcccct agatgtgaaa 1620
ctcaagctgt acgaccggag tctggagtca aaccagagc agctgcaggc catgaggcac 1680
attgttacgg gcaccaccg tccagcccc tacatcatct ttgggcctcc aggcaccggc 1740
aagactgtca cgttagtgga ggcaattaag caggtggtga agcacttgcc caaagcccac 1800
atcttggtc gcgctccatc caactcaggg gctgacctac tctgtcaaag gctccgggtc 1860
caccttccta gctccatcta ccgctcctg gccccagca gggacatccg catggtacct 1920

```

```

gaggacatca agccctgctg caactgggac gcaaagaagg gggagtatgt atttcccgcc 1980
aagaagaagc tgcaggaata ccgggtctta attaccaccc tcatcactgc cggcagggtg 2040
gtctcggccc agtttcccat tgatcacttc acacacatct tcatcgatga ggctggccac 2100
tgcatggagc ctgagagtct ggtagctata gcagggtga tggaaagtaa ggaaacaggt 2160
gatccaggag ggcagctggt gctggcagga gacctcggc agctggggcc tgtgctgct 2220
tccccactga ccagaagca tggactggga tactcactgc tggagcggct gctcatctac 2280
aactccctgt acaagaaggg ccctgatggc tatgaccccc agttcataac caagctgctc 2340
cgcaactaca ggtctcatcc caccatcctg gacattccta accagctcta ttatgaaggg 2400
gagctgcagg cctgtgctga tgtcgtggat cgagaacgct tctgccgctg ggcgggccta 2460
cctcgacagg gctttcccat catctttcac ggcgtaatgg gcaaagatga gcgtgaaggc 2520
aacagcccat ccttcttcaa ccctgaagag gctgccacag tgacttccta cctgaagctg 2580
ctcctggccc cctcctccaa gaagggcaaa gctcgctga gccctcgaag tgtgggcgtc 2640
atctccccgt accggaaaca ggtggagaaa atccgttact gcatcaccaa acttgacagg 2700
gagcttcgag gactggatga catcaaggac ttgaagggtg gttcagtaga agaattccaa 2760
ggccaagaac gaagcgtcat cctcatctcc accgtgcgaa gcagccagag ctttgtgcag 2820
ctggatctgg actttaatct gggtttcctt aagaacccca agaggttcaa tgtagctgtg 2880
acccgggcca aggcctgct catcatcgtg ggaaccccc ttctcctggg ccatgaccct 2940
gactggaaag tattcctgga gttctgtaaa gaaaacggag ggtataccgg gtgtcccttc 3000
cctgccaac tggacctgca acagggacag aatttactgc aagggtctgag caagctcagc 3060
ccctctacct cagggcccca cagccatgac tacctcccc aggagcggga ggggaagg 3120
ggcctgtctc tgcaagtga gccagagtgg aggaatgagc tctgaagaca cagcacccag 3180
ccttctcgca ccagccaagc cttaactgcc tgctgaccc tgaaccagaa ccagctgaa 3240
ctgccccctc aagggacagg aaggctgggg gagggagtgt acaacccaag ccattccacc 3300
ccctccccctg ctggggagaa tgacacatca agctgctaac aattggggga aggggaagga 3360
agaaaactct gaaac

```

<210> 31

<211> 2434

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1259937CB1

<400> 31

```

caaggatccg atgggtatat ggagtgtgag gtaatggatc attcatgtgg aaggatgcag 60
ggggtttttg agaccagggt ttggaagaga gttcagcact gctggtagtt ttgggaatca 120
cccatgtgca ggcgacacat gaggcagtaa ggaactctgc aggggtccct gagatttgga 180
aatgtaggga agagcaatgg attgaggtcc gaacctggag gatctgctat acgcagagct 240
gggaggaggg acagagtcag taccagagtc ggaaaaaagc aggggtggga ggggaacctg 300
agtcaggaga cttgcctggc aggcgctgcc ctgccagcag aggcctgaca gtggtttcca 360
tgaactgcat ccctgctgtg ggctgggaca gggccactga cacagtatcg gagcacagaa 420
ggggaaagga gcaggaggga ttccaactct gccagttagc agctgtgtgg ctttgggcat 480
gttacttaac ctctctgagc ctcatattat tcatccataa aatggaaata aaaataatac 540
ttttgtcaaa ggcgattgt gaatathtag atcctcagaa taatgcctgg cttgtagcaa 600
atggtagctg gaggaaaagg aagagaaaac caaagtcagc agctgaagga ttttcatatt 660
agaactgctc tggacctatc tggcagatgc agaagcacac acacacggag gggcatggat 720
ttgccccgcc cttagacatg ttgtgtcttc tcctggatcc ttgggtcccag gtgccttacc 780
tgagctcagg tgaatgtggc aagcagagcc ctctgggtgg gtgaatgctg tgtggcggcc 840
gtgctcctgg tgacacaggg acctcacaat ccctccctcc acggtctcct ctcatgtcct 900
cccagcctta ttttctcggt cctcttcctc ccaggcccgg aacttgccctg tttggctccc 960
caaccaggac gagccccctc ctggcagcag ctgtgccatc caagtggggg ataaagtccc 1020
ctatgacatc tgccggccag accactcagt gttgacctg cagctgcctg tgacagcctc 1080

```



```
cgtgagagag gtgatggcag cgttggccca ggaggatggc tggaccaagg ggcaggtgct 1140
ggtgaaggtc aattctgcag gtgatgccat tggcctgcag ccagatgccg gtggtgtggc 1200
cacatctctg gggctcaatg agcgtctctt tgttgtcaac ccacaggaag tgcattgagct 1260
gateccacac cctgaccagc tggggccccc tgtgggctct gctgaggggc tggacctggt 1320
gagtggcaag gacctggcag gccagctgac ggaccacgac tggagcctct tcaacagtat 1380
ccaccaggtg gagctgatcc actatgtgct gggcccccag catctgcggg atgtcaccac 1440
cgccaacctg gagcgcttca tgcgcgctt caatgagctg cagtactggg tggccaccga 1500
gctgtgtctc tgccccgtgc ccggcccccg ggcccagctg ctcaggaagt tcattaagct 1560
ggcgggccac ctcaaggagc agaagaatct caattccttc tttgccgtca tgtttggcct 1620
cagcaactcg gccatcagcc gcctagccca cacctgggag cggctgcctc acaaagtccg 1680
gaagctgtac tccgccctcg agaggctgct ggatccctca tgggaaccac gggatataccg 1740
actggccctc gccaaagtct cccctcctgt catcccttc atgccccttc ttctcaaaga 1800
catgaccttc attcatgagg gaaaccacac actagtggag aatctcatca actttgagaa 1860
gatgagaatg atggccagag ccgcgcggat gctgcaccac tgccgaagcc acaaccctgt 1920
gcctctctca ccactcagaa gccgagtttc ccacctccac gaggacagcc aggtggcgag 1980
gatttccaca tgctcggagc agtccctgag caccggagt ccagccagca cctgggctta 2040
tgtccagcag ctgaaggatca ttgacaacca gcgggaactc tcccgcctct cccgagagct 2100
ggagccatga ggaggggctg ggactggagc tggagcaggc acttgagacc gggaaagcca 2160
gggtgtgccc ggccaagata ctcacaggct ggccacagct gggcaaggct ctccgtggag 2220
tggactcgag tccctggagc aggcagtgtg gaggcagcca tccctgtga tgactggcag 2280
ctaaggagga cctcggagtg gaccgagcc aggaataacg aatgacccaa ggccaaggaa 2340
gggaggacag agaggcccca ggagtgggtg gagagtggag tgcgctggga cgttgtgtgc 2400
aatagagagg tctccacacc agaaaaaaaa aaaa 2434
```

<210> 32

<211> 892

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1452285CB1

<400> 32

```
cacgcctctc tctgcaccta cacttgcgct ccccaagtct ctctcgtgcg cagagcccag 60
gctgcgcttc cctggtcagg cacggcacgt ctggccggcc gccaggatgc aggccccgca 120
caaggagcac ctgtacaagt tgctggatgat tggcgacctg ggcgtgggga agaccagtat 180
catcaagcgc tacgtgcacc agaacttctc ctgcgactac cgggccacaa tcggcgtgga 240
cttcgcgctc aaggtgctcc actgggaccc ggagactgtg gtgcgcctgc agctctggga 300
tatcgcaggt caagaaagat ttggaaacat gacgagggtc tattaccgag aagctatggg 360
tgcatattat gtcttcgatg tcaccaggcc agccacattt gaagcagtgg caaagtggaa 420
aatgatttg gactccaagt taagtctccc taatggcaaa ccggtttcag tggttttgtt 480
ggccaacaaa tgtgaccagg ggaaggatgt gctcatgaac aatggcctca agatggacca 540
gttctgcaag gagcacggtt tcgtaggatg gtttgaaaca tcagcaaagg aaaatataaa 600
cattgatgaa gcctccagat gcctggatga acacatactt gcaaatgagt gtgacctaat 660
ggagtctatt gagccggacg tcgtgaagcc ccatctcaca tcaaccaagg ttgccagctg 720
ctctggctgt gccaaatcct agtaggcacc tttgctgggt tctggtagga atgacctcat 780
tgttccacaa attgtgcctc tatttttacc attttgggt aacgtcagga tagagatacc 840
acatgtggca agccaaagat ctatgcctcc atatgtgcct ttctgttagc tt 892
```

<210> 33

<211> 2288

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1812894CB1

<400> 33

```
cggtctcgagc ccagatggag gcaacagtac taggttgcag gacttgctaa gccggtaggt 60
ggcctgggggt aagagatggg aagagaggct gatccctgtc cccacagca ctggaggact 120
cccgggtccca ggaaggggca aatggtgagg ccgagtcagg tgagctcagc cggcttcggg 180
ctgagctggc aggcgccctg gcagaaatgg aaaccatgaa ggctgtggca gaggtgagcg 240
agagcacgaa ggccgaggct gtggctgcgg tgcagcggca gtgccaagag gaggtggcct 300
cgctgcaggc catcctgaaa gactccatca gcagctatga agcccagatc accgccctga 360
agcaggagcg acagcagcag cagcaggact gtgaggagaa ggagcgggag ctgggcccgc 420
tgaagcagct gctgtcccgg gcctaccccc tggactccct ggagaagcag atggaaaagg 480
cccacgagga ctccggagaag ctgcgggaga tcgtactgcc catggaaaag gagatcgagg 540
agctgaaggc gaagctgctg agggccgagg agctgattca ggagatccag agacgtcccc 600
ggcatgcccc ttccctgcac ggctccacgg agttgctgcc cctgtcccgg gatccatcgc 660
ccccgctgga gcctctggag gagctgagcg gagatggggg tccagccgct gaggccttcg 720
ctcacaactg cgatgacagc gcctccatct cctccttctc ccttggcggt ggggtcggca 780
gcagctcctc cctgccccaa agccgccagg gcctgagccc tgaacaggaa gagacggcct 840
cgctggtgtc tacgggcacc ctggttcccg agggcatcta cctgccccct cctggctacc 900
agctcgtccc agacactcag tgggagcagc tgcagacaga gggccgacag ctgcagaagg 960
acctggagag cgtcagtcgc gagcgggacg agctccaaga gggcctgaga cggagcaatg 1020
aggactgtgc caagcagatg caggtgtctc tggcccaggc ccagaactca gagcagctgc 1080
tgcggaccct gcaagggacc gtgagccagg cccaggagcg ggtgcagctg cagatggcgg 1140
agctggtcac caccacaag tgctgcacc atgaggtaaa gcggttgaat gaggaaaacc 1200
aagggctccg ggccgagcag ctgccatcct cagcccccca gggctcgcag caggagcagg 1260
gcgaggagga atcactgccc agctctgtgc cagagctgca gcagctgctg tgctgcacgc 1320
ggcaagaggc gagggcccgg ctgcaggccc aggagcacgg ggccgagcgc ctgcggatcg 1380
agatcgtgac gctgcgggag gctctggagg aggagacagt ggccagggcc agcctggagg 1440
ggcagctgag ggtgcagcgg gaggagacag aggtgctgga ggctccctg tgcagcctga 1500
ggacagagat ggagcgggtg cagcaggaac agagcaaggc ccagctccca gacctcctct 1560
cagaacagag ggccaagggt ctgcggctgc aggcagagct ggagaccagt gagcaggtgc 1620
agagggattt cgtgcgactg tcccaggccc tgcagggtgc cctagagcgg atccgccagg 1680
ctgagaccct ggagcaagtg cgcagcatca tggatgaggc gccactcacg gacgtcaggg 1740
acatcaagga cacctgaggg gtcaggatat cccaccccc accctgggaa agacgccttt 1800
ccccactcct gaaccatgag gcctcgtctt ggggtcttgg atggcttttc caccgtccct 1860
gagactgggg ttgaggggac tgacgggggc caccaccgcc ccgcctcca gcgcctcctc 1920
ccagggtggc tgggcctcct gttctcaggg atcacacctg ggtgaggggc ccaagccctc 1980
cccgaacca aaggtgcagg ctccaggcctg cggctttctg gctgctgtgc tgcctcctgg 2040
gctccagccc tcccctgccc ccagcccgtc ccctgcccag ggcacagcgg agccatgggg 2100
gctgggagtc cccatcagag gcagtgggtt gggccccggc cctgggacag gcagctgcct 2160
tctggtctgc atgacactaa gacgcttgtc cacagcggcg acccaggcct ccaagcttgc 2220
acagaggcaa ggccagactt ttccgtcggt tattttcaat aaataagcag ctccagcga 2280
aaaaaaaa
```

<210> 34

<211> 1813

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 1708, 1711, 1713, 1715

<223> Incyte ID No: a o r g o r c o r t, unknown, or other

<220>

<221> misc_feature

<223> Incyte ID No: 3074884CB1

<400> 34

```

gcacgaagga ggcccggctt caggtggccc tggcggagat gccggaagat gcggacgaga 60
acgccgagga ggagctgctg cggggagagc ctctgctgcc ggcggggacc cagcgcgtgt 120
gtctggttca ccctgacgtc aagtggggcc cggggaagtc gcagatgact cgagccgagt 180
ggcaggtggc ggaggccaca gcgctggtgc acacgctgga cggctggtcc gtggtgcaga 240
caatggtcgt gtccaccaa acgccggaca ggaagctcat ctttggcaaa gggaactttg 300
agcacctgac agaaaagatc cgagggtctc cagacgtcac gtgcgtcttc ctgaacgtgg 360
agaggatggc tgccccgacc aagaaagaac tggaagccgc ctggggcgctg gaggtgtttg 420
accgcttcac ggtcgtcctg cacatcttcc gctgtaacgc ccgcacgaag gagggccggc 480
ttcaggtggc cctggcggag atgccgctgc acaggtcgaa cttgaaaagg gacgtcgccc 540
acctgtaccg aggagtcggc tcgcgtaca tcatggggtc aggagaatcc ttcattgcagc 600
tgcagcagcg tctcctgaga gagaaggagg ccaagatcag gaaggccttg gacaggcttc 660
gcaagaagag gcacctgctc cgccggcagc ggacgaggcg ggagttcccc gtgatctccg 720
tggtggggta caccaactgc ggaaagacca cgctgatcaa ggcactgacg ggcgatgccg 780
ccatccagcc acgggaccag ctgtttgcc cgtggacgt cacggcccac gcgggcacgc 840
tgccctcacg catgaccgtc ctgtacgtgg acaccatcgg ctccctctcc cagctgccgc 900
acggcctcat cgagtccttc tccgccaccc tggaagacgt ggcccactcg gatctcatct 960
tgcacgtgag ggacgtcagc caccgcgagg cggagctcca gaaatgcagc gttctgtcca 1020
cgctgcgtgg cctgcagctg cccgccccgc tcctggactc catggtggag gttcacaaca 1080
aggtggacct cgtgcccggg tacagcccca cggaaaccga cgtcgtgccc gtgtctgccc 1140
tgcggggcca cgggctccag gagctgaaag ctgagctcga tgcggcggtt ttgaaggcga 1200
cggggagaca gatcctcact ctccgtgtga ggctcgcagg ggcgcagctc agctggctgt 1260
ataaggaggc cacagttcag gaggtggacg tgatccctga ggacggggcg gccgacgtga 1320
gggtcatcat cagcaactca gcctacggca aattccggaa gctctttcca ggatgaacgg 1380
acgccacag aggcctgcgg ggtgggggca tcgctgcctg gggagctgag gcgttaccgc 1440
tgtgttgggg gcagcttggt gtcaggtgca gcagggtcct ccttgtctgg ttctgcaccc 1500
gtctcgctcc cagccatttg ctgggatgac cgtgcaggcc ggtgacacgg ccgcacctgc 1560
cccaaagcgg gccgcccag cgtccactcc aagcctgagc atccacacaa tccagtgagg 1620
ccctcggtgc ctgctgtgaa ctgctttccc tcggaatgtt tccgtaacag gacattaaac 1680
ctttgttttt acttccgtga aaaaaaanac ngngnaaaaa aaaaaggggc ggcccgctcc 1740
tagaggattc caagccttac cgtaacgcgt tgcattggcg agcggtcata agcttcttct 1800
aatagggggt cac 1813

```

<210> 35

<211> 1733

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3452277CB1

<400> 35

```

ctacggcctg gaccgagtga ccaatccgaa tgaagtcaag gtaaaccaga aacaaacagt 60
cgttgctgtc aaaaaagaga tcatgtatta ccaacaggcc ttgatgaggt ccacagtga 120

```

gtcttctgtg tccctgggag ggattgtgaa atacagtgag cagttctcat ccaacgatgc 180
catcatgtca ggctgcctcc ccagcaaccc ctggatcacc gatgacaccc agttctggga 240
cttaaattgcc aaattggtgg aaatcccaac caagatgcga gtggaacgat gggccttcaa 300
cttcagcgaa ttgatccgag accccaaagg tcgacagagc ttccagtact tcctcaagaa 360
agaattcagt ggagagaatc tgggattctg ggaagcctgc gaggatctga agtatggaga 420
tcagtccaaa gtcaaggaga aagcagagga gatttacaag ctgttccttg ccccgggggc 480
gaggcgctgg atcaacatag atggcaaaac catggacatc acagtgaagg ggctgaagca 540
ccccaccgc tatgtgctgg acgccgcaca aaccacatt tacatgctca tgaagaagga 600
ttcttatgct cgctatttaa aatctccgat ctataaggac atgctggcca aagctattga 660
acctcaggaa accaccaaga aaagctccac cctccctttt atgcggcgct acctgcgctc 720
cagcccaagc cctgtcatcc tgagacagct ggaagaggaa gccaaaggccc gagaagcagc 780
caacactgtg gacatcaccc agccgggcca gcacatggct cccagccccc atctgaccgt 840
gtacaccggg acctgcatgc cccgtctcc ttctagcccc ttctcctcct cctgccgctc 900
ccccaggaag cctttcgctt caccagccg cttcatccgg cgaccagca ccaccatctg 960
cccctaccc atcagagtgg ccttggagag ctcacggggc ttggagcaga aaggggagtg 1020
cagcgggtcc atggccccc gtgggcccctc tgtcaccgag agcagcgagg cctccctcga 1080
cacctcctgg cctcgcagcc ggcccagggc cctcctaag gcccgcatgg ctctgtcctt 1140
cagcaggttt ctgagacgag gctgtctggc ctcacctgtc ttgcccaggc tctcacccaa 1200
gtgccctgct gtgtcccacg ggagggtgca gcccctgggg gacgtggggc agcagctgcc 1260
acgattgaaa tccaagagag tagcaaactt tttccagatc aaaatggatg tgcccacggg 1320
gagcgggacc tgcttgatgg actcggagga tgctggaaca ggagagtcgg gtgaccgggc 1380
cacagaaaag gaggtcatct gcccctggga gagcctgtaa ggaaagaggc aggctgagct 1440
gggggctctg gaccaggaag atgctctgac agatgccatg gtatgggcca caggacacac 1500
ttgctcgaga accaaagtgc atttgggtga catttgaaga ttggggagac aagatggggg 1560
agattgtggc aaagaatgct ctggctgggt accaggggccc aactccttct cctcttcctg 1620
accctccctc ccctgggcag aagaaacgca tgtggaccag aagactttcc ctgctgcctt 1680
aaaaccaata aaaggttaac ttaagtttc ttggaaaaaa aaaaaaaaag ggg 1733

<210> 36

<211> 1776

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4203832CE1

<400> 36

cccagccgag cccagcctag cccgagccca gcccagcgca accgcgagcc ccaagcccga 60
gccgcgcccc gcccagcgag agccctccag ccgctcacc cgcgtgccac cccagcgacc 120
ctcagccgct ctctgccctt ctctcgccc cgcgcccgc ctccggccc ctctgcccga 180
tgaaactggc cgcgatgatc aagaagatgt gcccagcgca ctccggagctg agtatcccgg 240
ccaagaactg ctatcgcatg gtcacctcgt gctcgtccaa ggtgggcaag acggccatcg 300
tgtcgcgctt cctcacgggc cgcttcgagg acgcctacac gcctaccatc gaggacttcc 360
accgcaagtt ctactccatc cgcggcgagg tctaccagct cgacatcctc gacacgtccg 420
gcaaccaccc gtccccgcc atgcggtgcc tctccatcct cacaggagac gttttcatcc 480
tggtgttcag tctggacaac cgcgactcct tcgaggaggt gcagcggctc aggcagcaga 540
tcctcgacac caagtcttgc ctcaagaaca aaaccaagga gaacgtggac gtgcccctgg 600
tcactctgcg caacaagggt gaccgcgact tctaccgcga ggtggaccag cgcgagatcg 660
agcagctggt gggcgacgac ccccagcgct gcgcctactt cgagatctcg gccaaagaaga 720
acagcagcct ggaccagatg ttccgcgcgc tcttcgccat ggccaagctg cccagcgaga 780
tgagcccaga cctgcaccgc aaggtctcgg tgcagtactg cgacgtgctg cacaagaagg 840
cgctgcggaa caagaagctg ctgcggggcc gcagcggcgg cggcggcggc gacccgggcg 900
acgccttttg catcgtggca cccttcgcgc gccggcccag cgtacacagc gacctcatgt 960


```

acatccgcga gaaggccagc gccggcagcc aggccaagga caaggagcgc tgcgtcatca 1020
gctaggagcc ccgccgcgct ggcgacacaa cctaaggagg acctttttgt taagtcaaata 1080
ccaacggccc ggtgcgcccc aggccgggag cgcgcgcgga ctggcgtctc cctccccggc 1140
gatccgcccc cagcactggg gaggcgccac tgaaccgaga agggacggtc atctgctccg 1200
gaaggaaaga gaacgggcca agactgggac tattccccac ccccggtccc ccattgaggc 1260
ccgccacccc cataactttg ggagcgaggg ccagcccgag ggtggattta tcttctcaaa 1320
gacctaaagag tgagcgcggg gtgggggagg gatgtgaagt tatccagcct ctgctaggct 1380
tcaagaaacc gtcattgccc cttgagggtc aggaccacag gggcattatc ttgtctgtga 1440
ttccgggttg ctgtgacagc cggtagagcc tctgccctcc cgaaactaag cgggggggag 1500
tgggtcaaat catagccaag tgacttgttt acatgtgagt gaaactgcac aaaggaacac 1560
aaaacaaaac ttgcacttta acggtagttc cgggtgtcaac atggacacga acaaacctt 1620
acccaggtgt ttatactgtg tgtgtgtgag gtctttaaag ttattgcttt atttggtttt 1680
ttaatatata ataaaaataa ttaaaatgga aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1740
aaaaaaaaaa aaaaaaaaaa gaaaaaaaaa aaaaaa 1776

```

<210> 37

<211> 1316

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 104368CB1

<400> 37

```

cggacgcgtg gggcggttt tctctttagc gctcagggcg ctgaccact gctcttcctc 60
ttaagaaagt gctccattcc ttccggcgcc cggagctgct ggcccaaagg gatccggagc 120
gagctagggc agacaccatg accacccttg atgataagtt gctgggggag aaactgcagt 180
actactatag cagcagttag gatgaggaca gtgaccacga ggacaaggac cgaggcagat 240
gtgccccagc cagcagttct gtgcctgcag aggctgagct ggcaggcgaa ggcattctcag 300
ttaacacagg cccaaaagggt gtgatcaatg actggcgccg cttcaagcag ttggagacag 360
agcagagggg ggagcagtg cgggagatgg aaaggctgat caagaagctg tcaatgactt 420
gcaggtccca tctggatgaa gaggaggagc aacagaaaca gaaagacctc caggagaaga 480
tcagtgggaa gatgactctg aaggagtgtt ccataatgaa tgaggaccaaa gatgatgaag 540
agtttctgca gcagtaccgg aagcagcgaa tggaagagat gcggcagcag cttcacaagg 600
ggccccaatt caagcaggtt tttgagatct ccagtggaga agggttttta gacatgattg 660
ataaagaaca gaaaagcatt gtcattcatg ttcataattt tgaggatggc attccaggga 720
ccgaagccat gaatggttgc atgatctgcc ttgccgcaga gtaccagct gtcaagttct 780
gcaaggtgaa gagctcagtt attggcgcca gcagtcagtt caccaggaat gcccttcctg 840
ccctgctgat ctataagggg ggtgaattga tcggcaattt tgttcgtgtt actgaccagc 900
tgggggatga tttctttgct gtggaccttg aagcttttct ccaggaattt ggattactcc 960
cagaaaagga agtcttggtg ctgacatctg tgcgtaactc tgccacgtgt cacagtgagg 1020
atagcgacct ggaaatagat tgaactgata gtctagttgc atagatttct cattgttttg 1080
gttggaatac acgtcattgt ttatttttgt tcctttgtct tctggctttt cagctgttct 1140
ttgtagtccc ttttattatg cataaaataa agaaattctt agattaaatc agaattgctg 1200
ataaccttgt agctagcaat aaggtgactt acagttgtat aaacaggaag ccaggctttt 1260
gaactgttta ctttaagattc tgtggtgtga catctctgtt attgtttcca gtcaat 1316

```

<210> 38

<211> 1554

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1441680CB1

<400> 38

```
gtggcttggtg gagtggcgac cgtagtgag gcggttgctg agacagacgc tgaggcgggt 60
aggaggagcc cgagccgtaa ggaagccgt gatgagggcc gtgttgacgt ggagagataa 120
agccgagcac tgtataaatg acatcgcat taagcctgat ggaactcaac tgattttggc 180
tgccggaagc agattactgg tttatgacac ctctgatggc accttacttc agcccctcaa 240
gggacacaaa gacactgtgt actgtgtggc atatgcgaag gatggcaagc gctttgcttc 300
tggatcagct gacaaaagcg ttattatctg gacatcaaaa ctggaaggca ttctgaagta 360
cacgcacaat gatgctatac aatgtgtctc ctacaatcct attactcatc aactggcatc 420
ttgttcctcc agtgactttg gggtgtggc tcctgaacag aagtctgtct ccaaacacaa 480
atcaagcagc aagatcatct gctgcagctg gacaaatgat ggtcagtacc tggcgctggg 540
gatgttcaat gggatcatca gcatacggaa caaaaatggc gaggagaaag taaagatcga 600
gcggccgggg ggctccctct cgccaatatg gtccatctgc tggaaacctt caagagagga 660
acgtaatgac atcctggctg tggctgactg gggacagaaa gtttccttct accagctgag 720
tggaaaacag attggaaggg atcgggcact gaactttgac ccctgctgca tcagctactt 780
tactaaaggc gagtacattt tgctgggggg ttcagacaag caagtatctc ttttcaccaa 840
ggatggagtg cggcttgagg ctgttgggga gcagaactcc tgggtgtgga cgtgtcaagc 900
gaaaccggat tccaactatg tgggtggtcg ctgccaggac ggcaccattt ccttctacca 960
gcttattttc agcacagtcc atggagttaa caaggaccgc tatgcctaca gggatagcat 1020
gactgacgtc attgtgcagc acctgatcac tgagcagaaa gttcggatta aatgcaaaga 1080
gcttgtcaag aagattgcc a tctacagaaa tcgattggct atccaactgc cagagaaaat 1140
cctcatctat gagttgtatt cagaggactt atcagacatg cattaccggg taaaggagaa 1200
gattatcaag aagtttgagt gcaacctcct ggtggtgtgt gccaatcaca tcatcctgtg 1260
ccaggagaaa cggctgcagt gcctgtcctt cagcggagtg aaggagcggg agtggcagat 1320
ggagtctctc attcgttaca tcaaggtgat cgggtggcct cctggaagag aaggcctctt 1380
agtggggctg aagaagatgt acttgtagt gtattcatc atattgattg taaaggatta 1440
tttttctc agtactgatg tccttgga tcttacctgg aaacatgttt gcaaaaaaca 1500
ttattgggtc tttcatcttt tttcttggtt ttacatattt gttcaataaa aata 1554
```

<210> 39

<211> 2320

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1494955CB1

<400> 39

```
gggcgcccgc gtactcacta gctgaggtgg cagtgggtcc accaacaatgg agctctcgca 60
gatgtcggag ctcatggggc tgctgggtgtt gcttggggctg ctggccctga tggcgacggc 120
ggcggtagcg cgggggtggc tgcgcgcggg ggaggagagg agcggccggc ccgcctgcca 180
aaaagcaaat ggatttccac ctgacaaatc ttcgggatcc aagaagcaga aacaatatca 240
gcggattcgg aaggagaagc ctcaacaaca caacttcacc caccgcctcc tggctgcagc 300
tctgaagagc cacagcggga acatatcttg catggacttt agcagcaatg gcaaatacct 360
ggctacctgt gcagatgatc gcaccatccg catctggagc accaaggact tcctgcagcg 420
agagcaccgc agcatgagag ccaacgtgga gctggaccac gccaccctgg tgcgcttcag 480
ccctgactgc agagccttca tcgtctggct ggccaacggg gacaccctcc gtgtcttcaa 540
gatgaccaag cgggaggatg ggggctacac cttcacagcc accccagagg acttccctaa 600
aaagcacaag gcgcctgtca tcgacattgg cattgctaac acaggaagt ttatcatgac 660
tgctccagt gacaccactg tcctcatctg gagcctgaag ggtcaagtgc tgtctaccat 720
```



```

caacaccaac cagatgaaca acacacacgc tgctgtatct ccctgtggca gatttgtagc 780
ctcgtgtggc ttcaccccag atgtgaaggt ttgggaagtc tgctttggaa agaaggggga 840
gttccaggag gtggtgcgag ccttcgaact aaagggccac tccgcggctg tgcactcggt 900
tgctttctcc aacgactcac ggaggatggc ttctgtctcc aaggatggta catggaaact 960
gtgggacaca gatgtggaat acaagaagaa gcaggacccc tacttgctga agacaggccg 1020
ctttgaagag gcggcggtg ccgcgcgtg ccgcctggcc ctctcccca acgcccaggt 1080
cttggccttg gccagtggca gtagtattca tctctacaat acccggcggg gcgagaagga 1140
ggagtgcctt gagcgggtcc atggcgagtg tatcgccaac ttgtcctttg acatcactgg 1200
ccgctttctg gcctcctgtg gggaccgggc ggtgcggctg tttcacaaca ctctggcca 1260
ccgagccatg gtggaggaga tgcaggcca cctgaagcgg gcctccaacg agagcacccg 1320
ccagaggctg cagcagcagc tgaccaggc ccaagagacc ctgaagagcc tgggtgccct 1380
gaagaagtga ctctgggagg gcccgcgca gaggattgag gaggaggat ctggcctcct 1440
catggcgctg ctgccatctt tcctcccagg tggaagcctt tcagaaggag tctcctgggt 1500
ttcttactgg tggcctgtc tcttccatt gaaactactc ttgtctactt aggtctctct 1560
cttcttgctg gctgtgactc ctccctgact agtggccaag gtgcttttct tcctcccagg 1620
cccagtgggt ggaatctgtc ccacactggc actgaggaga atggtagaga ggagaggaga 1680
gagagagaga atgtgatttt tggccttggt gcagcacatc ctcacacca aagaagtttg 1740
taaagtgtcc agaacaacct agagaacacc tgagtactaa gcagcagttt tgcaaggatg 1800
ggagactggg atagcttccc atcacagaac tgtgttccat caaaaagaca ctaagggtt 1860
tccttctggg cctcagttct atttgtaaga tggagaataa tcctctctgt gaactccttg 1920
caaagatgat atgaggctaa gagaatatca agtccccagg tctggaagaa aagtagaaaa 1980
gagtagtact attgtccaat gtcataaag tggtaaaagt gggaaccagt gtgctttgaa 2040
accaaattag aaacacattc cttgggaagg caaagttttc tgggacttga tcatacattt 2100
tatatgggtg ggacttctct cttcgggaga tgatatcttg tttaaggaga cctcttttca 2160
gttcatcaag ttcatacagat atttgagtgc ccactctgtg cccaaataaa tatgagctgg 2220
ggattaaata cgaataagac atggtttctg ccatcaaaga tggctggtgg gagagagaga 2280
tacaccctta ttaagtgtt tgtgttagtt tattcatagc 2320

```

<210> 40

<211> 879

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1508161CB1

<400> 40

```

gaagaatttg ttcaggcgtt cgtgcagaag gacccttttg ataatgacaa gagttgctac 60
agtgaacgga agaaaacacg aaacttagaa gcttacgtgg aatggtttaa tcgcctcagc 120
tacttggttg ctacagaaat ctgtatgcct gttaagaaaa aacaccgagc aagaatgatt 180
gagtatttca ttgacgtagc tcgggagtggt tttaacattg gcaacttcaa ctcttgatg 240
gcgataatct ctggtatgaa tatgagccca gtctctcgac taaaaaaaac ttgggcaaaa 300
gtgaagactg caaaatttga cattcttgag catcagatgg acccttcaag caatttctat 360
aattatcgaa cagctcttcg tggggcagca caaaggtctt taactgctca tagtagtaga 420
gaaaagattg tgataccatt cttcagtcct ttaatcaaag atatttattt cctcaatgag 480
ggttggtgca accgccttcc caatggccat gtcaattttg agaaattttg ggaactggcc 540
aaacaagtga gtgaatttat gacatggaaa caagtggagt gtccatttga gagggaccgg 600
aagatcttgc agtatctgct cacagtacca gtcttcagtg aagatgctct ctacttggct 660
tcttatgaga gtgaaggacc tgaaaatcat atagagaaag acagatggaa gtctttaagg 720
tcgagcctct taggcagagt ttaacacatg ggagctgcct gcctgctgct gctgctgctt 780
cctgcagatc atggaggggc tggcctttgt tttctggcat ctcgtaccac gaacactcat 840
gaagaccctg cagtcattgg agcaccggg tcagcaaag 879

```

<210> 41
<211> 2248
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 1811877CB1

<400> 41
ggaggaccag gaggacatca ctgcctttga cctcagccct gacaacgagg tgctggtgac 60
agccagtcgg gcattgctgc tggctcagtg ggcctggcaa gagggcagcg ttacccgcct 120
gtggaaggcg atacacacgg ccccgctggc caccatggcc ttcgacccca cctccactct 180
gctagccaca ggtggctgtg atggggccgt gcgcgtctgg gacatcgtgc ggcactacgg 240
gacacaccac ttccgaggct cgcgggtgt cgtgcaccta gtggccttcc acccggaacc 300
tacacgcctg ctgctcttct cctcggccac ggatgccgcc atccgcgtgt ggtcactgca 360
ggaccggtca tgcttggtg tgctgactgc ccactacagc gccgtcacct cactggcctt 420
cagcgccgac ggccacacca tgctcagctc cggccgtgac aagatatgta tcatctggga 480
ccttcagagc tgccaggcca cgaggaccgt gcctgtgttt gagagcgtgg aggctgctgt 540
gctgttgcca gaggagccag tgtcccagct ggggtgtgaag tccccagggc tgtactttct 600
gacagctggc gaccaaggca ctctgcgcgt gtgggaggca gcttctgggc agtgtgtgta 660
cacgcaggcc cagccgcccgg gccctgggca ggagctgacc cactgcaccc tggcacacac 720
cgccggcgtg gtcttcaccg ccaccgccga ccacaacctg ttgctctacg aggctcgtc 780
cctgcggctg cagaaacagt tcgctggcta cagtgaggag gttttggatg tccggtttct 840
tgggcccag gactcccacg ttgtcgtggc ctccaatagc ccctgcctaa aagtgtttga 900
gctgcagacg tcagcctgcc agatcctcca cggccacacg gatatcgtcc tggccctgga 960
tgtgttccgg aaggggtggc tctttgccag ctgtgccaag gatcagagcg tccgtatctg 1020
gagaatgaac aaggctggcc aggtgatgtg cgtggctcag ggttccggtc acacacacag 1080
tgtgggcacc gtctgctgct ctaggctgaa ggagtccttc ctggtgacag gcagccagga 1140
ctgcactgtg aagctgtggc ctcttcccaa agccttgctg tccaagaaca cagccccaga 1200
caacggccct atcctcctgc agggccagac cactcagcgc tgccatgata aggacatcaa 1260
cagcgtggct attgccccca acgacaagct gctggccaca ggctcacagg accgcacggc 1320
caagctctgg gccctgccac agtgccagct gctgggtgtc ttctcaggcc accggcgtgg 1380
cctctggtgc gtccagttct ctcccatgga ccagggtgctg gccacggcct cagctgatgg 1440
caccatcaag ctctgggcac tccaggactt cagctgtctc aagacatttg aggggcacga 1500
tgcttctgtg ctgaaggtgg cctttgtgag ccgtggcacg cagctgctgt ccagcggttc 1560
ggatggcctc gtgaagctct ggaccatcaa gaacaacgag tgtgtgcgga cgctggatgc 1620
ccacgaggac aaggtctggg ggctgcaact cagccggctg gacgaccacg ccctcactgg 1680
ggccagtgac tcccaggtca tcctctggaa ggatgtgacc gaggcggagc aggcagagga 1740
gcaggccagg caagaggagc aggtggtcag gcagcaagag ctggacaacc tgctgcatga 1800
gaagcggtag ctgcggggcg tgggcctggc catctccctg gatcggcccc acaccgtgct 1860
gactgtcatc caggccatcc ggagggaccc tgaggcctgc gagaagctgg aagccaccat 1920
gctccgactg cggcgcgacc agaaagaggc cctgctgcgc ttctgcgtca cgtggaacac 1980
caactcgcgg cactgccacg agggccaggc cgtgctgggt gtgctcttga ggcgagaggc 2040
ccccgaggag ctgctggcct acgaaggcgt gcgggcagcg cttgaggccc tgctgcccta 2100
cactgagcgg cactttcagc ggctcagcag gaccctccag gccgcgctt tcttggactt 2160
cctgtggcac aacatgaagc tccctgtgcc ggccgcccgc cccaccccct gggaaaccca 2220
taaaggcgca ctgccctaaa aaaaaaaaaa 2248

<210> 42
<211> 2146
<212> DNA
<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1848674CB1

<400> 42

```
gttattggca agttcccctg cagttgtttg ggctgtccct gtggctggtt ctggggtgtg 60
cggccagcca tggagcgctc tgggcccgag gaagtgcagc gctcagacgc gtcgggaccg 120
gacccgcagc ttgcggtcac catgggcttc acgggggttcg gtaaaaaagc tcgcacattt 180
gacttggaag caatgtttga acaaactcga aggacagctg tggaaagaag tcgcaaaaca 240
ctggaagcaa gagaaaaaga ggaagaaatg aacagagaga aagaattaag aagacaaaat 300
gaagatattg agccaacatc ctcaagatca aatgtggtca gagattgctc caaatcatct 360
tccagggata cgagcagcag tgaagtga cagagttctg actcttctga tgatgagtta 420
attggccctc ctttaccctc taaaatggta ggaaaaccag ttaattttat ggaggaagat 480
atcctcggtc ctttacctcc acctcttaat gaagaagaag aagaagcaga ggaagaagaa 540
gaggaagagg aggaagagga aaatcctgtt cacaagattc ctgactcgca tgagataacg 600
ctgaagcatg gcactaaaac agtgtctgct ttgggtcttg atccctcagg tgcccgtttg 660
gtgacaggag gatatgacta tgatgttaag ttttgggatt ttgctggaat ggatgcttct 720
tttaaggcat ttcgatccct tcagccctgt gagtgccatc agatcaagtc attacagtat 780
agtaacacag gagacatgat tcttgttgta tctggaagct ctgaggccaa ggtgattgac 840
agagatgggt ttgaagtaat ggaatgtata aaaggagacc agtatattgt ggacatggcc 900
aacaccaagg gtcatacagc aatgcttcat actggctcat ggcattccaa aataaaggga 960
gaatttatga cttgctcaaa tgatgcgact gtgaggacgt ggggaagtga aaatccaaag 1020
aagcaaaaaa gtgtgtttta accacggacg atgcaaggca aaaaagtcatt tcccactacg 1080
tgcacatata gtagagatgg aaacctcata gcagctgcct gccagaatgg aagcatacag 1140
atctgggacc gaaatttgac tgttcacctt aagttccact ataaacaggc tcatgactcg 1200
ggcacagaca cttcttgctg gactttttcc tatgatggta atgtccttgc ctctcgtgga 1260
ggtgacgatt cattaaaatt atgggacatc cgacaattta ataaaccact tttttcagcc 1320
tcgggtcttc ccaccatgtt cccaatgact gactgctgtt tcagtccaga tgataagctc 1380
atagtcactg gtacatctat tcaaagagga tgtggcagcg gcaaacttgt tttctttgag 1440
cgtaggactt tccaaagggt gtatgaaata gacatcacag atgcgagtggt tgttcgctgc 1500
ctgtggcatc caaagctgaa ccagatcatg gttggaactg gaaatggatt ggctaaagtc 1560
tattacgacc ccaacaagag tcagagggga gcaaaattat gtgtgggtta aaccagcgg 1620
aaggcaaac aagctgagac tctaactcag gactacatca tcacctca tgccttgctc 1680
atgttcctg agccccgcca acggagtaca aggaaacagc tggagaagga cagactggat 1740
ccctgaagt cgcataaacc tgaacctcct gtagcaggcc caggctcgtg tggccgagtt 1800
ggaaccacg ggggcactct ctcttcctat attgtgaaga acattgcttt ggacaagacc 1860
gatgacagta atcctcgga agccattttg cgtcatgcca aagcagcaga agacagccca 1920
tattgggttt ctccagcata ttccaagact cagcccaaaa ccatgtttgc ccaagttgaa 1980
tctgatgatg aggaagcaaa gaatgagcca gaatggaaaa aacgtaaaat ttgaagaatc 2040
tcatttgaga gctgtttgca tgagtgggag gggatatggga caggtttggg tttttttttt 2100
atgctcatga aattaaat tcatTTTTAT gaaaaaaaaa aaaaaa 2146
```

<210> 43

<211> 714

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2012970CB1

<400> 43

```
ggatggcgag cagcggaggc gaggcggtga cgagagcagc ggctccgcca ttggacgagg 60
```

```

aggcctgagg gacggggccag cgggtgcacaa gaagagaccg aggcgggtgg ccccgagaga 120
gccagggcca tggaggccaa catgccgaag cggaaggagc ctggcaggtc tctccgtatc 180
aaagtcattc ccatgggcaa cgccgaagtg gggaaaagct gtattataaa gcgatactgt 240
gagaaaagat tctgtgtctaa atacctggca acaattggaa ttgactatgg agtcacaaag 300
gtacacgtca gagacagaga aatcaaagtt aacatctttg atatggctgg acatcccttc 360
ttctatgagg ttcgaaatga gttttacaag gacacacagg gtgtgatact ggtctatgat 420
gttggggcaga aagactcctt tgacgccctt gatgcgtggc tggcagaaat gaagcaagag 480
cttggacctc atggaaacat ggaaaatatt atattttagt tttgtgccaa caagattgat 540
tgtaccaaac atcgtgtgtg agatgaaagt gaaggacgtc tttgggctga aagcaaaggg 600
ttcctgtact ttgaaacttc agcacaactt ggagaaggca ttaatgagat gttccagata 660
catcttggat agaactaatg gataaattag tctgttttaa aaagaaaaaa aaaa 714

```

<210> 44

<211> 1779

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2254315CB1

<400> 44

```

caggaggaag atggcggcgt ccgcagctgc cgctgagctc caggcttctg ggggtccgcg 60
gcacccagtg tgtctgttgg tgttgggaat ggcgggatcc gggaaaacca cttttgtaca 120
gaggctcaca ggacacctgc atgccaagg cactccaccg tatgtgatca acctggatcc 180
agcagtacat gaagtccctt ttccctgcaa tattgatatt cgtgatactg taaagtataa 240
agaagtaatg aaacaatatg gacttggacc caatggcggc atagtgcctt cactcaatct 300
ctttgctacc agatttgatc aggtgatgaa atttattgag aaggcccaga acatgtccaa 360
atatgtgttg attgacacac ctggacagat tgaggatatt acctgggtcag cttctgggac 420
aattatcact gaagcccttg catcctcatt tccaacagtt gtcattctatg taatggacac 480
atcgagaagt accaaccag tgaccttcatt gtccaacatg ctctatgcct gcagcatctt 540
atacaaaacc aagctgcctt tcattgtggt catgaataaa actgacatca ttgaccacag 600
ctttgcagtg gaatggatgc aggattttga ggctttccaa gatgccttga atcaagagac 660
tacatacgtc agtaacctga ctgcttcaat gagcctgggtg ttagatgagt ttacagctc 720
actcaggggtg gtgggtgtct ctgctgttct gggtactgga ttagatgaac tctttgtgca 780
agttaccagt gctgccgaag aatatgaaag ggagtatcgt cctgaatatg aacgtctgaa 840
aaaatcactg gccaacgcag agagccaaca gcagagagaa caactggaac gccttcgaaa 900
agatatgggt tctgtagcct tggatgcagg gactgccaaa gacagcttat ctctgtgct 960
gcacccttct gatttgatcc tgactcgagg aaccttggat gaagaggatg aggaagcaga 1020
cagcgatact gatgacattg accacagagt tacagaggaa agccatgaag agccagcatt 1080
ccagaatttt atgcaagaat cgatggcaca atactggaag agaaacaata aataggagac 1140
tttagcacac ttcaattggt tctagaagtc cagaattttg gacctccacg tgaaagaact 1200
gttcttacct ctgaactggg ggctcccata agggataatt ttcttcagag tagcaaagtt 1260
tctcttatta gagaaatctt gtgactcaga tgaagtcagg gatagaagac ccttggacct 1320
ggcagggttaa tgctgattat tccttggcct ttcccttgta tttatgcaag gaaggatata 1380
ctgagctgat actcttccaa gcctacaact tcaagtttta tcatttgaac tcaagtactt 1440
ttgctgctga ggaatggaat caaaagaacg tagtctcctg gtgaccacct cagatctcta 1500
ttattaggct agatgtatag cctctactcc ccagcttct tgctcttgac cctgcactgt 1560
aagttgccct tctattagca gccaaaggaa agggaaacat gagcttatcc agaacgggtg 1620
cagagtctcc ttggcaatca accaacgttg ctatgaaata tgcctcacac tgtatagctc 1680
attataggac gtcaggtttg ttgaaaaaag tgggcaagac atgattaatg aatcagaatc 1740
ctgtttcatt ggtgacttgg ataaagactt tttaattttt 1779

```


<210> 45
<211> 2234
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2415545CB1

<400> 45
cccccttctga aaaatggttt caccgcgtcc ttctctagaa cgggtcgggc cgctaccact 60
gtccggcccg gaggggaact gttttctccg gaagtgacaa cacgctgact aggaaaagga 120
ggaggcgggg cagtggggcc ttccggcgcg actatggaag gagccggcta cagggtggtg 180
tttgagaagg gcggagtgtt cctgcacacc agcgctaaga agtatcagga ccgagactct 240
ctcatcgctg gtgtcatccg tgcgtggaa aaggacaatg acgtcctcct gcactgggct 300
cctgtagagg aggctggaga ttccacccaa atcctcttct ccaagaagga ctccagtggg 360
ggtgactcat gtgcttctga ggaggaacca acctttgacc ccggctatga acctgactgg 420
gctgtcatca gcactgtgcg gccacagccc tgccactcag agcccacgag aggtgcagag 480
cccagctgcc ccaggggctc ctgggccttc tcagtgagtc tgggggagct aaagtccatc 540
cgccgctcca agccaggcct cagctgggccc tacctgggtc tggtgaccca ggctggaggt 600
tccttgcccg cactgcactt ccaccgcggg ggcacccgcg ccctgctccg cgtcctcagc 660
cgctacctgc tgttgccag ctccccgcag gactcccgcc tctacctgtt cttccccac 720
gactcctctg ctctctccaa ctcttccac cacctgcagc tctttgacca ggacagctcc 780
aatgtggtgt cacgcttctt ccaggatccc tactccacca ccttcagcag cttctccga 840
gtgaccaact tcttccgggg tgccctgcag ccacagcctg agggagccgc ctccgacctt 900
cccccgccac ccgacgatga gcccgagcct ggattcgagg tcatttctct tgtggagctg 960
gggcctcggc caaccgtgga gcggggccct ccagttacag aggaggagtg ggcacgccac 1020
gtgggcccctg aaggtcgctt gcagcaggtc cctgagctga agaaccggat cttctcgggg 1080
ggtctgagcc ccagcctgcg gcgcgaggcc tggaagtctc tcctagggta cctcagctgg 1140
gaaggcacag ctgaggagca caaggccac atacgcaaga aaacggatga gtatttccgc 1200
atgaagctgc agtggaatc tgtgagccct gagcaggagc ggagaaactc acttctgcat 1260
ggataccgca gcctcatcga aagggatgtg agccgcactg acaggaccaa caagttctac 1320
gagggctccc agaaccggg gctgggcccgt ctgaacgata tcctcctcac ctactgcatg 1380
tatcacttcg acctcggtta cgtccagggc atgagtgate ttctctcccc gatcctctac 1440
gtcattcaga acgaggtgga tgctttctgg tgtttctgtg gcttcattgga gctcgtgcaa 1500
gggaactttg aagagagcca ggagaccatg aagcggcaac tcgggcgact gctgctgctc 1560
ctgaggggtg tggacccctt gctctgcgac ttccctggatt cccaggactc cggtctctctc 1620
tgcttctgtt tccggtggct gctcatctgg ttcaagaggg aattcccctt cccggatgtc 1680
cttcggctgt gggaggtgct gtggacaggg ctccctggcc ccaatctgca cctgctgggtg 1740
gcctgcgcca tcctggacat ggagagggac acctcatgc tgtccggctt cggctccaat 1800
gagatcctca agcacatcaa cgagctgact atgaagctga gcgtggagga cgtgctgacc 1860
cgcgccgagg ccctgcaccg ccagctaacc gcctgcaccc gagctgcccc acaacgtgca 1920
ggagatcctg gggctggccc cgccacgcag agccccacag cccctcgccc accgctccc 1980
cgctgcctct gtacgcccac ccgggccccg cccaccccgc cgccctccac ggacacagcc 2040
ccgcagcccg acagcagcct ggagatcctg cccgaggagg aggacgaggg cgccgactcc 2100
taaccccgcc aggcagcctc gttctgcaca ggcactttag cccgagccag gcacacctgc 2160
gagggggcag gtgtgctccg ccgcctgctt gataagctgg cttcattaaa ctgacacttc 2220
tcaaaaaaaaaaaaa 2234

<210> 46
<211> 3150
<212> DNA
<213> Homo sapiens

<220>

<221> unsure

<222> 96, 97, 99, 3070-3072, 3074, 3078, 3080-3082, 3085-3087, 3091,

<222> 3099, 3100, 3103, 3107, 3110-3112, 3114, 3115, 3121, 3123, 3125,

<222> 3128, 3136, 3138, 3140, 3141, 3143, 3145, 3147, 3149

<223> Incyte ID No: a or g or c or t, unknown, or other

<220>

<221> misc_feature

<223> Incyte ID No: 2707969CB1

<400> 46

```

acacaggagc aatgcaaaat ttgataaagc atttttctgt cagatcagct gagccctact 60
gcccttcctc tcaagattcc tggagaccca gatgttnngna tctttcattg acaacaaaat 120
aatgtgtcat gatgatgatg ataaagaccc tgtactccgg gtatttgatt cccgagttga 180
caagatcagg ctgttgaatg ttcggacacc tactctccgt acatccatgt accagaagtg 240
taccactgtg gatgaagcag agaaagcaat tgagctgcgt ctggcaaaaa ttgaccatac 300
tgcaattcac ccacatttac ttgacatgaa gattggacaa gggaaatatg agccgggctt 360
cttccctaag ctgcagtctg atgtactttc cactgggcca gccagcaaca agtggacgaa 420
aaggaatgcc cctgcccagt ggaggcggaa agatcggcag aagcagcaca cagaacacct 480
gcgtttagat aatgaccaga gggagaagta catccaggaa gccaggacta tgggcagcac 540
tatccgccag cccaaactgt ccaacctctc tccatcagtg attgccaga ccaattggaa 600
gtttgtagag ggctgtctga aggaatgccg caataagacc aagaggatgc tgggtgaaaa 660
gatgggccga gaagctgtgg agctagggca tggggagggtg aacatcacag ggggtggaaga 720
gaacaccctg attgccagcc tttgtgatct cctggaaagg atctggagtc atggactaca 780
agtgaacag gggaaatcag ccttatggtc ccacctgtta cattatcagg acaaccggca 840
gagaaaactc acatcaggaa gcctcagtac ctcaggaata cttcttgatt cagaacgtag 900
gaagtctgat gccagctcac tcatgcctcc cctgaggatc tccctgattc aggatatgag 960
gcacatccag aacatcgggg aaatcaagac tgatgtggga aaggccagag catgggtgcg 1020
actgtccatg gaaaaaaagt tactttccag acacctgaag cagctcctct cagaccatga 1080
gctcaccaaa aagttatata agcgctatgc cttcctgcgc tgtgatgacg agaaggagca 1140
gttcctctat cacctcctgt ctttcaatgc cgtcgattac ttttgcttca ccaatgtctt 1200
cacaactatc ctgatcccgt accacattct gatcgtagca agcaagaagc tgggggggctc 1260
catgttcact gccaacccat ggatctgtat atcaggagaa ttgggtgaga cacagatcat 1320
gcagattccc aggaatgtgc tagagatgac cttcgagtgc cagaacttgg ggaagcttac 1380
tactgtccag attggccatg ataactctgg gctgtatgcc aaatggctgg tggagtatgt 1440
gatggtcagg aatgagatca caggacatac ctacaagttc ccgtgtggcc ggtgggttagg 1500
gaagggcatg gatgatggaa gcctggagcg gatcctagtt ggggagctgc tcacatccca 1560
gcctgagggtg gatgagaggc catgccggac cccgccgtg cagcagtcct ccagtgtcat 1620
ccggagggtt gttaccatct caccacaaca caagcccaag ctgaacactg ggcagatcca 1680
ggagtccatc ggggaggcag tcaatggcat tgtgaagcac ttccataagc ctgagaaaga 1740
gcgaggcagt ctgacgctgt tgctctgtgg agagtgtggc cttgtctcgg ccttggaaaca 1800
ggctttccag catggattta aatcgccccg gctcttcaaa aatgtcttca tttgggattt 1860
cctggaaaaa gcacaaacct attatgagac attagagaag aatgaagtag tccctgagga 1920
aaactggcat acaagagccc ggaacttctg ccgatttgct actgcaatca acaatactcc 1980
ccggaacatc ggcaaggatg gcaagtttca gatgctggtg tgcttgggag ccagagatca 2040
cctcctacac cactggattg ccctgctggc tgactgcccc atcactgcac acatgtatga 2100
ggatgtggca ctgatcaaag accatacact tgtcaattcc ttgattcgtg tgctgcagac 2160
attgcaggag ttcaacatca cgctggagac gtcccttgct aagggtcatc acatctgacc 2220
tcccagcacc agccagcagc aggactgaga aagactcacc ctgcagctct gacctttttt 2280
cccaaaggga cttaaagcat tgtgcaggag taggagacaa aatgtacact cactgtaaaa 2340
agagaactag aggatttttg gaataaataa tctatttttag agttttattg ctgatttgct 2400
ttttacacac tttcatgtga aagagtgata gggagaggga gcgaggctgg tgccgcttat 2460
tttgaagctg gtgccctccc tcgccgtggc cacatgctgg aagcctgagg cctccctgga 2520
ctgagcctgt ggcactgcgt gcgggacagt tatgtttcct tgccccgtcg cattaatgag 2580

```



```
gcccttccac atcattttta aactaatgtt tttctatatt aacattatta tggatatttg 2640
gctttcatag gccacacaca ggtgtgctgc gcgggaagcc ccatgctcca atcaaaggga 2700
tttttagtag tgcctctaag caagcaccga tgagtcagtc ccacgtattt tcttttttgt 2750
cagtattggt tgggaaggag acatgccggg atgtgtcatc gtgccaaata ccacatttcc 2820
tgttggcaca gtttcacaga agtaaacata agcatgtttt aacagggtttt tcttttcttt 2880
tttctttttt aaaatgtttt atttatttaa cccgccattg tgtgttttta agtattttct 2940
ttttttaagg aaaggaaaag cttgtcacia tctaactggc tatgttatta ttattaaatt 3000
tatgttttgc aacttagaaa ccagctacag tatggcccac ttaataaaac acctgaaaca 3060
aaaaaaaaagn nngngggngn nngtnnngag naggagggnn ggngggnggn nngnngggag 3120
ntnanttntg ggggtgngngn ngngnangnt 3150
```

<210> 47

<211> 1806

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2817769CB1

<400> 47

```
gtctgcgcgc aggtgccgct cggcgcccgg gccgcccgtt ccgcggtgtgt cgccgccgtc 60
gtgcgtgccg ctccggcgag gggacggggc tgcgttctct cctccttctt ccccgccctc 120
agctgccggc aggaccttct tctcgctgcc gctgggaccc cgtgtcatcg cccaggccga 180
gcacgatgcc ccctaaaaag ggaggtgatg gaattaaacc accccaatc attggaagat 240
ttggaacctc actgaaaatt ggtattgttg gattgccaaa tgttgggaaa tctactttct 300
tcaatgtgtt aaccaatagt caggcttcag cagaaaactt cccgttctgc actattgatc 360
ctaattgagag cagagtacct gtgccagatg aaagggttga ctttctttgt caataccaca 420
aaccagcaag caaaattcct gccttttctaa atgtgggtga tattgctggc cttgtgaaag 480
gagctcacia tgggcagggc ctgggggaatg cttttttatc tcatattagt gcctgtgatg 540
gcatctttca tctaacacgt gcttttgaag atgatgatc cacgcacgtt gaaggaaagt 600
tagatcctat tcgagatata gaaataatac atgaagagct tcagcttaaa gatgaggaaa 660
tgattggggc cattatagat aaactagaaa aggtggctgt gagaggagga gataaaaaac 720
taaaacctga atatgatata atgtgcaaag taaaatcctg ggttatagat caaaagaaac 780
ctgttcgctt ctatcatgat tggaatgaca aagagattga agtgttgaat aaacacttat 840
ttttgacttc aaaaccaatg gtctacttgg ttaatctttc tgaaaaagac tacattagaa 900
agaaaaacaa atggttgata aaaattaaag agtgggtgga caagtatgac ccagggtgctt 960
tggtcattcc ttttagtggg gccttggaac tcaagttgca agaattgagt gctgaggaga 1020
gacagaagta tctggaagcg aacatgacac aaagtgtctt gccaaagatc attaaggctg 1080
ggtttgcagc actccaacta gaatactttt tcaactgcagg cccagatgaa gtgcgtgcat 1140
ggaccatcag gaaagggact aaggctcctc aggtgcagg aaagattcac acagattttg 1200
aaaagggatt cattatggct gaagtaatga aatacgaaga ttttaaagag gaaggttctg 1260
aaaatgcagt caaggctgct ggaaagtaca gacaacaagg cagaaattat attgttgaag 1320
atggagatat tatcttcttc aaatttaaca cacctcaaca accgaagaag aaataaaatt 1380
tagttattgc tcagataaac atacaacttc caaaaggcat ctgattttta aaaaattaaa 1440
atcttctgaaa accaatgcga caaataaagt tggggagatg ggaatctttg acaacaaat 1500
tatttttatt tgttttaaaa ttaaaatact gtgtaccccc cccactcca tgaaatgcag 1560
gttcactaaa tgtgaacagc tttgcttttc acgtgattaa gaccctactc caaattgtag 1620
aagcttttca ggaaccatat tactctcatg atacttcatt aatctccatc atgtatgcca 1680
agcctgacac atttgacagt gaggacaatg tggcttgctc ctttttgaat ctacagataa 1740
tgcattgttt acagtactcc agatgtctac actcaataaa acatttgaca aaacaaaaaa 1800
aaaaaa 1806
```

<210> 48
<211> 2880
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2917557CB1

<400> 48
gaggaggagg aggaggaaga agaggaagaa gaagatgaag aaagtgaaga agaggaggaa 60
gaggagggag aaagtgaagg cagtgaaggt gatgaggaag atgaaaaggt gtcagatgag 120
aaggattcag ggaagacatt agataaaaag ccaagtaaag aaatgagctc agattctgaa 180
tatgactctg atgatgatcg gactaaagaa gaaagggctt atgacaaagc aaaacggagg 240
attgagaaac ggcgacttga acatagtaaa aatgtaaaca ccgaaaagct aagagcccct 300
attatctgcg tacttgggca tgtggacaca gggaagacaa aaattctaga taagctccgt 360
cacacacatg tacaagatgg tgaagcaggt ggtatcacac aacaaattgg ggccaccaat 420
gttcctcttg aagctattaa tgaacagact aagatgatta aaaattttga tagagagaat 480
gtacggattc caggaatgct aattattgat actcctgggc atgaatcttt cagtaatctg 540
agaaatagag gaagctctct ttgtgacatt gccatttttag ttgttgatat tatgcatggt 600
ttggagcccc agacaattga gtctatcaac cttctcaaat ctaaaaaatg tcccttcatt 660
gttgactca ataagattga taggttatat gattggaaaa agagtcctga ctctgatgtg 720
gctgctactt taaagaagca gaaaaagaat acaaaagatg aatttgagga gcgagcaaag 780
gctattattg tagaatttgc acagcagggt ttgaatgctg ctttgtttta tgagaataaa 840
gatccccgca cttttgtgtc tttggtacct acctctgcac atactggtga tggcatggga 900
agtctgatct accttcttgt agagttaact cagaccatgt tgagcaagag acttgcacac 960
tgtgaagagc tgagagcaca ggtgatggag gttaaagctc tcccggggat gggcaccact 1020
atagatgtca tcttgatcaa tgggcgtttg aaggaaggag atacaatcat tgttcctgga 1080
gtagaagggc ccattgtaac tcagattcga ggccctctgt tacctcctcc tatgaaggaa 1140
ttacgagtga agaaccagta tgaaaagcat aaagaagtag aagcagctca gggggtaaaag 1200
attcttgga aagacctgga gaaaacattg gctggtttac cctccttgt ggcttataaa 1260
gaagatgaaa tccctgttct taaagatgaa ttgatccatg agttaaaagca gacactaaat 1320
gctatcaaat tagaagaaaa aggagtctat gtccaggcat ctacactggg ttctttggaa 1380
gctctactgg aatttctgaa aacatcagaa gtgccctatg caggaattaa cattggccca 1440
gtgcataaaa aagatgttat gaaggcttca gtgatgttg aacatgacct tcagtatgca 1500
gtaatttttg ccttcgatgt gagaattgaa cgagatgcac aagaaatggc tgatagttaa 1560
ggagttagaa tttttagtgc agaaattatt tatcatttat ttgatgcctt tacaaaatat 1620
agacaagact acaagaaaca gaaacaagaa gaatttaagc acatagcagt atttccctgc 1680
aagataaaaa tccctccctca gtacattttt aattctcgag atccgatagt gatgggggtg 1740
acggtggaag caggtcaggt gaaacagggg acacccatgt gtgtcccaag caaaaatttt 1800
gttgacatcg gaatagtaac aagtattgaa ataaaccata aacaagtgga tgttgcaaaa 1860
aaaggacaag aagtttgtgt aaaaatagaa cctatccctg gtgagtcacc caaaatgttt 1920
ggaagacatt ttgaagctac agatattctt gttagtaaga tcagccggca gtccattgat 1980
gcactcaaag actggttcag agatgaaatg cagaagagtg actggcagct tattgtggag 2040
ctgaagaaag tatttgaaat catctaattt tttcacatgg agcaggaact ggagtaaag 2100
caatactgtg ttgtaatatc ccaacaaaaa tcagacaaaa aatggaacag acgtatttgg 2160
acactgatgg acttaagtat ggaaggaaga aaaatagggt tataaaatgt tttccatgag 2220
aaaccaagaa acttacactg gtttgacagt ggtcagttac atgtccccac agttccaatg 2280
tgccgtgttca ctcacctctc ccttcccca ccttctcta cttggctgct gttttaaagt 2340
ttgcccttcc ccaaatttgg atttttatta cagatctaaa gctctttcga ttttatactg 2400
attaaatcag tactgcagta tttgattaac caagcttctg cagattttgt gattcttggg 2460
acttttttga cgtaagaaat acttctttat ttatgcatat tcttcccaca gtgatttttc 2520
cagcattctt ctgccatatg cctttagggc ttttataaaa tagaaaatta ggcattctga 2580
tatttcttta gctgctttgt gtgaaaccat ggtgtaaaag cacagctggc tgctttttac 2640
tgcttgtgta gtcacgagtc cattgtaatc atcacaatc taaaccaaac taccaataaa 2700

gaaaacagac atccaccagt aagcaagctc tgttaggctt ccatgttagt gtagcttctc 2760
 tcccacaagt tgcctccta ggacaagaat tatcttacia actaaactat catcacacta 2820
 ccttgtatgc cagcacctgg taacagtaga gatttttata cattaatctt gatctgtttt 2880

<210> 49

<211> 1109

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3421335CB1

<400> 49

cccacgcgtc cgctcgctct tgggtcatgc ctggccagca gaaagcagct ccatagggga 60
 ggagagccac gcaggatctc acagctgcag tctaatagta acacagagga ttcagcagtg 120
 gccaccatgg gttctgtgaa ttccagaggt cacaaggcgg aagcccaggt ggtgatgatg 180
 ggcctggact cggcgggcaa gaccacgctc ctttacaagc tgaagggcca ccagctgggtg 240
 gagaccctgc cactgtttgg tttcaacgtg gagcctctga aagctcctgg gcacgtgtca 300
 ctgactctct gggacgttgg ggggcaggcc ccgctcagag ccagctggaa ggactatctg 360
 gaaggcacag atactcctgt gtacgtgctg gacagcacag atgaagcccg cttacccgag 420
 tcggcggctg agctcacaga agtctgaac gacccaaca tggctggcgt ccccttcttg 480
 gtgctggcca acaagcagga ggcacctgat gcacttccgc tgcttaagat cagaaacagg 540
 ctgagtctag agagattcca ggaccactgc tgggagctcc ggggctgcag tgccctcact 600
 ggggaggggc tgcccagagg cctgcagagc ctgtggagcc tcctgaaatc tcgcagctgc 660
 atgtgtctgc aggcgagagc ccatggggct gagcgcggag acagcaagag atcttgatcc 720
 agacagagca gcatatcttt gtcatacaa actagaagaa ccagctgatc cttgagaaat 780
 ttacgcttag tctatcaaac aagaaatgct ggcttggccc ggtggctcat gcctgtaatc 840
 ccagcactgt gggagaccac ggtgggggaa tcccttgagc ccaggagtgt gagagcaaca 900
 tcacaacacc ccatttctac taataatcaa aaaattggcc gggcatggtg gcatgtgcct 960
 gtagtcccag ctacttggga ggctgaggca ggagaatcgc ttgagcccaa gaggtagagg 1020
 ttgcagttag ccaagatcgc gccactgcac tccagtctgg gcaacagagt gagaccctgt 1080
 tctagtgggtg ataataataa tgatgtagt 1109

<210> 50

<211> 2407

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 605761CB1

<400> 50

ctttcgctgg cgccattacc tgagttctcc tccagcgttt ccgcaccctc tccgattagc 60
 ggtcccagga gtttccaagg taaccgcgca gtagggcgga tctcattagg cggaaagcga 120
 aacccggaag tgacgtctct accgggtgtc agcagcgaga gggttcgaag atggcggcgc 180
 gcaaggggtcg gcgtcgcacg tgtgaaaccg gggaacccat ggaagccgag tccggcgaca 240
 caagttccga gggcccggcc caggtctacc tgcccggccg ggggcccgcg ctacgcgaag 300
 gggaggagct ggtcatggac gaggaggcct atgtgctcta ccaccgagcg cagactggcg 360
 cccctgtct cagctttgac atagtccggg atcacctggg agacaaccgg acagagcttc 420
 ctcttacact ttacttgtgt gctgggaccc aggttgagag cgcccagagc aacagactga 480
 tgatgcttcg gatgcacaat ctgcatggga caaagccccc accctcagag ggcagtgatg 540

aagaagaaga ggaggaagat gaagaggatg aagaagagcg gaaacctcag ctggagctgg 600
ccatgggtgcc ccactatggt ggcatacaacc gagttcgggt gtcattggctg ggtgaagagc 660
ctgtggctgg ggtgtggtca gagaagggcc aggtggaggt gtttgcgctg cggcggcttc 720
tgcaggtggt ggaggagccc caggccctgg cagccttcct ccgggatgag caggcccaaa 780
tgaagcccat cttctccttc gctggacaca tgggcgaggg ctttgcctt gactggtccc 840
cccgggtgac cggtcgctg ctgaccggtg actgtcaaaa gaacatccac ctctggacac 900
ctacggacgg cggctcctgg cacgtggacc agcggccatt cgtgggccac acacgctctg 960
tggaggacct gcagtgggtca ccgactgaga acacggtgtt tgcctcctgc tcagctgacg 1020
cctccatccg catctgggac atccgggcag cccccagcaa ggcctgcatg ctcaccacag 1080
ccaccgcca tgatggggac gtcaatgtca tcagctggag ccgcccggag cccttcctgc 1140
tcagtggcgg ggatgatggg gccctcaaga tctgggacct tcggcagttc aagtctggtt 1200
ccccagtggc caccttcaag cagcacgtgg ccccggtgac ctccgtcgag tggcaccccc 1260
aggacagcgg ggtctttgca gcctcgggtg cagaccacca gatcacacag tgggacctgg 1320
cagtggagcg ggacctgag gcgggcgacg tggaggccga ccccgactg gccgacctcc 1380
cgcagcagct gctgttcgtg caccagggcg agaccgagct gaaggagctg cactggcacc 1440
cgcagtggcc agggctcctg gtcagcacgg cgtgtcagg cttcaccatc ttccgcacca 1500
tcagcgtctg aggcgtccca ctggctctga tcttgcttcc tgcttggaag ctgaagtcga 1560
attgggctcc cctggaaggg gttcattcag gtctgttgac tgagactggc cggcctgtgg 1620
gctgccgtga tggattctgt ttgacgtatt gttctctaga aggcctggct ctgatccagt 1680
gacctcttc accaaagaac tcggtttaac cagggtcttg taagaccact cccaccaga 1740
gacttgtgtg gcctggtgtg gcctgtgtgt cggattcctt cctgtcagct gtgacctt 1800
tgacctgtgt cccagaacc cagttttttg tttgtttgtt tgagacggag tcttggtctg 1860
tcgcccaggc tggagtgcag tagcacgac ttggctcact gcaacctccg cctcctgggt 1920
taaagtgatt ctctcagctc agtctcccag gtagctggga ttacaggcat gtgccaccac 1980
acccggttaa tttttgtatt tttagtagag acggggtttc accatgttgg ccaggctgg 2040
ctcaaattct tgatctcaag tgatctgtcc gcccggcct cccagagtgc tgggttggga 2100
ttacaggcgt gagccaccgc gtccggctca ggaccagtt ttggctgctg gttcccagca 2160
ggggactcgg gggatataca gtggctgcac caaattggag gtgtgggttc ctccaacaca 2220
atgtgtctct gccggtgtc ttccgtccag ctgggtttgg ccaggatttc tccgtgtggg 2280
ggctacatgc gacctctcc cctcctcct gacttttagag gctggtgctg tgcgggagg 2340
aaggtcaggg ctctgagca gcaataaagg accaggaaga ggcctgaggt gtaaaaaaaa 2400
aaaaaaa 2407

<210> 51

<211> 1158

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 483862CB1

<400> 51

ggaagaccgt cccggatggc ctccgggact gccagtgtgt ggaggtgagc tccgggattg 60
ccggcattcc cgcttctgct ggttgcttca tgctgcaggc tgccggccgtc agccctcgtc 120
cgcatgtgtg gcgctgaggt gccggggcag caagtgcacat gtcgtcgggc ctccgcgccg 180
ctgacttccc ccgctggaag cgccacatct cggagcaact gaggcgccgg gaccggctgc 240
agagacaggc gttcgaggag atcatcctgc agtataacaa attgctggaa aagtcagatc 300
ttcattcagt gttggcccag aaactacagg ctgaaaagca tgacgtacca aacaggcacg 360
agataagtc cggacatgat ggcacatgga atgacaatca gctacaagaa atggcccaac 420
tgaggattaa gcaccaagag gaactgactg aattacacaa gaaacgtggg gagttagctc 480
aactggtgat tgacctgaat aaccaaattgc agcgggaagg caggagatg cagatgaatg 540
aagcaaaaat tgcagaatgt ttgcagacta tctctgacct ggagacggag tgcctagacc 600
tgcgcaactaa gctttgtgac cttgaaagag ccaaccagac cctgaaggat gaatatgatg 660

```

ccctgcagat cacttttact gccttggagg gaaaactgag gaaaactacg gaagagaacc 720
aggagctggt caccagatgg atggctgaga aagcccagga agccaatcgg cttaatgcag 780
agaatgaaaa agactccagg aggcggcaag cccggctgca gaaagagctt gcagaagcag 840
caaaggaacc tctaccagtc gaacaggatg atgacattga ggtcattgtg gatgaaactt 900
ctgatcacac agaagagacc tctcctgtgc gagccatcag cagagcagcc acgtaagtag 960
gcaggtttgg gccagggaaa agacagcttg aggagcaata tgaaggcaca tctgtggaca 1020
tgacaaagaa tgcagtcaga tgcacccaac cccttactcc ttttctggga caccagcgt 1080
cgaacacacc acagaggtgt ctagtctttc tcagttcacc tctgcttaat gggagggaag 1140
cagaacacgg gtggcttc                                     1158

```

<210> 52

<211> 1026

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1256777CB1

<400> 52

```

ctgcgggect ctctccgtcg ccatggaaac gaaagcggcc aagtagagct ccgtcctgac 60
gcgcccgcct ccgtgggctc cggccggcta agccgcggcg gacaactatg ctgaaagcca 120
agatcctctt cgtggggcct tgcgagagt gaaaaactgt tttggccaac tttctgacag 180
aatcttctga catcactgaa tacagcccaa cccaaggagt gaggatccta gaatttgaga 240
acccgcatgt taccagcaac aacaaaggca cgggctgtga attcgagcta tgggactgtg 300
gtggcgatgc taagtgtgag tcctgctggc cggccctgat gaaggatgct catggagtgg 360
tgatcgtctt caatgctgac atcccaagcc accggaagga aatggagatg tggatttcct 420
gctttgtcca acagccgtcc ttacaggaca cacagtgtat gctaattgca caccacaaac 480
caggctctgg agatgataaa ggaagcctgt ctttgtcgcc acccttgaac aagctgaagc 540
tgggtgcactc aaacctggaa gatgaccctg aggagatccg gatggaattc ataaagtatt 600
taaaaagcat aatcaactcc atgtctgaga gcagagacag ggaggagatg tcaattatga 660
cctagccagc cttcacctgg gactgccaca tccccagtga aatcagcatg tttctcggtg 720
cagatctgaa atcacatcca gctcctgatg ttttcttctc cctctgactg cagaggaagt 780
gttcctacct gcaggaaggc acctgtcaca cagggcgttc actcagacca tctgtgctct 840
gccctgagtt cagttgagaa aatcctatta tcaaatttgg atttcctggc cccagaactt 900
cccaaagacc tgtaaaatgg agggatttac cacctcacat atgtccagtt aaacagtttg 960
tggacttgta accgtcgcag cccaatgata caacagtagt ttaatcacgt gaaaaaaaaa 1020
aaaaaa                                     1026

```

<210> 53

<211> 2456

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2198779CB1

<400> 53

```

gacgagcgac gagatgacgg aggagcgggt ggccaacgca tgccggcagc cgggtgtaaac 60
aaggcctcgc gccgctgcgg gtctctgcgac cgctcctggc tgggatttcg attggctcgg 120
cagagaggtt acctggaaat ccaacaccgc ccaacacccc tcccgtccc cagtcggggg 180
acttcgatag gattggagaa ggagttgaca ggaggagccc ccgcacagga cctaagaatg 240

```



```

ctgtgaccag aagatgggat cgcggaacag cagcagtgca ggatccgggt ccggagaccc 300
ctccgagggc ttgccccgaa gaggggctgg cctgcgtcgg agtgaggaag aggaagaaga 360
ggatgaagat gtggatctgg cccaggtact ggcctatctc ctccgcagag gccaagtga 420
gttgggtgcag ggaggagggtg cagcaaattt acaattcatt caggccctct tggactcaga 480
ggaagagaat gacagagctt gggatggtcg tcttggggat cgatacaacc cacctgtgga 540
tgctaccctt gacacccggg agctggaatt caatgagatc aagacacaag tggactggc 600
cacagggcag ctggggctta ggcgggcccgc ccagaagcac agctttcctc gaatgttgca 660
ccagagagaa cggggcctct gccatcgggg aagcttctcc cttggagaac agtctcgagt 720
gatatctcac ttcttgccca atgatctggg cttcactgat agctactctc agaaggcttt 780
ctgtggcatc tacagcaaag atgggtcaa atcatgtct gcttgccaag accagacaat 840
ccgactctat gactgccgat atggccgttt ccgtaaattc aagagcatca aggcccgca 900
tgtaggctgg agcgtcttgg atgtggcctt caccctgat gggaaacct tctctactc 960
tagctggtct gattacattc atatctgcaa tatctatggg gagggagata cacacactgc 1020
cctggatctc aggccagatg agcgtcgctt tgctgtcttc tccattgctg tctcctcaga 1080
tggacgagaa gtactaggag gggccaatga tggctgcctg tatgtctttg accgagaaca 1140
gaaccggcgc acccttcaga ttgagtccca tgaggatgat gtgaatgcag tggcctttgc 1200
tgatataagc tcccaaattc tgttctctgg gggagatgat gccatctgca aagtgtggga 1260
tcgacgcacc atgcgggagg atgaccccaa gcctgtgggt gcaactggctg gacaccagga 1320
tggcatcacc ttcattgaca gcaagggtga tgcccgggat ctgatctcca actctaaaga 1380
ccagaccatc aaactctggg atatccgacg cttttccagc cgggaaggca tggaaagctt 1440
acgccaggct gccacacagc aaaactggga ctatcgggtg cagcaagtgc ccaaaaaagg 1500
gtttactctg catccctacc cagcctggcg gaagctgaag ctcccagggg acagctcctt 1560
gatgacctac cggggccacg gagtgtgca caccctcatc cgctgccggg tctcccccat 1620
tcatagcact ggccagcagt tcatctacag tggctgctcc actggcaaag tggttgtgta 1680
cgaccttcta agtggccaca ttgtgaagaa gctgaccaac cacaaggcct gtgtgcgtga 1740
cgtcagttgg cacccttttg aagagaagat tgtcagcagt tcgtgggacg ggaacctgcg 1800
tctgtggcag taccgccagg ctgagtactt ccaggatgac atgccagaat ctgaggaatg 1860
tgccagcgcc cctgccccag tgccccaatc ctctacaccc ttttcctcac ccagtagat 1920
ccaacctcca gcccatata ggggtgaacct cttgataagc tctctgcctc ctctccctt 1980
tctcccttgt ggggaatgtt tggaggaatc actggcattt gatggggaat aacataagcc 2040
tgggctctga gcctcagctg agccttgaa gattctcccc atggggcaga gtggtctcct 2100
tacgtgctca caccagtcg gcttgggtcc ctatctctgg ccagagtttg gcaggactgc 2160
cattatctgg ggtgtggcct ctgccagcaa gagaagtgtc ctgggtgttt ttaatcatgt 2220
ttgaatgtta ggggttgat cctagagtag atgcctgagg ccacatctga acagacctgt 2280
cagccaggcc tgccaggtct tcacgttgag gattcaactg gccaatcaca ggacaggtgt 2340
cctggccttt ctctctgagg tctctagggg aggggcatgg gtaagggtgt tctctcagca 2400
ccctcctggg gtggggatta tgtctgctgt catgtctggg tctttaaggt aggaca 2456

```

<210> 54

<211> 1771

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2226116CB1

<400> 54

```

cggctcgagt taatatatttc gttgagcgga accttgctat tccataagag gatgtgtcca 60
gtgttggtga agatttcatg ttttaaattc tttgtacaga aatcctgctc ccaagtcaca 120
gataggctga cgggtcagag ggcaagacgt gaccagggc cgagagggtg agtgaccagg 180
aaaatcggat tcatcagttc acttgtttgt ttcagaaacg tgcacaaaga cctgctgcat 240
gagggcctcg tcttcagttt ctgtttcatg cccagcatta aaccaagtat ctcattttgc 300
caatttgact tctgtagggg ccattggcacc tgcaagggtg ttctcagcaa gattgaggac 360

```



```
cgtgtttcag ggcgtggggc attgggcttt gtccacatgg gctggcctga agcccagccg 420
gctactgcca cagcgggctt ctcccaggct gctctcggtc ggccgtgcgg acctcgccaa 480
gcatcaggaa ctcccgggga agaagctgct ctctgagaaa aagctgaaaa ggtactttgt 540
ggactatcgg agagtgttg tctgtggagg aaacggaggc gctggggcaa gctgcttcca 600
cagtgaagccc cgaaggagt ttggaggccc tgatggaggg gacggaggca acggtggaca 660
cgtcattctg agagttgacc agcaagtcaa gtccctgtcg tcggtcctgt cgcggtacca 720
gggtttcagt ggagaagatg gagggagtaa aaactgcttc gggcgcagtg gcgccgtcct 780
ctacatccgg gtccccgtgg gcacgctggg gaaggaggga ggcagagttg tggccgacct 840
gtcttgctg ggagatgagt acattgccgc gctgggcggg gcaggaggga aaggcaaccg 900
cttcttcctg gccaaacaaca accgtgcccc tgtgacctgt acccctggac agccaggaca 960
gcagcgagtt ctccacctgg agctcaagac ggtggccccc gccggaatgg tgggattccc 1020
caacgccggg aagtcctcac tgetccgggc catttcaaac gccagaccgg ccgtggcttc 1080
ctaccggttc accacctga agccccacgt cgggatcgtc cactacgaag gccacctaca 1140
aatagcagtg gccgacatcc ccggcatcat acgaggcgcc caccagaaca ggggtctggg 1200
gtccgccttc ctccaggaca tcgagcgctg ccgctttctc ttgttcgtgg tggatctttc 1260
tcagcctgag ccgtggactc aagttgacga tttaaaatat gaactggaga tgtatgaaaa 1320
gggcctgtct gcgaggcccc acgcaatcgt cgcaaacaag attgacctcc ctgaagccca 1380
agccaatctg tcccagctcc gggatcactt gggacaggag gtcacgtgac tgtcggcggt 1440
gaccggcgag aacctggagc agctgctggt gcacctgaag gtgctgtatg acgcctacgc 1500
ggaggccgag ctgggcccagg gccgccagcc gctcaggtgg tagccacgcc agagcggggg 1560
cgctctggg cctctgtctg agcaaacctg ggtgtgaatt cgggtggtttt gaatgcataa 1620
agtgccttgt ggacacgggg gagttgtggt gcttctgggt ctctgggccc cgctgctgg 1680
cctgagatgc cctcatgttg ggaagcatte cgtgcccccc accccgcctg ccctccgtat 1740
ttcctgcacc tgtcagcctg cgctgactga t 1771
```

<210> 55

<211> 2724

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2504472CB1

<400> 55

```
gctgaccagt tggcgacatg gtggcacccg tgctggagac ttctcacgtg ttttgctgcc 60
caaaccgggt gcggggagtc ctgaactgga gctctgggcc cagaggactt ctggcctttg 120
gcacgtcctg ctccgtgggt ctctatgacc ccctgaaaag ggttggtgtt accaacttga 180
atggtcacac cgcccagatc aattgcatac agtggatttg taaacaggat ggctcccctt 240
ctactgaatt agtttctgga ggatctgata atcaagtgat tcaactggaa atagaggata 300
atcagctttt aaaagcagtg catcttcaag gccatgaagg acctgtttat gcggtgcatg 360
ctgtttacca gaggaggaca tcagatcctg cattatgtac actgatcgtt tctgcagctg 420
cagattctgc tgttcgactc tggctctaaa aggggtccaga agtaatgtgc cttcagactt 480
taaactttgg aaatggattt gctttggctc tctgcttata ttttttgcca aatactgatg 540
taccaatatt agcatgtggc aatgatgatt gcagaattca catatttgct caacaaaatg 600
atcagtttca gaaagtgtt tctctctgtg gacatgagga ttggattaga ggagtggaa 660
gggcagcctt tggtagagat cttttcctag caagctgttc acaagattgc ctgataagaa 720
tatggaagct gtatataaag tcaacatctt tagaaactca ggatgacgat aacataagac 780
tgaaagaaaa tacttttacc atagaaaatg aaagtgttaa aatagcattt gctgttactc 840
tgagacagat gctagccggg catgaaaact gggtaaatgc agttcactgg caacctgtgt 900
tttaciaaga tgggtgtccta cagcagccag tgagattatt atctgcttcc atggataaaa 960
ccatgattct ctgggctcca gatgaagagt caggagtttg gctagaacag gttcgagtag 1020
gtgaagtagg tgggaatact ttgggatttt atgattgcca gttcaatgaa gatggctcca 1080
tgatcattgc tcatgctttc cacggagcgt tgcacctttg gaaacagaat acagttaacc 1140
```

```

caagagagtg gactccagag attgtcattt caggacactt tgatgggtgtc caagacctag 1200
tctgggatcc agaaggagaa tttattatca ctgttggtac tgatcagaca actagacttt 1260
ttgctccatg gaagagaaaa gaccaatcac aggtgacttg gcatgaaatt gcaaggcctc 1320
agatacatgg gtatgacctg aaatgttttg caatgattaa tcggtttcag tttgtatctg 1380
gagcagatga aaaagttctt cgggtttttt ctgcacctcg gaattttgtg gaaaattttt 1440
gtgccattac aggacaatca ctgaatcatg tgctctgtaa tcaagatagt gatcttccag 1500
aaggagccac tgtccctgca ttgggattat caaataaagc tgtcttttcag ggagatatag 1560
cttctcagcc ttctgatgaa gaggagctgt taactagtac tgggttttgag tatcagcagg 1620
tggcctttca gccctccata cttactgagc ctcccactga ggatcatctt ctgcagaata 1680
ctttgtggcc tgaagttcaa aaactatatg ggcacggtta tgaaatatatt tgtgttactt 1740
gtaacagttc aaagactctg cttgcctcag cttgtaaggc agctaagaaa gagcatgcag 1800
ctatcattct ttggaacact acatcttggg aacagggtgca gaatttagtt tccacagtt 1860
tgacagtcac gcagatggcc ttctcaccta atgagaagtt cttactagct gtttccagag 1920
atcgaacctg gtcattgtgg aaaaagcagg atacaatctc acctgagttc gagccagttt 1980
ttagtctttt tgccttcacc aacaaaatta cttctgtgca cagtagaatt atttggtctt 2040
gtgattggag tcctgacagc aagtatttct tctactgggag tcgagacaaa aagggtggtt 2100
tctggggtga gtgcgactcc actgatgact gtattgagca caacattggc cctgctcct 2160
cagtccctga cgtgggtggg gctgtgacag ctgtcagcgt ctgccagtg ctccacctt 2220
ctcaacgata cgtggttgca gtaggattgg agtgtggaaa gatttgctta tatacctgga 2280
aaaagactga tcaagttcca gaaataaatg actggacca ctgtgtagaa acaagtcaaa 2340
gccaaagtca tacactggct atcagaaaat tatgctggaa gaattgcagt ggaaaaactg 2400
aacagaagga agcagaaggt gctgagtggt tacactttgc aagctgtggt gaagatcaca 2460
ctgtgaagat acacagagtc aataaatgtg cactgtaatg gacttaataa ctacatgctt 2520
gcagtcactg gtatctttaa atattatcat gtaaacaggt catctttacc ttcataactg 2580
aattgagttt ctgggttttt tttttttttg agatggagtc ttgctttgtc acaacctcca 2640
cctcccaggt tcaagcgatt ctctttcttc agcctcctga gtagctggga ctagaggcac 2700
accaccatgc ccggetaatt tttg

```

<210> 56

<211> 2963

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3029920CB1

<400> 56

```

ggccgaagag gctggcaggt ggcgccgtgg ggtgggtgct cctgggtgaga ggagtccact 60
ccgtgcgtgc gggcggaggc cggcccccca gagccgccga catgaagaaa gacgtgcgga 120
tcctgctggt gggagaaact agagttggga agacatcact gattatgtct ctggtcagtg 180
aagaatttcc agaagaggtt cctccccggg cagaagaaat caccattcca gctgatgtca 240
ccccagagag agttccaaca cacattgtag attactcaga agcagaacag agtgatgaac 300
aacttcatca agaaatatct caggctaatt tcatctgtat agtgtatgcc gttaacaaca 360
agcattctat tgataaggta acaagtcgat ggattcctct cataaatgaa agaacagaca 420
aagacagcag gctgccttta atattggttg ggaacaaatc tgatctggtg gaatatagta 480
gtatggagac catccttctt attatgaacc agtatacaga aatagaaacc tgtgtggagt 540
gttcagcgaa aaacctgaag aacatatcag agctctttta ttacgcacag aaagctgttc 600
ttcatcctac agggcccttg tactgcccag aggagaagga gatgaaacca gcttgtataa 660
aagcccttac tcgtatatatt aaaatatctg atcaagataa tgatgggtact ctcaatgatg 720
ctgaactcaa cttcttttcag aggattttgt tcaacactcc attagctcct caagctctgg 780
aggatgtcaa gaatgtagtc agaaaacata taagtgtagg tgtggctgac agtgggttga 840
ccctgaaagg ttttctcttt ttacacacac tttttatcca gagagggaga cacgaaacta 900
cttggactgt gcttcgacga tttgggttat atgatgacct ggatttgaca cctgaatatt 960

```

```

tggtccccct gctgaaaata cctcctgatt gcactactga attaaatcat catgcatatt 1020
tattttctcca aagcaccttt gacaagcatg atttgatag agactgtgct ttgtcacctg 1080
atgagcttaa agattttattt aaagttttcc cttacataacc ttgggggcca gatgtgaata 1140
acacagtttg taccaatgaa agaggctgga taacctacca gggattcctt tcccagtgga 1200
cgctcacgac ttatttagat gtacagcggg gcctggaata ttggggctat ctaggctatt 1260
caatattgac tgagcaagag tctcaagctt cagctgttac agtgacaaga gataaaaaga 1320
tagacctgca gaaaaaacia actcaaagaa atgtgttcag atgtaatgta attggagtga 1380
aaaactgtgg gaaaagtgga gttcttcagg ctcttcttgg aagaaactta atgaggcaga 1440
agaaaattcg tgaagatcat aaatcctact atgcgattaa cactgtttat gtatatggac 1500
aagagaaata cttgttggtg catgatattc cagaatcgga atttctaact gaagctgaaa 1560
tcatttgtga tgttgatatg ctggtatatg atgtcagcaa tcccaaattc tttgaatact 1620
gtgccaggat ttttaagcaa cactttatgg acagcagaat accttgctta atcgtagctg 1680
caaagtcaga cctgcatgaa gttaaacaag aatacagtat ttcacctact gatttctgca 1740
ggaaacacaa aatgcctcca ccacaagcct tcacttgcaa tactgctgat gccccagta 1800
aggatatctt tgttaaattg acaacaatgg ccatgtatcc gcacgtgaca caagctgacc 1860
tcaagagctc cacgttttgg ctccgagcaa gttttggtgc tactgttttt gcagttttgg 1920
gctttgctat gtacaaagca ttattgaaac agcgatgata taaaaagaaa tactgtccct 1980
acaaaaaaca aatactttta tgtacattct gaatgcttta agttctgcta gaattattga 2040
gatatttata catgcagagt tactttatta atatttgtaa ttcatgcata agagtatttt 2100
aatgatagtt ataactgcag tattggctag catatggaaa gaaaacagct aacagccaaa 2160
ctaaaatggc taaattccag aggccaaaag ggaatatttt gtaaataatat gtacatattc 2220
aggcaagata tggctctcca agctgagttc tagaaatgat gtttctagac atttctaagt 2280
ggattattga gtgctcactt ggctcactct tctaggttta agttagccca gagattgtat 2340
ttactcatgg atcactttat ttatttcaca ttactcaga atgattcctt gggttctata 2400
aggacataag gtacaatttg ccattgtctc tccattttta aaaacataca agtcagtgtc 2460
agcttaccaa catgacattt tttcagtcag ttgtggtagg ccagccttga agccatcgca 2520
cagtctagaa acttgtgtag ctgagtggtc agctcacctt taagggtgaa gttaggtaaa 2580
agcaattagc agaggcgtta tctatgtgat tatgttgctt ccttgctcagt atgttgaatt 2640
ttatagccct ttcaatgaaa taaaaaaaaa atttgtatat taccaatgtt tttagttaa 2700
ataaagagtc acccttacta ctgttgaatt tcatcccaag tgtaaatcat tctataatgg 2760
ctgtgtctgt tatagtatat tacagtaact gcatgtgtca ccaagtgttc tatatcaggc 2820
taggataacc tagaggcagt aattttttta atgataaaat aaatctaata aatataaact 2880
ctcatgataa acctattttt tccatcatca gccttttcaa gtatttaaata aaataactgc 2940
tgtgtactgt gaaaaaaaaa aaa 2963

```

<210> 57

<211> 3332

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 3332415CB1

<400> 57

```

gcctggcaga ggctggcggg catcgtgccc gtccctgccg gtctcccggg caccggcca 60
ccgccccacc ccctcctccc tgccatggag cccgagctgg acgctcagaa gcagcctcga 120
ccgcgaggagc gaagccgccc ggcctctggg ctccagcagg agggagcgac ggggccttcg 180
gccgacacct ccgggtcggg gctggacggg agatgttccc ttcggagagg cagctccttc 240
acattcttaa cacctggccc caactgggac ttcactttga aaagaaaacg cagagagaaa 300
gatgatgatg ttgtaagcct tagcagcctt gatctgaagg agccaagcaa taaaagagtt 360
cgacctctgg ctggtgtcac gtccttgcca aatttaattc ctctgtgaag aaatggagct 420
gtcagacgtt ttgggtcaaac aatacagtc tttacccttc gtggtgacca cagatcccca 480
gcctctgccc agaagttttc tagcaggtca acagtcccaa caccggcca gagaggagc 540

```

agtgcaactgt ggtcagagat gctggacatc accatgaagg agtctctcac caccagggag 600
atcagacggc aggaggcaat atatgaaatg tcccgagggtg aacaggattt aattgaggat 660
ctcaaacttg caagaaaggc ctaccatgac cccatgttaa agttgtccat catgtcagaa 720
gaggaactca cacatatatt tgggtgatctg gactcttaca tacctctgca tgaagatttg 780
ttgacaagaa taggagaagc aaccaagcct gatggaacag tggagcagat tggtcacatt 840
ctcgtgagct ggttaccgcg cttgaatgcc tacagagggtt actgtagtaa ccagctggca 900
gccaaagctc ttcttgatca aaagaaacag gatccaagag tccaagactt cctccagcga 960
tgtctcgagt ctcccttcag tcgaaaacta gatctttgga gtttcctaga tatccctcga 1020
agtcgcctag tcaaataccc tttactgtta aaagaaattc ttaaacacac tccaaaagag 1080
caccctgatg ttcagcttct ggaggatgct atattgataa tacagggagt cctctctgat 1140
atcaacttga agaaaggtga atccgagtgc cagtattaca tcgacaagct ggagtacctg 1200
gatgaaaagc agagggaccc cagaatcgaa gcgagcaaag tgcctgctgtg ccatggggag 1260
ctgcggagca agagtggaca taaactttac attttctgtt ttcaagacat cttgggttctg 1320
actcggccccg tcacacggaa cgaacggcac tcttaccagg tttaccggca gccaatccca 1380
gtccaagagc tagtcctaga agacctgcag gatggagatg tgagaatggg aggctccttt 1440
cgaggagctt tcagtaactc agagaaaagc aaaaatatct ttagaattcg cttccatgac 1500
ccctctccag cccagtctca cactctgcaa gccaatgacg tgttccacaa gcagcagtgg 1560
ttcaactgta ttcgagcggc cattgcccc ttccagtcgg caggcagtc acctgagctg 1620
cagggcctgc cggagctgca cgaagagtgt gaggggaacc acccctctgc gaggaaactc 1680
acagcccaga ggagggcatc cacagtttcc agtggtactc aggtagaagt tgatgaaaac 1740
gcttacagat gtggctctgg catgcagatg gcagaggaca gcaagagctt aaagacacac 1800
cagacacagc cgggcatccg aagagcgagg gacaaagccc tttctggtgg caaacggaaa 1860
gagactttgg tgtagagaag gctctgtgtg ttaactgatg ggagagactg tttgtttata 1920
aatgtgtaca gttttgtttt ctctgaaggg gagcatcata gggttacttt ataccagttg 1980
taacattttc attgtttttg gttgttcttt tttcttttt taatggcagc taaagatata 2040
cagattactg ttaaattgca gtcctttttt ttttaaagat attttcttga gttatttaga 2100
acatggtaag cctggtattt tttaatcaaa caaaatatat atgaaatggg ttttctctta 2160
attctggatt catcatggct ttctaatacc aattgtata tttacaatat tcaccaaacc 2220
ttagaatttt gcaaagtctg gaattctgcc agtggttctt tgctaagcct tgcattgcaa 2280
atttgaaatt ttaacattgg cacccaaac ctacatggaa tgtatgtctg gagtatttca 2340
aactttacat tgaaacataa tttccttgga aaacaaacca taagcctgag gaggttttta 2400
tcaactggaa tgctttatat tagtttgttt ttcactgtac attcctcatt ttacattcat 2460
ttaacctgcc gattatttaa tttttttatt gtaaagtagt ttttagcatt tgcttttatt 2520
tttttacttt gatgcctttt caaattggca tgtctttaa gtatttttct tcctgattaa 2580
aaatgtgtgt gtatgtgtgt gtgtgtgtgt atatatatat atttttttta atcacattaa 2640
ttttaccaag tgaaaccaag ccatactgtt tttgagccaa ttaagaaaat tgccattttt 2700
aaagtgtagc atttcaggg aaagacccat gaaatggctt gatgtattct agactactga 2760
aagaaaacca cttcaaagat tttgttgaaa gtttttagtgt tgtctgaaat gcaagaggga 2820
aggtgattgg tagtgagtta aaagaaaaag agaggaaaag agagtagttt tgtcttcaag 2880
taaaatgtct gggtgtgcca gacatttcac aagtgtgaaa ggagatagga gaagctcaac 2940
ttgagggcgt gtagtaagtt gtagaaggct cgagggggacg tggacttatt tgccttggtt 3000
tgcaatacct gcaaataatg agtttgaaaa gaaacaatga aatgtgttaa aaatttgacc 3060
atattagata aattttggtg gatttagtca taagatggaa aaagactggg gaatctttta 3120
ttacaaaatg tttctgttaa aatgggatca tcatctttga aaggggggag gaggagtaaa 3180
agcccgatta taatggtgat caattcaagt cagtgttgac tattctgtga aatatatttg 3240
gccagtggaa atgataatca gaaaagactg taaatagatc catccaaatg atttctctgt 3300
acaaatgaat gatactatta aaaaaaaaaa aa 3332

<210> 58

<211> 2617

<212> DNA

<213> Homo sapiens

<220>

<223> Incyte ID No: 4031536CB1

<400> 58

tttagtaatg	tgcctgtatt	acatgtagag	agtattcgtc	aaccaagagg	agtttttaaaa	60
tgtcaaaacc	gggaaaacct	actctaaacc	atggcttggg	tcctgttgat	cttaaaagt	120
caaaagagcc	tctaccacat	caaactgtga	tgaggatatt	tagcattagc	atcattgccc	180
aaggcctccc	tttttgtcga	agacggatga	aaagaaagtt	ggaccatggg	tctgaggtcc	240
gctctttttc	tttgggaaag	aaaccatgca	aagtctcaga	atatacaagt	accactgggc	300
ttgtaccatg	ttcagcaaca	ccaacaactt	ttggggacct	cagagcagcc	aatggccaag	360
ggcaacaacg	acgccgaatt	acatctgtcc	agccacctac	aggcctccag	gaatggctaa	420
aaatgtttca	gagctggagt	ggaccagaga	aattgcttgc	tttagatgaa	ctcattgata	480
gttgtgaacc	aacacaagta	aaacatatga	tgcaagtgat	agaaccccag	tttcaacgag	540
acttcatttc	attgctccct	aaagagttgg	cactctatgt	gctttcattc	ctggaaccca	600
aagacctgct	acaagcagct	cagacatgtc	gctactggag	aattttgggt	gaagacaacc	660
ttctctggag	agagaaatgc	aaagaagagg	ggattgatga	accattgcac	atcaagagaa	720
gaaaagtaat	aaaaccaggt	ttcatacaca	gtccatggaa	aagtgcatac	atcagacagc	780
acagaattga	tactaactgg	aggcgaggag	aactcaaadc	tcctaagggtg	ctgaaaggac	840
atgatgatca	tgtgatcaca	tgcttacagt	tttgtggtaa	ccgaatagtt	agtggttctg	900
atgacaacac	tttaaaagtt	tggtcagcag	tcacaggcaa	atgtctgaga	acattagtgg	960
gacatacagg	tggagtatgg	tcatacacia	tgagagacaa	catcatcatt	agtggatcta	1020
cagatcggac	actcaaagt	tggaatgcag	agactggaga	atgtatacac	accttatatg	1080
ggcatacttc	cactgtgcgt	tgtatgcac	ttcatgaaaa	aagagttgtt	agcggttctc	1140
gagatgccac	tcttaggggt	tgggatattg	agacaggcca	gtgtttacat	gttttgatgg	1200
gtcatgttgc	agcagtccgc	tgtgttcaat	atgatggcag	gaggggtgtt	agtggagcat	1260
atgattttat	ggtaaagggt	tgggatccag	agactgaaac	ctgtctacac	acgttgcagg	1320
ggcatactaa	tagagtctat	tcattacagt	ttgatgggat	ccatgtgggtg	agtggatctc	1380
ttgatacatc	aatccgtgtt	tgggatgtgg	agacagggaa	ttgcattcac	acgttaacag	1440
ggcaccagtc	gttaacaagt	ggaatggaac	tcaaagacaa	tattcttgtc	tctgggaatg	1500
cagattctac	agttaaaatc	tgggatattca	aaacaggaca	gtgtttacaa	acattgcaag	1560
gtcccaacaa	gcatacagat	gctgtgacct	gtttacagtt	caacaagaac	tttgtaatta	1620
ccagctcaga	tgatggaact	gtaaaactat	gggacttgaa	aacgggtgaa	tttattcgaa	1680
acctagtac	attggagagt	ggggggagtg	ggggagttgt	gtggcggatc	agagcctcaa	1740
acacaaagct	ggtgtgtgca	gttggggagtc	ggaatgggac	tgaagaaacc	aagctgctgg	1800
tgctggactt	tgatgtggac	atgaagtga	gagcagaaaa	gatgaatttg	tccaattgtg	1860
tagacgatat	actccctgcc	cttccccctg	caaaaagaaa	aaaagaaaag	aaaaagaaaa	1920
aaatcccttg	ttctcagtgg	tgcaggatgt	tggcttgggg	caacagattg	aaaagacctt	1980
cagactaaga	aggaaaagaa	gaagagatga	caaaccataa	ctgacaagag	aggcgtctgc	2040
tgtctcatca	cataaaaggc	ttcacttttg	actgagggca	gctttgcaaa	atgagacttt	2100
ctaaatcaaa	ccaggtgcaa	ttattttctt	attttcttct	ccagtgggtc	ttgggcagtg	2160
ttaatgctga	aacatcatta	cagattctgc	tagcctgttc	ttttaccact	gacagctaga	2220
cacctagaaa	ggaactgcaa	taatatacaa	acaagtactg	gttgactttc	taattagaga	2280
gcatactgaa	caaaaagtca	tttttctgga	gtggaaaagc	ttaaaaaaat	tactgtgaat	2340
tgtttttgta	cagttatcat	gaaaagcttt	tttttttttt	tttgccaacc	attgccaatg	2400
tcaatcaatc	acagtattag	cctctgttaa	tctatttact	gttgcttcca	tatacattct	2460
tcaatgcata	tgttgctcaa	aggtggcaag	ttgtcctggg	ttctgtgagt	cctgagatgg	2520
atttaattct	tgatgctggg	gctagaagta	ggtcttcaaa	tatgggattg	ttgtcccaac	2580
cctgtactgt	actcccagtg	gccaaactta	tttatgc			2617



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, 16/18, A61K 38/17, G01N 33/68		A3	(11) International Publication Number: WO 00/31263 (43) International Publication Date: 2 June 2000 (02.06.00)	
(21) International Application Number: PCT/US99/28013 (22) International Filing Date: 23 November 1999 (23.11.99) (30) Priority Data: 60/109,592 23 November 1998 (23.11.98) US 60/118,610 4 February 1999 (04.02.99) US 60/127,990 6 April 1999 (06.04.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 60/109,592 (CIP) Filed on 23 November 1998 (23.11.98) US 60/118,610 (CIP) Filed on 4 February 1999 (04.02.99) US 60/127,990 (CIP) Filed on 6 April 1999 (06.04.99) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive, #12, Mountain View,			CA 94040 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). YANG, Junming [CN/US]; 7136 Clarendon Street, San Jose, CA 95129 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 14 September 2000 (14.09.00)	
(54) Title: GTPASE ASSOCIATED PROTEINS				
(57) Abstract <p>The invention provides human GTPase associated proteins (GTPAP) and polynucleotides which identify and encode GTPAP. The invention also provides expression vectors, host cells, antibodies, agonist, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of GTPAP.</p>				

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/28013

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C07K16/18 A61K38/17 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MOOSLEHNER K ET AL: "STRUCTURE AND EXPRESSION OF A GENE ENCODING A PUTATIVE GTP-BINDING PROTEIN IDENTIFIED BY PROVIRUS INTEGRATION IN A TRANSGENIC MOUSE STRAIN" MOLECULAR AND CELLULAR BIOLOGY 1991, vol. 11, no. 2, 1991, pages 886-893, XP000891270 ISSN: 0270-7306 abstract; figure 1	1-12
A	WO 98 37196 A (LUDWIG INST CANCER RES) 27 August 1998 (1998-08-27) abstract; claims 1-52; examples 1-8	1-20
A	WO 94 16069 A (SCHERING CORP ; NAKAFUKU MASATO (JP); KAZIRO YOSHITO (JP)) 21 July 1994 (1994-07-21) abstract; claims 1-39	1-6, 9-15
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

24 March 2000

Date of mailing of the international search report

05. 07. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

Intern ial Application No

PCT/US 99/28013

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 91 15582 A (CETUS CORP) 17 October 1991 (1991-10-17) abstract; claims 1-46; example 10 ---	1-16,19, 20
A	WO 90 00607 A (CETUS CORP) 25 January 1990 (1990-01-25) abstract; claims 1-55; figures 3,4 -----	1-14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/28013

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 19,20 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 17 18 20
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
claims 1-20 partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 17 18 20

Claims 17,18,20 refer to an antagonist and agonist and the use of antagonist of polypeptide of claim 1 without giving a true technical characterization. Moreover, no such compound is defined in the application. In consequence, the scope of said claims is ambiguous and vague, and their subject-matter is not sufficiently disclosed and supported (art.5 and 6 PCT). No search can be carried out for such speculative claims the wording of which, is in fact, a mere recitation of the results to be achieved.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-20 (partially)

A protein with amino acid with seq.id. 1 and corresponding nucleotide sequence with seq.id. 30 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

2. Claims: 1-20 (partially)

A protein with amino acid with seq.id.2 and corresponding nucleotide sequence with seq.id. 31 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

3. Claims: 1-20 (partially)

A protein with amino acid with seq.id.3 and corresponding nucleotide sequence with seq.id. 32 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

4. Claims: 1-20 (partially)

A protein with amino acid with seq.id.4 and corresponding nucleotide sequence with seq.id. 33 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

5. Claims: 1-20 (partially)

A protein with amino acid with seq.id.5 and corresponding nucleotide sequence with seq.id. 34 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

6. Claims: 1-20 (partially)

A protein with amino acid with seq.id.6 and corresponding nucleotide sequence with seq.id. 35 , method for detecting a polynucleotide, expression vector ,host cell , method for

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

7. Claims: 1-20 (partially)

A protein with amino acid with seq.id.7 and corresponding
nucleotide sequence with seq.id. 36 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

8. Claims: 1-20 (partially)

A protein with amino acid with seq.id.8 and corresponding
nucleotide sequence with seq.id. 37 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

9. Claims: 1-20 (partially)

A protein with amino acid with seq.id.9 and corresponding
nucleotide sequence with seq.id. 38 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

10. Claims: 1-20 (partially)

A protein with amino acid with seq.id.10 and corresponding
nucleotide sequence with seq.id. 39 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

11. Claims: 1-20 (partially)

A protein with amino acid with seq.id.11 and corresponding
nucleotide sequence with seq.id. 40, method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

12. Claims: 1-20 (partially)

A protein with amino acid with seq.id.12 and corresponding nucleotide sequence with seq.id. 41 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

13. Claims: 1-20 (partially)

A protein with amino acid with seq.id.13 and corresponding nucleotide sequence with seq.id. 42 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

14. Claims: 1-20 (partially)

A protein with amino acid with seq.id.14 and corresponding nucleotide sequence with seq.id. 43 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

15. Claims: 1-20 (partially)

A protein with amino acid with seq.id.15 and corresponding nucleotide sequence with seq.id. 44 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

16. Claims: 1-20 (partially)

A protein with amino acid with seq.id.16 and corresponding nucleotide sequence with seq.id. 45 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

17. Claims: 1-20 (partially)

A protein with amino acid with seq.id.17 and corresponding nucleotide sequence with seq.id. 46 , method for detecting a polynucleotide, expression vector ,host cell , method for

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

18. Claims: 1-20 (partially)

A protein with amino acid with seq.id.18 and corresponding
nucleotide sequence with seq.id. 47 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

19. Claims: 1-20 (partially)

A protein with amino acid with seq.id.19 and corresponding
nucleotide sequence with seq.id. 48 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

20. Claims: 1-20 (partially)

A protein with amino acid with seq.id.20 and corresponding
nucleotide sequence with seq.id. 49 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

21. Claims: 1-20 (partially)

A protein with amino acid with seq.id.21 and corresponding
nucleotide sequence with seq.id. 50 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

22. Claims: 1-20 (partially)

A protein with amino acid with seq.id.22 and corresponding
nucleotide sequence with seq.id. 51 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

23. Claims: 1-20 (partially)

A protein with amino acid with seq.id.23 and corresponding nucleotide sequence with seq.id. 52 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

24. Claims: 1-20 (partially)

A protein with amino acid with seq.id.24 and corresponding nucleotide sequence with seq.id. 53 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

25. Claims: 1-20 (partially)

A protein with amino acid with seq.id.25 and corresponding nucleotide sequence with seq.id. 54 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

26. Claims: 1-20 (partially)

A protein with amino acid with seq.id.26 and corresponding nucleotide sequence with seq.id. 55 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

27. Claims: 1-20 (partially)

A protein with amino acid with seq.id.27 and corresponding nucleotide sequence with seq.id. 56 , method for detecting a polynucleotide, expression vector ,host cell , method for producing a polypeptide , pharmaceutical composition , antibody , agonist and antagonist , method for preventing a disorder

28. Claims: 1-20 (partially)

A protein with amino acid with seq.id.28 and corresponding nucleotide sequence with seq.id. 57 , method for detecting a polynucleotide, expression vector ,host cell , method for

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

29. Claims: 1-20 (partially)

A protein with amino acid with seq.id.29 and corresponding
nucleotide sequence with seq.id. 58 , method for detecting a
polynucleotide, expression vector ,host cell , method for
producing a polypeptide , pharmaceutical composition ,
antibody , agonist and antagonist , method for preventing a
disorder

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: 1st Application No

PCT/US 99/28013

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9837196 A	27-08-1998	AU 6661298 A EP 0981613 A	09-09-1998 01-03-2000
WO 9416069 A	21-07-1994	AU 6083894 A CA 2153486 A EP 0679185 A JP 8507204 T	15-08-1994 21-07-1994 02-11-1995 06-08-1996
WO 9115582 A	17-10-1991	AU 7554691 A EP 0537155 A	30-10-1991 21-04-1993
WO 9000607 A	25-01-1990	US 5104975 A AT 156518 T AU 627764 B AU 4034989 A DE 68928242 D DE 68928242 T EP 0466688 A EP 0649908 A US RE35171 E US 5234839 A US 5760203 A US 5763573 A US 5830684 A	14-04-1992 15-08-1997 03-09-1992 05-02-1990 11-09-1997 18-12-1997 22-01-1992 26-04-1995 05-03-1996 10-08-1993 02-06-1998 09-06-1998 03-11-1998

